

## The radioprotective effect of a wheat germ diet on rat myocardial tissue exposed to X-rays

### Efecto radioprotector de una dieta de germen de trigo en tejido miocárdico de ratas expuestas a rayos X

María Bayona-Caballero<sup>1</sup>, Judith Alayo-Zavaleta<sup>1</sup>, César Lombardi-Pérez<sup>2</sup>, Carmen Marín-Tello<sup>1</sup>

<sup>1</sup> Universidad Nacional de Trujillo. Facultad de Farmacia y Bioquímica. Trujillo Perú.

<sup>2</sup> Universidad Privada Antenor Orrego. Escuela de Medicina Veterinaria y Zootecnia. Trujillo Perú

<http://dx.doi.org/10.30827/ars.v60i4.10226>

#### Artículo Original Original Article

##### Correspondencia

Correspondence

Carmen Marín-Tello. Human Physiology and Physiopathology. Facultad de Farmacia y Bioquímica. Universidad Nacional de Trujillo. Perú. Ciudad Universitaria Av. Juan Pablo II S/N. Phone: 044-474839.  
E-mail: cmarin@unitru.edu.pe

##### Financiación

Fundings

No fundings received

##### Conflicto de interés

Competing interest

The authors report no conflict of interests

##### Agradecimientos

Acknowledgements

Received: 05.08.2019  
Accepted: 09.09.2019

#### RESUMEN

**Introducción:** Determinar si una dieta con germen de trigo (*Triticum aestivum*) tiene un efecto radioprotector sobre el tejido miocárdico de ratas albinas (*Rattus rattus var. Albinus*)

**Método:** Ratas con un peso entre 200 - 250 g fueron divididas en 4 grupos de 6 cada uno: Dos grupos fueron alimentadas con una dieta regular de germen de trigo durante 16 días antes y después de una única exposición a un grupo con 18 mSv de rayos X. Los otros dos grupos fueron alimentados con una dieta estándar, uno de ellos, fue expuesto a los rayos X. Los animales fueron sacrificados y se realizó un estudio histopatológico del tejido cardiaco.

**Resultados:** Las ratas alimentadas con una dieta estándar y expuesta a los rayos X, presentaron marcada hiperemia de vasos sanguíneos, necrosis, presencia de fibrocitos en el tejido conectivo, pérdida de la arquitectura y disposición radial muscular. Las ratas expuestas y alimentadas con dieta de trigo presentaron solo una ligera necrosis y presencia de fibrocitos. Las ratas no expuestas a rayos X presentaron un miocardio saludable.

**Conclusiones:** La dieta del germen de trigo puede tener un efecto radioprotector sobre el miocardio de rata.

**Palabras clave:** Trigo; Rayos X; Corazón.

#### ABSTRACT

**Introduction:** To determine whether a diet of wheat (*Triticum aestivum*) germ has a radioprotective effect on albino rat (*Rattus rattus var. Albinus*) myocardial tissue.

**Method:** Rats between 200 and 250 g were divided into 4 groups of 6 each: Two groups were fed either a wheat diet or regular diet 16 days before and after a single exposure to 18 mSv of X-rays. The other two groups were fed the same diets but not exposed to X-rays. The animals were sacrificed, and heart tissue was submitted to histopathological study.

**Results:** Rats fed a standard diet and exposed to X-rays presented marked hyperemia of blood vessels, necrosis, presence of connective tissue fibrocytes, loss of muscle architecture and radial arrangement. Exposed rats fed a wheat diet presented with only light necrosis and the presence of fibrocytes. Rats not exposed to X-rays had healthy myocardia.

**Conclusions:** Wheat germ diet may have a radioprotective effect on rat myocardium.

**Keywords:** Triticum; X-Rays; heart.

## INTRODUCTION

Ionizing radiation, such as X-rays, damages cells because it has sufficient energy to break chemical bonds in and cause damage to vital cellular components, causing cell death, cell repair, or cancer. This damage can be caused directly to cellular components or indirectly through the formation of reactive oxygen species from water. The severity of radiation damage depends on both absorbed dose size and frequency of exposure. A large dose may kill, but repeated small doses can increase cancer risk.

On average, a person receives a radiation dose of 3 mSv/year from both artificial and natural sources, with radon gas and medical procedures such as X-ray imaging and radiotherapy among the most important sources. Recently, the use of these imaging and therapeutic procedures has increased. Although this trend is helpful, it also makes radiation hygiene for both medical personnel and patients an increasingly important consideration. Radiation hygiene may include preparation of technical guides using the ALARA principle (As Low As Reasonably Achievable), education of medical staff and supporting a culture of radio-protection, using newer technologies that maintain quality of treatment but decrease dose, and performing risk-benefit analysis of procedures that involve ionizing radiation<sup>(1)</sup>.

Although radiation hygiene is helpful in limiting exposure, it will be difficult to eliminate all exposure. Of particular concern are parts of the body that may be exposed during common procedures. Centrally located organs like the heart would have the greatest exposure. For example, 40% of all X-ray images taken worldwide involve the thorax<sup>(2)</sup> giving the heart a dose each time. It also may receive collateral doses during radiotherapy treatments of thoracic tumors, which may have secondary effects. For instance, breast cancer radiotherapy carries the risk of cardiac death years later<sup>(3)</sup>. Although the heart is not considered highly sensitive to radiation, a dose of 40 Gy broken into a conventional radiotherapy regimen of 10 Gy/week can cause some myocardial degeneration. A dose greater than 60 Gy for the whole heart can cause death by pericardial effusion and constrictive pericarditis.

Given this potential for collateral damage in spite of good radiation hygiene, the possibility of improving the body's resistance to and ability to recover from radiation doses is appealing. One promising area of research is supplementation with radioprotective foods and substances before, during, and after radiation dose. One such food, whole wheat, has been found to have cardioprotective properties<sup>(4)</sup> most likely due to its antioxidant, nutritionally essential polyunsaturated fatty acids, and vitamin content. Found at

highest concentrations in the germ, these components may improve cellular repair, decrease inflammation<sup>(5)</sup>, improve electrical properties of the myocardium<sup>(6)</sup> and scavenge free radicals post-exposure<sup>(7,8,9,10)</sup>.

Although the impact of diet on the development of resistance to radiation damage is often overlooked<sup>(11)</sup>, some research using gastric, esophageal, and hepatic rat tissue<sup>(7,12)</sup>, is promising in that animals fed with wheat germ better maintain cellular morphology post exposure to X-rays. Given the frequent exposure of the heart to concomitant radiation during medical procedures, here we aim to extend previous research by evaluating the radioprotective effects of a wheat germ-enriched diet on cardiac tissue from rats exposed to X-rays.

## METHODS

### Animals and diet:

The study took place in the animal facility and toxicology laboratory of the School of Pharmacy and Biochemistry of the National University of Trujillo (UNT). Animals were treated in accordance with international guidelines for handling of experimental animals. The study was carried out using two diets - wheat germ (WG) or the standard animal facility diet of corn and barley, which does not contain wheat (NW). Germ from a native variety of wheat (*Triticum aestivum*) was obtained on the open market in Otuzco province, Peru and was confirmed by the *Herbarium Truxillensis* of UNT.

Male rats *Rattus rattus* var. Albinus between 200 and 250 g were obtained from the Animal Facility of the National Health Institute, Peru. On day 1, the rats were assigned to be in one of 4 groups of 6 animals each: NW diet, no radiation exposure (NWNE); WG diet, no radiation exposure (WGNE); NW diet, radiation exposure (NWE); WG diet, radiation exposure (WGE). All animals were housed in individual cages and received 10 g of food daily. Water was provided *ad libitum*. Tiles were placed in the cages so that the animals had access to all of the food provided.

### Radiation treatment of animals:

On day 16, rats of groups NWE and WGE were immobilized with pentobarbital in saline (3 mg/kg). The rats were exposed to 18 mSv of X-rays in a standard X-ray imaging device at an ES-SALUD hospital at El Porvenir district, La Libertad Region, Peru. The animals continued with their diet for an additional 16 days after exposure.

All animals were sacrificed with a dose of 12 mg/kg pentobarbital in saline on day 32. The heart was extracted and preserved in 10% formaldehyde.

### Histopathological Study:

Longitudinal and transversal slices were made from the extracted hearts and were stained with hematoxylin and eosin stain. A pathologist analyzed the slices for tissue characteristics, without knowing the identities of the samples.

### RESULTS

The presence of radiation-damaged cells was observed in samples exposed to X-rays (Fig. 1 C and D; Fig. 2 C and D). However, the severity of tissue damage is clearly different: qualitatively comparing Fig 1D and Fig 2C makes it clear that the NWE group suffered more cardiac damage (necrosis, hyperemia and karyolysis) when compared to the WGE group which had less damage (no karyolysis, reduced necrosis and hyperemia).

In contrast, cellular viability, indicated by the presence of fusiform nuclei and proper cellular architecture

was maintained in groups not exposed to radiation (compare Fig. 1 A and B to Fig. 2 A and B). Myocardocytes, myofibrils, and connective tissue cells and nuclei have normal morphology. Taken together, it seems that the WG and NW diets maintain cardiac tissue viability in the absence of ionizing radiation, but the WG diet decreases the severity of radiation-induced cardiac damage.

### DISCUSSION


The use of ionizing radiation plays an important role in the medical sciences, such as in diagnostic imaging and radiotherapy. Although useful, ionizing radiation can cause tissue damage<sup>(14)</sup> by directly damaging vital cellular apparatuses or by generating reactive oxygen species (ROS) or other free radicals. Damage to DNA from these reactive species induces apoptosis, which includes an ordered fragmentation of internucleosomal DNA strands and condensation of nuclear chromatin, which causes disappearance of the nucleus<sup>(15)</sup>. These effects were observed in rat myocardium (Fig. 1 C and D, Fig. 2 C and D) exposed to a single 18 mSv dose from an imaging X-ray machine.

Of the two groups exposed to X-rays, the group fed wheat germ (WGE) experienced less myocardial damage than the group fed the standard diet (NWE). A likely explanation is that a diet of wheat germ provides some protection from the quantity of radiation-induced damage (Fig. 1 D and Fig. 2 C) in myocardium. A probable mechanism for this protection may be that wheat germ components, such as vitamins, organic acids, sterols, lipids, glutamic acid, proteins, antioxidants, and others<sup>(16,17,18,19)</sup> may protect against ionizing radiation by directly absorbing it or by neutralizing or inhibiting the formation of free radicals<sup>(20,21)</sup> as compared


to the standard diet. This would leave fewer free radicals available to damage DNA. Wheat germ components may also stimulate cellular repair mechanisms. Additionally, antioxidants in wheat germ may be beneficial because they improve cardiac contraction through improved regulation of the flow of calcium ions<sup>(13)</sup> maintaining myocardium during and after radiation exposure. Wheat germ contains lipids and sterols that maintain cell membranes that can also attenuate the effect of radiation damage, especially capillary membranes. This damage can initiate inflammatory processes that ultimately lead to heart disease<sup>(22)</sup>.

Although medical procedures involving ionizing radiation procedures have become safer due to improvements of radiation hygiene protocols, it is not possible to eliminate collateral exposure to ionizing radiation. Exposure of this type is of particular concern for organs like the heart because its central location frequently puts it in risk of exposure to ionizing radiation from medical imaging and radiotherapy of the thorax. Therefore, increasing the resistance of heart to radiation damage, such as through dietary supplementation<sup>(23,24)</sup> is an appealing approach. The evidence that a wheat germ diet in rat can be radioprotective provides promise for additional dietary protocols to help build resistance to radiation damage.

Future studies that could complement and extend these results could include testing for biological markers of cell damage and cytogenetic assays<sup>(25)</sup> among others.



**Figure 1. Representative micrographs of myocardium of rats fed the NW diet. (A)** Micrograph of group NWNE myocardium at 400x. 1) Nucleus of cardiac muscular cells. 2) Fusiform nucleus of connective tissue cells. 3) Areas free of fibrils. **(B)** Micrograph of group NWNE 100x showing branching and anastomosis of myocardium and vascularized connective tissue. 1) Fusiform nuclei of capillary endothelium. 2) Nucleus of cardiac muscular cells. **(C)** Micrograph of the ventricular section of group NWE myocardium at 100 x showing marked hyperemia in blood vessels h) and loss of muscular architecture (\*) with an absence of radial ordering. Ventricular section. **(D)** Micrograph of the ventricular section of group NWE myocardium at 600 x, indicating the presence of anuclear cardiac cells, nuclei undergoing karyolysis cc), and f) Presence of fibrocytes in connective tissue. Ventricular section



**Figure 2. Representative micrographs of myocardium of rats fed the WG diet. (A) and (B)** Micrographs of group WGNE at 1000 x. **(A)** Marked staining of nuclei of cardiac muscle (m), fibrocytes (f), and endothelial cells (e) indicates tissue viability. **(B)** Image indicates apparently normal pathology with branching and anastomosis of myocardium. Both muscular and fibrocyte cell nuclei are fusiform. **(C)** Micrograph of group WGE transversal cut at 400 x. A few anuclear muscle fibers indicate radiation-caused necrosis (\*). Transversal cut **(D)** Micrograph of group WGE transversal cut at 1000 x. Presence of myocardial nuclei (\*) and fibrocytes (arrow) indicate continued viability of the tissue with minimal radiation damage. Transversal cut

**Study Limitation:** A sham treatment group was not used.

## REFERENCES

1. Soffia P, Ubada C, Miranda P, Rodríguez JL. Radioprotección al día en radiología diagnóstica: Conclusiones de la Conferencia Iberoamericana de Protección Radiológica en Medicina (CIPRaM) 2016. *Rev Chil Radiol.* 2017;23(1):15–19.
2. World Health Organization (WHO). Department of Public Health, Environmental and Social Determinants of Health (PHE). Comunicación sobre los riesgos de la Radiación en la imagenología Pediátrica. 2016. Available from: [http://www.who.int/ionizing\\_radiation/pub\\_meet/summary-es.pdf?ua=1](http://www.who.int/ionizing_radiation/pub_meet/summary-es.pdf?ua=1)
3. Taylor CW, McGale P, Povall JM, Thomas E, Kumar S, Dodwell D, et al. Estimating Cardiac Exposure From Breast Cancer Radiotherapy in Clinical Practice. *Int J Radiat Oncol.* 2009;73(4):1061–1068. doi: 10.1016/j.ijrobp.2008.05.066
4. Aune D, Keum NN, Giovannucci E, Fadnes LT, Boffetta P, Greenwood DC, et al. Whole grain consumption and risk of cardiovascular disease, cancer, and all cause and cause specific mortality: systematic review and dose-response meta-analysis. *BMJ.* 2016;353:g4398. doi: 10.1136/bmj.g4398

- ysis of prospective studies. *BMC Med.* 2016;14(1):1-14. doi: 10.1136/bmj.i2716
5. Valenzuela R, Tapia G, González M VA. Omega-3 fatty acids (EPA and DHA ) and its applications in diverse clinical situation. *Rev Chil Nutr.* 2011;38:356-367.
  6. Carrero J, Martín-Bautista E, Baró L, Fonollá J, Jiménez J, Boza J LE. Cardiovascular effects of omega-3-fatty acids and alternatives to increase their intake. *Nutr Hosp.* 2005;20(1):63-69. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/15762422>
  7. Hoyos M, Flores P. Tipos de Radiación, Aplicaciones, Beneficios y Riesgos. *Rev. Act. Clin. Med [edición electrónica]*. 2013 [cited 2019 Ago 04]. Available from: [http://www.revistasbolivianas.org.bo/scielo.php?script=sci\\_arttext&pid=S2304-37682013001000003&lng=es](http://www.revistasbolivianas.org.bo/scielo.php?script=sci_arttext&pid=S2304-37682013001000003&lng=es).
  8. Almén A, Lundh C, Bath M. Challenges assessing radiation risk in image-guided treatments - Implications on optimisation of radiological protection. *Journal of Radiological Protection.* 2018;38(3):1064-1076. doi: 10.1088/1361-6498/aacc83
  9. Fernandez-Gil B, Abdel Moneim AE, Ortiz F, Shen YQ, Soto-Mercado V, Mendivil-Perez M, et al. Melatonin protects rats from radiotherapy-induced small intestine toxicity. *PLoS One.* 2017;12(4):1-21. doi: 10.1371/journal.pone.0174474
  10. Fuentes L, Felipe S, Valencia V. Efectos biológicos de los Rayo-X en la práctica de Estomatología. *Rev Habanera Ciencias Médicas.* 2015;14(3):337-347.
  11. Pérez-Ruíz JM. Impact of nutritional assessment in patients with heart failure. *Nutr Hosp.* 2017;34(6):1265-1266. doi: 10.20960/nh.1677
  12. Marín-Tello C, Guevara-Vásquez AM, Mejía D, Sánchez C, Lombardi-Pérez C. Efecto de *Triticum aestivum* (trigo) sobre la arquitectura de los hepatocitos de *Rattus rattus* var. albinus irradiados con Rayos X. *Pharmaciencia.* 2013;(1):16-23.
  13. Rosero-Salazar D, Ortiz-Salazar M S-ML. Miocardiocitos conductentes ventriculares. *Univ y Salud.* 2015;17(2):262-70.
  14. Ramírez ME, Rojas M. La necrosis, un mecanismo regulado de muerte celular. *Iatreia.* 2010;23(2):166-177.
  15. Bialostozky D, Rodríguez-Diez G ZC. Apoptosis detection in cardiovascular diseases through nuclear cardiology spect images. *Arch Cardiol Mex.* 2008;78(2):217-28.
  16. Fuchs-Tarlovsky V, Bejarano-Rosales M, Gutiérrez-Salmeán G, Casillas M<sup>3</sup>, López-Alvarenga J, Ceballos-Reyes G. M. Effect of antioxidant supplementation over oxidative stress and quality of life in cervical cancer. *Nutr. Hosp.* 2011;26(4):819-826. doi: 10.3305/nh.2011.26.4.4894
  17. Abilés J, Moreno-Torres R, Moratalla G, Castaño J, Abúd RP, Mudarra A, et al. Effects of supply with glutamine on antioxidant system and lipid peroxidation in patients with parenteral nutrition. *Nutr. Hosp.* 2008;23(4):332-339.
  18. Fischer N, Seo E-J, Efferth T. Prevention from radiation damage by natural products. *Phytomedicine.* 2018;47:192-200. doi: 10.1016/j.phymed.2017.11.005
  19. Kim W, Kang J, Lee S, Youn B. Effects of traditional oriental medicines as anti-cytotoxic agents in radiotherapy. *Oncol Lett.* 2017;13(6):4593-4601. doi: 10.3892/ol.2017.6042
  20. González-Pumariega M, Fuentes-León F, Vernhes M, Schuch AP, Martins CF, Sánchez-Lamar Á. El extracto acuoso de *Cymbopogon citratus* protege al ADN plasmídico del daño inducido por radiación UVC. *Ars Pharm.* 2016; 57(4): 193-199
  21. Marín-Tello C, Matos-Deza L, Aliaga-Arauco J, Lombardi-Pérez C, Castañeda-Marín E, Rengifo-Penadillos R, et al. *Lepidium meyenii* (maca) and the cerebral stimulation for mobile phones: some answers in an animal model. 3rd International Brain Stimulation Conference. *Brain Stimul* 2019;12(2):555. doi: <https://doi.org/10.1016/j.brs.2018.12.835>
  22. Schultz-Hector S, Klaus-Rüdiger T. Radiation-induced cardiovascular diseases: Is the epidemiologic evidence compatible with the radiobiologic data? *Int. J. Radiation Oncology Biol. Phys.* 2007;67(1):10-18.
  23. Luna J, Amaya E, de Torres M.<sup>a</sup>V, Peña M.<sup>a</sup>C, Prieto I. Nutrientes y radioterapia; revisión de la literatura. *Nutr Hosp.* 2015;32(6):2446-2459. doi: 10.3305/nh.2015.32.6.9596
  24. MINISTERIO DE SALUD. "Tablas Peruanas de composición de los Alimentos." Lima Perú; 2017. 1-146 p.
  25. Castro I. Cytogenetic indicators for the identification of ionizing radiation exposure in humans. *Acta med costarric.* 2013;55(3):110-117.