

Revisión

Retinol, β -carotene, α -tocopherol and vitamin D status in European adolescents; regional differences and variability: A review

J. Valtueña¹, C. Breidenassel², J. Folle² and M. González-Gross^{1,2}

¹Department of Health and Human Performance. Faculty of Physical Activity and Sport Sciences (INEF). Universidad Politécnica de Madrid. Spain. ²Institut für Ernährungs- und Lebensmittelwissenschaften - Humanernährung. Rheinische Friedrich-Wilhelms Universität Bonn. Germany.

Abstract

Currently, blood levels to define vitamin deficiency or optimal status in adolescents are extrapolated from adults. This may be not adequate as vitamin requirements during adolescence depend on the process of sexual maturation, rapid increasing height and weight, among other factors. In order to establish the state of the art, Medline database (www.ncbi.nlm.nih.gov) was searched for studies published in Europe between 1981 and 2010 related to liposoluble vitamin status in adolescents. A comparison of the vitamin status published in the reviewed articles was difficult due to the lack of studies, lack of consensus on cut-off levels indicating deficiency and optimal vitamin levels and the different age-ranges used. In spite of that, deficiency prevalence varied for vitamin D (13-72%), vitamin A (3%), E (25%) and β -carotene (14-19%). Additional factors were considered as possible determinants. We conclude that it is necessary to establish a consensus on acceptable ranges and cut-offs of these vitamins during adolescence. Representative data are still missing; therefore, there is a high need to get deeper into the investigation on liposoluble vitamins in this population group.

(*Nutr Hosp.* 2011;26:280-288)

DOI:10.3305/nh.2011.26.2.4971

Key words: *Vitamins. Adolescence. Nutritional status. Risk factors.*

ESTADO DE RETINOL, β -CAROTENO, α -TOCOPHEROL Y VITAMINA D EN ADOLESCENTES EUROPEOS; DIFERENCIAS REGIONALES Y VARIABILIDAD: REVISIÓN

Resumen

En la actualidad, los diferentes valores sanguíneos que definen un estado óptimo o deficiente de vitaminas liposolubles en los adolescentes son extrapolados de los adultos. Sin embargo, podría no ser lo adecuado debido a que los requerimientos vitamínicos de los adolescentes están marcados por el proceso de maduración sexual y crecimiento entre otros factores. Para establecer el punto de partida, la base de datos Medline (www.ncbi.nlm.nih.gov) ha sido el medio utilizado para la búsqueda de los estudios publicados sobre el estado en vitaminas liposolubles en adolescentes europeos entre los años 1981 y 2010. Comparar los diferentes resultados obtenidos en los diferentes estudios fue difícil debido a la carencia de estudios, a la falta de consenso en los puntos de corte que indican deficiencia y estado óptimo y a los diferentes rangos de edad utilizados. A pesar de esto, en función de los estudios, se observa una variabilidad en la prevalencia de deficiencia de vitamina D (13-72%), vitamina A (3%), E (25%) y β -caroteno (14-19%). Otros factores adicionales fueron considerados como posibles determinantes del estado vitamínico. Se identifica la necesidad de establecer un consenso sobre los rangos aceptables y puntos de corte de estas vitaminas para este grupo de población y profundizar en la investigación de las vitaminas liposolubles en el periodo de la adolescencia.

(*Nutr Hosp.* 2011;26:280-288)

DOI:10.3305/nh.2011.26.2.4971

Palabras clave: *Vitamins. Adolescence. Estado nutricional. Factores de riesgo.*

Correspondence: Jara Valtueña Santamaría.
Departamento de Salud y Rendimiento Humano.
Facultad de Ciencias de la Actividad Física y del Deporte-INEF.
Universidad Politécnica de Madrid.
C/ Martín Fierro, 7.
28040 Madrid.
E-mail: jara.valtueña@upm.es

Recibido: 24-VI-2010.
1.ª Revisión: 14-IX-2010.
Aceptado: 17-IX-2010.

Abbreviations

BMI: Body Mass Index.

RBP: Retinol Binding Protein.

Introduction

Vitamin deficiency, at least on a subclinical level, could be an unrecognized health problem in children and adolescents in Europe because they are not routinely screened for in these population groups. Especially adolescents are considered as a risk group for malnutrition because of their increasing needs in nutrients and energy intake for an adequate growth and development varying with age.¹ Additionally to that, the risk for malnutrition is enhanced because although the access to high quality food and nutrition is continuously improving in European countries,² dietary patterns and food choices are not optimal and not according to recommended guidelines.¹

Vitamins are specifically involved in multiple cellular and tissue processes,³ and there is increasing evidence that links deficiencies with chronic diseases like cardiovascular and cerebrovascular disease, cancer, diabetes and osteoporosis in adulthood,¹⁻⁸ though data are scarce for younger ages. But deficiency stages at these early ages could contribute to risk factors,^{2,5,8-10} beside other already described implications like higher risk for osteomalacia,^{8,11,12} impaired cognitive function and concentration problems,^{13,14} hyperactivity⁹ and immune system impairment.^{8,13,15} For identifying deficiency stages, reference values are currently extrapolated from adult data for most vitamins.¹⁶

As already mentioned, little information exists on the vitamin status of representative populations in European countries, particularly for Eastern Europe. Lambert et al. published a review on the contribution of essential nutrients and nutritional status of European adolescents, but vitamins were barely mentioned.¹⁷ Complementary to that review, we published a review on B vitamin status in European adolescents some years ago.¹⁸ The aim of the present review is to get more information about the existing data to give a better overview on fat soluble vitamin (A, E, D and provitamin β -carotene) status in European adolescents, with a specific focus on blood levels, reference values and assessment methods.

Material and methods

Those studies which evaluated blood fat soluble vitamin levels in European adolescents were included in this review. The database Medline (www.ncbi.nlm.nih.gov) was used for searching studies published between 1981 and 2010. Terms like "European adolescents", "liposoluble or fat soluble vitamins", "vitamin status and values", "antioxidants", names of European coun-

tries and technical terms of the different liposoluble vitamins as well as combinations out of these terms were entered in the data base for the search. In addition, references of relevant articles were also used to get further information. Because the adolescence period goes from childhood to adulthood, age ranges vary in different articles and data for a broader age-range had to be included (from 9 to 18 years).

Results

A total of 17 articles including around 4503 subjects from different European countries were found focused on the assessment of liposoluble vitamins in adolescents, from North (Finland, Sweden, Ireland, Poland and Denmark), East (Yugoslavia, Austria and Hungary), West (France and Great Britain) and South (Italy and Greece) Europe. Analysing the number of subjects, only three studies included a sample size of less than 50 persons. Two studies analysed a population between 50-100, seven between 100 and 500 and four studies between 500 and 1,000 subjects (table I).

Vitamin D status in European adolescents

The main sources of vitamin D are subcutaneous skin synthesis under the influence of ultraviolet light (290-315 nm) and food.¹⁹ Vitamin D deficiency has been identified as a common problem in Europe.¹⁹⁻²¹ Due to the geographical situation of our continent, located at high latitude of 40° N in Madrid (Spain) and 60° N in Oslo (Norway), the skin synthesis of vitamin D may not compensate a low nutritional intake. Moreover, vitamin D intake in Europe is in general low (\leq 2-3 mg/d)²¹. In all the reviewed studies vitamin D status was assessed by means of serum 25-hydroxycholecalciferol [25(OH)D], which is the main circulating vitamin D metabolite, representing not only the consumed amount through diet and supplements but also the subcutaneous synthesis.^{19,22,23} One of the most important applications of vitamin D assessment in adolescence is the determination of abnormal serum concentrations of calcium for bone health. Moderately low levels of vitamin D may have adverse effects on calcium homeostasis leading to bone loss, even without symptoms of osteomalacia.^{19,24,28} Some studies report a positive association between serum [25(OH)D] and bone mineral content in adolescents.^{25,26} Inadequate vitamin D levels have been related to other diseases such as diabetes, multiple sclerosis and cancer.^{27,28}

Even vitamin D reference values for deficiency have been set at 27.5 nmol/L in children,¹⁶ there is still a lack of consensus on the deficient and sufficient 25(OH)D concentration, that varies from 20 nmol/L to 100 nmol/L in studies. A proposed scale for the adult population defined hypovitaminosis D as 25(OH)D concentrations below 100 nmol/L, insufficient vitamin D at

Table I
Studies on the nutritional status of fat-soluble vitamins in adolescents

	Country	Year	Author	Subjects		Vitamins	Additional parameters	
				Age (years)	Gender			
North	Finland	1997	Lehtonen-Veromaa et al. (1999) ³⁰	9-15	♀	D		
	Sweden	N.D.	Drott et al. (1993) ⁵¹	12-13, 13-14,	♂,♀	A, E	RBP	
				14-15, 15-16				
Ireland, Finland, Poland, Denmark	2002/03	Andersen R et al. (2005) ³⁶		12-13	♀	D		
East	Yugoslavia	N.D.	Buzina et al. (1982) ³⁰	12-15		A		
	Slovaquia	N.D.	Krajcovicova et al. (1997) ⁵⁴	11-14	♂,♀	A, E	β-C	
				13-16	♂,♀	A, E	β-C	
	Austria	1991	König y Elmadafa (2000) ³⁵ , ASNS		10-12, 13-14, 15-19	♂,♀	D	
	Austria	1991	Elmadafa et al. (1994) ⁵ , ASNS		10-12, 13-14, 15-18	♂,♀	A, E, K	
	Austria	N.D.	Marktl et al. (1982) ⁵²		11-12	♂,♀	A, E	
	France	N.D.	Guillemant et al. (2001) ⁴		13-16	♂	D	
France	1988	Herberg et al. (1994) ³⁰		10-14, 14-18	♂,♀	A, E	β-C, VitE/Chol	
France	1988/89	Herbeth et al. (1991) ⁴³		10-15	♂,♀	A, E	β-C	
France	1985/6	Malvy et al. (1993) ³⁶		12-13, 14-16	♂,♀	A, E	RBC; β-C, VitE/Chol	
France	1985/86	Malvy et al. (1989) ⁴⁷		9-12, 12-14, 14-16	♂,♀	A, E	β-C, VitE/Chol	
Great Britain	1997	Gregory et al. (2000) ³⁷		11-14, 15-18	♂,♀	D, A, E	β-C, VitE/Chol	
Greece	N.D.	Hassapidou et al. (1996) ⁵⁵		13-14	♂,♀	E		
Italy	N.D.	Bonofiglio et al. (1999) ³⁸		11-16	♀	D		

N.D.: Not documented; β-C: β-carotene; ASNS: Austrian Study on Nutritional Status of Schoolchildren.

concentrations of 25(OH)D below 50 nmol/L and a vitamin D deficiency when values are below 25 nmol/L.²⁹ The German, Swiss and Austrian Nutrition societies (D-A-CH) also agree to the latter data, that serum 25(OH)D levels between 20 to 25 nmol/L present a suboptimal state.^{16,30} There is a consensus that serum 25(OH)D concentrations below 10 nmol/L lead to risk of rickets, osteomalacia, increase in cardiovascular and cancer risk factors among others.^{8,24,31-34}

Several factors can influence vitamin D status. Regional differences can be appreciated throughout Europe (table II). The highest values for vitamin D were found in French teenagers,⁴ the lowest values were found in an Austrian study.³⁵ Depending on the cut offs used (varying from 15 nmol/L to 25 nmol/l), vitamin D deficiency ranged from 13 to 72% due to geographical differences.

Seasonal differences have been studied by several authors.^{4,20,36,37} In both French and Austrian children significant differences in serum concentrations between summer and winter were found, showing higher concentrations in summer.^{4,37} The prevalence of suboptimal vitamin D status was 5% lower in summer due to more adequate sun exposure in Finnish children.²⁰

The influence of age and Tanner stage on vitamin D status was analyzed in several studies with a significant decrease in concentrations of 25(OH)D levels with increasing age in both sexes ($p < 0.05$),^{35,37} although in British children aged 15 and above some slight increases were observed. Contrary to that, Bonofiglio et al. found a significant higher concentration of

25(OH)D in Italian postmenarcheal girls compared to premenarcheal ones ($p < 0.01$).³⁸ This could be due to the higher concentrations of vitamin D binding protein caused by the increase of estrogens levels.

The influence of body composition on vitamin D levels was studied by Smotkin-Tangorra et al.,³⁹ who found an inverse correlation between 25(OH)D levels and body mass index (BMI). Regarding supplementation, Guillemant et al.⁴ and Lehtonen-Veromaa et al.²⁰ observed a significant effect on 25(OH)D serum levels ($p = 0.000$) in French and Finnish children (table II).

A, E and provitamin β -carotene vitamin status in European adolescents

Vitamin E and the provitamin β -carotene have an important role as antioxidants in the human body.⁴⁰ Current lifestyle of adolescents with a low intake of fruits and vegetables^{17,41} could create a pro-oxidant situation with serious long-term consequences. Also vitamin A during adolescence is critical due to its role in cell growth, vision and tissue development. High oxidative stress and free radical levels are involved in the development of cardiovascular disease and other undesirable metabolic situations in adulthood,^{40,41} but risk factors are already established during childhood and adolescence.⁴²

Little is known about the reference blood levels of these vitamins during adolescence.⁴³ In adults, the World Health Organization (WHO) defines a defi-

Table II
Vitamin D status

Age (years)	9	10	11	12	13	14	15	16	17	18	19
Authors											
Lehtonen-Veromaa et al. ²⁰ Finland				♀ (1) ¹ 33.9 ± 13.9 nmol/l ♀ (2) ¹ 62.9 ± 15.0 nmol/l (summer) ♀ (3) ¹ 33.7 ± 11.4 nmol/l							
Guillemant et al. ⁴ France					♂ (1) ¹ 71.6 ± 19.9 nmol/l (summer) ♂ (2) ¹ 20.4 ± 6.9 nmol/l (winter) ♂ (3) ¹ 52.4 ± 16.3 nmol/l (summer) ♂ (4) ¹ 21.4 ± 6.1 nmol/l (winter)						
König y Elmadfa ³⁵ Austria		9.9 ± 6.1 nmol/l			9.2 ± 6.9 nmol/l		11.2 ± 8.1 nmol/l				
Gregory et al. ³⁷ Great Britain			♂ 56.7 ± 24.05 nmol/l ♀ 54.4 ± 23.52 nmol/l			♂ 56.7 ± 24.05 nmol/l ♀ 54.4 ± 23.52 nmol/l					
Bonofiglio et al. ³⁸ Italy				♀ 17.9 ± 1.1 ng/ml ² ♀ 44.75 ± 2.75 nmol/l ²⁺³ ♀ 23.2 ± 0.7 ng/ml ⁴ ♀ 58 ± 1.75 nmol/l ⁴⁺³							
Andersen R et al. ³⁶ North of Europe				♀ 29.4 (20.3, 38.3) nmol/l							

Serum concentrations of 25(OH)D Mean ± SD o Mean (interquartils).

¹blood extraction number. ²premenarcheal group. ³1 nmol/l = 0.4 ng/ml. ⁴postmenarcheal group.

ciency of vitamin A through plasma retinol values.⁴⁴ Concentrations lower than 10 µg/dL (0.35 µmol/L) indicate a deficiency and values between 10 and 20 µg/dL (0.7 µmol/L) are referred to as incipient deficiency.⁴⁵

Beta-carotene plasma levels are considered as a good indicator for fruit and vegetable consumption, with great importance because of its antioxidant power. Low levels are correlating with a higher risk of chronic diseases.⁴⁶ Although plasma values above 0.3 µmol/L are considered as acceptable in adults, β-carotene plasma levels above 0.5 µmol/L seem to have the best preventing effects.⁴⁵ To determine the nutritional status of vitamin E the level of serum α-tocopherol is normally used.⁴⁵ A strong correlation between total blood fat (triglycerides), cholesterol and α-tocopherol concentration in plasma has been found.⁴⁷

According to D-A-CH the normal α-tocopherol serum concentration for adults is between 12 and 46 µmol/L,³⁰ but for adolescents there are no explicit values. Although the limit value for an adequate nutri-

tional status is defined between 10-16 µmol/L,⁴⁸ the guide value for primary prevention of cardiovascular disease and cancer is higher than 30 µmol/L.³⁰

Data on vitamin A, E and β-carotene serum levels from existing European studies are presented in tables III-V. Deficiency states vary between less than 3% for vitamin A,^{9,15,49-50} 25% for β-carotene and 14-19% for α-tocopherol.^{9,15,50}

Gender seems to have no influence on serum blood levels. Regarding age, differences have been observed. In several studies, vitamin A and Retinol Binding Protein (RBP) levels increased according to age until puberty, where they reached adult levels.^{37,47,50,51} Plasma β-carotene and α-tocopherol levels seemed to be more stable through the age span in French children and adolescents.^{47,50,51} In British children, β-carotene levels increased with age in both sexes (p < 0.01).³⁷ And Winkelhofer-Roob et al.⁴¹ found that age was a significant predictor of plasma α-tocopherol concentration in Swiss subjects aged 0.4 to 38.7 years. Assessing vita-

Table III
Vitamin A status

Age (years)	9	10	11	12	13	14	15	16	17	18
Authors										
Drott et al. ⁵¹ Sweden				429 µg/l (211-599) ¹ 1.5 µmol/l (0.74-2.10) ⁴	924 µg/l (255-1,463) ¹ 3.23 µmol/l (0.89-5.12) ⁴	670 µg/l (343-1,112) ¹ 2.34 µmol/l (1.20-3.90) ⁴	576 µg/l (410-805) ¹ 2.01 µmol/l (1.43-2.81) ⁴			
Buzina et al. ⁵⁰ Yugoslavia				♂: 34.5 ± 8.8 µg/dl ² 1.21 ± 0.31 µmol/l ⁴						
Krajcovicova et al. ⁵⁴ Slovaquia				Vegetarians: 1.91 ± 0.02 µmol/l ² Omnivores: 1.54 ± 0.07 µmol/l ²						
Molnar et al. ⁴⁰ Hungary							Normal-weighted: 1.87 ± 0.51 µmol/l ²			
							Obese: 1.8 ± 0.51 µmol/l ²			
Hercberg et al. ⁵⁰ France				♂: 1.32 µmol/l (0.73/1.98) ³ ♀: 1.37 µmol/l (0.95/1.89) ³		♂: 1.56 µmol/l (1.16/2.30) ³ ♀: 1.6 µmol/l (1.27/2.59) ³				
Malvy et al. ⁴⁷ France		40.3 ± 9.4 µg/dl ² 1.41 ± 0.33 µmol/l ⁴		43.7 ± 9.8 µg/dl ² 1.53 ± 0.34 µmol/l ⁴		51.7 ± 12.2 µg/dl ² 1.81 ± 0.43 µmol/l ⁴				
Malvy et al. ⁵⁶ France				♂: 44.30 µg/dl (28.4-66.4) ¹ 1.55 µmol/l (1.0-2.32) ⁴ ♀: 42.15 µg/dl (21.8-64.0) ¹ 1.47 µmol/l (0.76-2.24) ⁴		♂: 51.50 µg/dl (25.4-88.7) ¹ 1.80 µmol/l (0.9-3.10) ⁴ ♀: 45.90 µg/dl (33.1-83.2) ¹ 1.60 µmol/l (1.16-2.90) ⁴				
Elmadfa et al. ¹⁵ Austria		31.6 ± 7.2 µg/dl ² 1.10 ± 0.25 µmol/l ⁴		33.5 ± 6.3 µg/dl ² 1.17 ± 0.22 µmol/l ⁴		38.6 ± 10.2 µg/dl ² 1.35 ± 0.36 µmol/l ⁴				
Marktl et al. ⁵³ Austria				♂: 156 ± 27.9 UI/dl ² 1.62 ± 0.30 µmol/l ⁴⁺⁵ ♀: 158 ± 26.3 UI/dl ² 1.66 ± 0.27 µmol/l ⁴⁺⁵						
Gregory et al. ³⁷ Great Britain				♂: 1.27 ± 0.27 µmol/l ² ♀: 1.37 ± 0.285 µmol/l ²			♂: 1.53 ± 0.392 µmol/l ² ♀: 1.51 ± 0.389 µmol/l ²			

Retinol Plasmatic concentrations.

¹Median (Range). ²Mean ± SD. ³Mean (Percentil 5/Percentil 95). ⁴1 µmol/l = 28.6 µg/dl. ⁵1 UI = 0.3 g.

Table IV
β-carotene status

Age (years)	9	10	11	12	13	14	15	16	17	18
<i>Authors</i>										
<i>Krajcovicova et al.</i> ⁵⁴ <i>Slovaquia</i>			Vegetarian: 0.46 ± 0.03 μmol/l ² Omnivores: 0.26 ± 0.02 μmol/l ²							
<i>Molnar et al.</i> ⁴⁰ <i>Hungary</i>							Normal-weighted: 0.4 ± 0.304 μmol/l ²			
							Obese: 0.26 ± 0.112 μmol/l ²			
<i>Hercberg et al.</i> ⁵⁰ <i>France</i>			♂: 0.64 μmol/l (0.22/1.79) ³ ♀: 0.58 μmol/l (0.17/1.44) ³			♂: 0.60 μmol/l (0.25/1.85) ³ ♀: 0.60 μmol/l (0.18/2.65) ³				
<i>Malvy et al.</i> ⁴⁷ <i>France</i>		499 ± 276 μg/l ² 0.93 ± 0.46 μmol/l ⁴		626 ± 388 μg/l ² 1.17 ± 0.72 μmol/l ⁴		525 ± 304 μg/l ² 0.98 ± 0.57 μmol/l ⁴				
<i>Malvy et al.</i> ⁵⁶ <i>France</i>			♂: 480 μg/l (135-1,250) ¹ 0.89 μmol/l (0.25-2.33) ¹ ♀: 527 μg/l (153-1,405) ¹ 0.98 μmol/l (0.29-2.62) ¹			♂: 487 μg/l (335-1,465) ¹ 0.91 μmol/l (0.62-2.73) ¹ ♀: 442 μg/l (210-1,310) ¹ 0.82 μmol/l (0.39-2.44) ¹				
<i>Elmadfa et al.</i> ¹⁵ <i>Austria</i>			33.2 ± 18.6 μg/dl ² 0.62 ± 0.35 μmol/l ⁴		33.7 ± 20.8 μg/dl ² 0.63 ± 0.39 μmol/l ⁴		40.8 ± 29.6 μg/dl ² 0.76 ± 0.55 μmol/l ⁴			
<i>Marktl et al.</i> ⁵² <i>Austria</i>			♂: 32.2 ± 21.9 μg/dl ² 0.60 ± 0.41 μmol/l ⁴ ♀: 25.3 ± 21.6 μg/dl ² 0.47 ± 0.40 μmol/l ⁴							
<i>Gregory et al.</i> ³⁷ <i>Great Britain</i>			♂: 0.297 ± 0.2108 μmol/l ² ♀: 0.272 ± 0.2370 μmol/l ²				♂: 0.249 ± 0.1781 μmol/l ² ♀: 0.257 ± 0.1580 μmol/l ²			

β-carotene plasmatic concentrations.

¹Median (Range). ²Mean ± SD. ³Mean (Percentil 5/Percentil 95). ⁴1 μmol/l = 536 μg/l.

mins by pubertal development, a positive correlation was found with vitamin A ($p < 0.001$), and a negative correlation with β-carotene and α-tocopherol serum levels with increasing Tanner stage ($p < 0.05$).⁴³

Differences in vitamin status have also been observed in relationship to BMI. In the study by Marktl et al.,⁵² obese adolescents presented higher vitamin A levels than normal-weighted. In the study by Molnar et al.⁴⁰, over-weighted adolescents with metabolic syndrome presented lower values of α-tocopherol and β-carotene than normal weighted ones ($p < 0.01$). However, in over-weighted adolescents without metabolic syndrome, only β-carotene concentrations were significantly lower compared to normal-weighted ones.

The effect of smoking on antioxidant status was studied in Britain by Crawley et al.⁵³ showing that teenage smokers had a lower intake of antioxidant nutrients, fruits, vegetables and cereals, being particularly significant among girls, but no blood levels were reported. An opposite effect was found among vegetarian subjects that presented higher levels of vitamin A, α-tocopherol and β-carotene than omnivore ones ($p < 0.001$).⁵⁴

In the same way as for vitamin D, seasonal differences have been observed in plasma α-tocopherol, cholesterol and retinol concentrations higher in winter

than in the other seasons⁴¹. In regard to regional differences increased α-tocopherol values were observed in South Europe.⁵⁵

Relationship between the different vitamins has also been studied. Serum retinol had a positive correlation with β-carotene and α-tocopherol levels^{37,47,56} and the latter with cholesterol.^{43,47,56} Drott et al. on the contrary found an inverse relationship between vitamin A and vitamin E serum concentrations.⁵¹

The influence of socioeconomic status has been studied by Marktl et al.⁵². Higher β-carotene and vitamin E serum concentrations were found in children of central Europe whose parents had a high socioeconomic background.

General discussion

There are very few published studies on nutritional vitamin status in European adolescents and data are not available for all countries, especially in the eastern part of Europe. Moreover, the percentage of representative studies is low and in some countries the sample is not representative at all of the general population. A comparison of the vitamin status published in the reviewed articles is difficult due to the lack of consensus on cut-

Table V
α-tocopherol status

Age (years)	9	10	11	12	13	14	15	16	17	18
Authors										
<i>Drott et al.</i> ⁵¹ Sweden				9.1 mg/l (3.2-13.2) ¹ 21.16 μmol/l (7.4-30.7) ⁴	6.1 mg/l (2.9-9.7) ¹ 14.9 μmol/l (6.7-22.5) ⁴	8.2 mg/l (3.0-13.2) ¹ 19.07 μmol/l (7.0-30.7) ⁴	7.5 mg/l (5.3-10.2) ¹ 17.44 μmol/l (12.3-23.7) ⁴			
<i>Krajcovicova et al.</i> ⁵⁴ Slovaquia				Vegetarians: 29.17 ± 0.96 μmol/l ² Omnivores: 22.94 ± 0.62 μmol/l ²						
<i>Molnar et al.</i> ⁴⁰ Hungary							Normal-weighted: 29.17 ± 0.96 μmol/l ²			
				Obese: 17.1 ± 4.65 μmol/l ²						
<i>Hercberg et al.</i> ⁵⁰ France				♂: 18.3 μmol/l (13.4/24.1) ³ ♀: 20.5 μmol/l (14.7/25.9) ³			♂: 18.3 μmol/l (13.4/24.1) ³ ♀: 20.5 μmol/l (14.7/25.9) ³			
<i>Malvy et al.</i> ⁴⁷ France		10.0 ± 2.0 mg/l ² 23.25 ± 4.65 μmol/l ⁴		9.7 ± 2.5 mg/l ² 22.65 ± 5.81 μmol/l ⁴		10.0 ± 2.1 mg/l ² 23.72 ± 4.88 μmol/l ⁴				
<i>Malvy et al.</i> ⁵⁶ France				♂: 8.30 mg/l (6.8-13.1) ¹ 19.30 μmol/l (15.81-30.46) ⁴ ♀: 8.9 mg/l (1.1-12.2) ¹ 20.7 μmol/l (2.56-28.37) ⁴		♂: 9.10 mg/l (5.8-15.7) ¹ 21.16 μmol/l (13.49-36.51) ⁴ ♀: 9.40 mg/l (3.7-13.7) ¹ 21.86 μmol/l (8.6-31.86) ⁴				
<i>Elmadfa et al.</i> ¹⁵ Austria		0.88 ± 0.21 mg/dl ² 20.46 ± 4.88 μmol/l ⁴			0.85 ± 0.19 mg/dl ² 19.77 ± 18.37 μmol/l ⁴		0.89 ± 0.20 mg/dl ² 20.7 ± 4.65 μmol/l ⁴			
<i>Marktl et al.</i> ⁵² Austria				♂: 1.17 ± 0.23 mg/dl ² 27.21 ± 5.35 μmol/l ⁴ ♀: 1.07 ± 0.22 mg/dl ² 24.88 ± 5.12 μmol/l ⁴						
<i>Gregory et al.</i> ²⁷ Great Britain				♂: 19.7 ± 4.02 μmol/l ² ♀: 20.5 ± 3.48 μmol/l ²			♂: 19.2 ± 3.97 μmol/l ² ♀: 19.8 ± 3.97 μmol/l ²			
<i>Hassapidpou et al.</i> ⁵⁵ Greece					0.79 mg/l (0.7-1.02) ¹ 18.37 μmol/l (16.28-23.27) ⁴					

α-tocopherol serum concentrations.

¹Median (Range). ²Mean ± SD. ³Mean (Percentil 5/Percentil 95). ⁴1 μmol/l = 0.43 mg/l.

off levels indicating deficiency and optimal vitamin levels and the different age-ranges used. Also, the way of presenting the results is not uniform, using the mean ± standard deviation (SD) or the largest median plus 5 and 95 percentiles. Its variation is also due to the variety of laboratory methods, lack of standard reference preparations and calibration materials, and shortages of these issues.³²⁻³⁵

Regarding the existing data on liposoluble vitamin status of the European adolescents reviewed, a general hypovitaminosis problem was found for vitamin D and β-carotene in both gender. However, one study carried out in the US by Yetley et al. concluded that males have higher vitamin D intakes and 25(OH)D concentrations than females.⁵⁷

There seem to be several factors that influence liposoluble vitamin status; age, season, BMI, smoking or socioeconomic status should be taken into account in future vitamin studies as possible modifiers of

vitamin status in adolescents. Vitamin D decreases according to age. In the before mentioned US study, hypovitaminosis D (< 27.5 nmol/L) was observed in 1% of infants and children aged < 11 years, 5% of adolescents aged 12-19 years, and 6% of adults aged < 20 years.⁵⁸ Serum α-tocopherol was found to increase slightly with age.⁵⁸

Several studies agree that vitamin D status exhibits a seasonal variation with higher values during the summer period.^{4,20,25,28} Both serum 25(OH) D concentration^{10,59,60} and biomarkers of bone turnover,⁶¹ positively correlate with bone mineral content. PTH concentrations tend to increase during winter⁶² being associated with bone absorption,^{11,12} that is related with bone mineral content.

Current research also suggests that obese adolescents present states of malnutrition.⁶³ This means that, despite an excess of body fat, certain nutrients may be at risk.²⁷ An inverse relation between antioxidant vita-

mins and vitamin D with BMI³⁹ has been observed.⁶⁴

Regarding health costs, Grant et al. suggested for Western countries of Europe that the increase in the mean serum 25(OH)D levels up to 40 ng/mL would have a positive impact on the reduction of direct and indirect economic burden of disease. For 2007, the reduction was estimated at € 187,000 million/year.⁶⁵

Due to the low 25(OH)D concentrations, especially during winter months, some researchers emphasize to improve vitamin D intake through supplements.⁶⁶ The fortification of food with vitamin D is not common in Europe, without any regulation in some countries, with the exception of some Scandinavian countries. Vitamin D is considered a pharmacological agent for children when the dose is higher than the levels recommended by the Nordic countries of 5 mg/d (200 IU/d) and till now the data about the influence of vitamin D supplementation or fortified foods in children and adolescents is scarce.⁶⁷ On the contrary, seasonal vitamin A intake is not so important. The recommendations in some countries like Ireland, Italy and the United Kingdom are lower than the average for all age ranges in men and women, without any significant deficiencies in these countries respect those with higher recommendations (DACH countries and Romania).

Also other behavioural and health aspects should be taken into account in future studies regarding vitamin status in adolescents. The lower intakes of antioxidant nutrients, fruits, vegetables and cereals by the teenage smokers in comparison with non smokers already mentioned agree with those found by Jain et al.⁶⁸ who observed an increased oxidative stress and a compromised antioxidant defence system in smokers.⁶⁸ Lenders et al.⁶⁴ found that concentrations of 25(OH)D were directly correlated with physical activity levels ($p < 0.05$).

Conclusions

Currently, there are no reference values for blood vitamin levels in adolescents. Analysing the available existing data on nutritional status in European adolescents, the comparability of them implies many difficulties due to the lack of data and reference values as well as the lack of consensus on methodological approaches. The importance of transnational studies in order to standardize the procedures and methodologies and the way to express the results to enable comparison should be highlighted.

References

1. Prentice A, Branca F, Decsi T, Michaelsen KF, Fletcher RJ, Guesry P et al. Energy and nutrient dietary reference values for children in Europe: methodological approaches and current nutritional recommendations. *Br J Nutr* 2004; 92 (2 Suppl.): 83S-146S.
2. Rolland-Cachera MF, Bellisle F, Deheeger M. Nutritional status and food intake in adolescents living in Western Europe. *Eur J Clin Nutr* 2000; 54 (1 Suppl.): 41S-46S.
3. Koletzko B, De la Guéronnière V, Toschke AM, von Kries R. Nutrition in children and adolescents in Europe: what is the scientific basis? Introduction. *Br J Nutr* 2004; 92 (2 Suppl.): 67S-73S.
4. Guillemand J, Le HT, Maria A, Allemandou A, Pérès G, Guillemand S. Wintertime vitamin D deficiency in male adolescents: effect on parathyroid function and response to vitamin D3 supplements. *Osteoporos Int* 2001; 12 (10): 875-9.
5. Pilz S, Dobnig H, Winklhofer-Roob B, Riedmüller G, Fischer JE, Seelhorst U, Wellnitz B et al. Low Serum Levels of 25-Hydroxyvitamin D Predict Fatal Cancer in Patients Referred to Coronary Angiography. *Cancer Epidemiol Biomarkers Prev* 2008; 17 (5): 1228-33.
6. Cheng S, Massaro JM, Fox C.S, Larson MG, Keyes MJ, McCabe EL et al. Adiposity, Cardiometabolic Risk, and Vitamin D Status: The Framingham Heart Study. *Diabetes* 2010; 59 (1): 242-8.
7. Holick MF. The Vitamin D Epidemic and its Health Consequences. *J Nutr* 2005; 135 (11): 2739S-48S.
8. Bischoff-Ferrari H. Health effects of vitamin D. *Dermatol Ther* 2010; 23 (1): 23-30.
9. Elmadfa I, Godina-Zarfl B, Dichtl M, König JS. The Austrian Study on Nutritional Status of 6- to 18-year-old pupils. *Bibl Nutr Dieta* 1994; (51): 62-7.
10. Scharla SH, Scheidt-Nave C, Leidig G, Woitge H, Wüster C, Seibel MJ et al. Lower serum 25-hydroxyvitamin D is associated with increased bone resorption markers and lower bone density at the proximal femur in normal females: a population-based study. *Exp Clin Endocrinol Diabetes* 1996; 104 (3): 289-92.
11. Dawson-Hughes B, Harris SS, Krall EA, Dallal GE. Effect of calcium and vitamin D supplementation on bone density in men and women 65 years of age or older. *N Engl J Med* 1997; 337: 670-6.
12. Bischoff-Ferrari HA, Dietrich T, Orav EJ, Dawson-Hughes B. Positive association between 25-hydroxy vitamin D levels and bone mineral density: a population-based study of younger and older adults. *Am J Med* 2004; 116: 634-9.
13. Slinin Y, Paudel ML, Taylor BC, Fink HA, Ishani A, Canales MT et al. 25-Hydroxyvitamin D levels and cognitive performance and decline in elderly men. Osteoporotic Fractures in Men (MrOS) Study Research Group. *Neurology* 2010; 74 (1): 33-41.
14. Annweiler C, Allali G, Allain P, Bridenbaugh S, Schott AM, Kressig RW et al. Vitamin D and cognitive performance in adults: a systematic review. *Eur J Neurol* 2009; 16 (10): 1083-9.
15. Elmadfa I, Bartens C, Jakob E, König J. Nutritional status and the immune system: fat-soluble vitamins and other nutrients. *Bibl Nutr Dieta* 1994; (51): 136-41.
16. Dietary reference intakes (DRI). Institute of medicine (2000).
17. Lambert J, Agostoni C, Elmadfa I, Hulshof K, Krause E, Livingstonstone B, Socha P et al. Dietary intake and nutritional status of children and adolescents in Europe. *Br J Nutr* 2004; 92 (2 Suppl.): 147S-211S.
18. Al-Tahan J, González-Gross M, Pietrzik K. B vitamin status and intake in European adolescents. A review of the literature. *Nutr Hosp* 2006; 21 (4): 452-65.
19. Scharla SH. Prevalence of subclinical vitamin D deficiency in different European countries. *Osteoporos Int* 1998; 8 (2 Suppl.): 7S-12S.
20. Lehtonen-Veromaa M, Möttönen T, Irjala K, Kärkkäinen M, Lamberg-Allardt C, Hakola P et al. Vitamin D intake is low and hypovitaminosis D common in healthy 9- to 15-year-old Finnish girls. *Eur J Clin Nutr* 1999; 53 (9): 746-51.
21. Ovesen L, Andersen R, Jakobsen. Geographical differences in vitamin D status, with particular reference to European countries. *J Proc Nutr Soc* 2003; 62 (4): 813-21.
22. Zerwekh JE. Blood biomarkers of vitamin D status. *Am J Clin Nutr* 2008; 87 (Suppl.): 1087S-91S.
23. Millen AE, Bodnar LM. Vitamin D assessment in population-based studies: a review of the issues. *Am J Clin Nutr* 2008; 87 (Suppl.): 1102S-5S.
24. Lamberg-Allardt C, Viljakainen HT. 25-Hydroxyvitamin D and functional outcomes in adolescents. *Am J Clin Nutr* 2008; 88 (Suppl.): 534S-6S.
25. Cranney A, Weiler HA, O'Donnell S, Puil L. Summary of evidence-based review on vitamin D efficacy and safety in relation to bone health. *Am J Clin Nutr* 2008; 88 (2): 513S-519S.

26. Lehtonen-Veromaa MK, Mottonen TT, Nuotio IO, Irtala KM, Leino AE, Viikari JS. Vitamin D and attainment of peak bone mass among peripubertal Finnish girls: a 3-y prospective study. *Am J Clin Nutr* 2002; 76 (6): 1446-53.
27. Gordon CM, DePeter KC, Feldman HA, Grace E, Emans SJ. Prevalence of vitamin D deficiency among healthy adolescents. *Arch Pediatr Adolesc Med* 2004; 158 (6): 531-7.
28. Stoffman N, Gordon CM. Vitamin D and adolescents: what do we know? *Curr Opin Pediatr* 2009; 21: 465-471
29. McKenna MJ, Freaney R. Secondary hyperparathyroidism in the elderly: means to defining hypovitaminosis D. *Osteoporos Int.* 1998; 8 (2 Suppl.): 3S-6S.
30. Ernährung, D.G.f. Referenzwerte für die Nährstoffzufuhr. Frankfurt: Umschau/Braus GmbH 2000; 1.
31. Carter GD, Carter R, Jones J, Berry J. How accurate are assays for 25-hydroxyvitamin D? Data from the international vitamin D external quality assessment scheme. *Clin Chem* 2004; 50: 2195-7.
32. Carter GD, Carter CR, Gunter E, Jones J, Jones G, Makin HL et al. Measurement of vitamin D metabolites: an international perspective on methodology and clinical interpretation. *J Steroid Biochem Mol Biol* 2004; 89-90, 467-71.
33. Binkley N, Krueger D, Cowgill CS, Plum L, Lake E, Hansen KE et al. Assay variation confounds the diagnosis of hypovitaminosis D: a call for standardization. *J Clin Endocrinol Metab* 2004; 89: 3152-7.
34. Lips P, Chapuy MC, Dawson-Hughes B, Pols HA, Holick MF. An international comparison of serum 25-hydroxyvitamin D measurements. *Osteoporos Int* 1999; 9: 394-7.
35. Koenig J, Elmadfa I. Status of calcium and vitamin D of different population groups in Austria. *Int J Vitam Nutr Res* 2000; 70 (5): 214-20.
36. Andersen R, Molgaard C, Skovgaard LT, Brot C, Cashman KD, Chabros E et al. Teenage girls and elderly women living in northern Europe have low winter vitamin D status. *Eur J Clin Nutr* 2005; 59: 533-41.
37. Gregory JR, Lowe S, Bates CJ, Prentice A, Jackson LV, Smithers G et al. National Diet and Nutrition Survey: young people aged 4 to 18 years. Report of the diet and nutrition survey 2000; London: HMSO.
38. Bonofiglio D, Maggiolini M, Marsico S, Giorno A, Catalano S, Aquila S et al. Critical years and stages of puberty for radial bone mass apposition during adolescence. *Horm Metab Res* 1999; 31 (8): 478-82.
39. Smotkin-Tangorra M, Purushothaman R, Gupta A, Nejati G, Anhalt H, Ten S. Prevalence of vitamin D insufficiency in obese children and adolescents. *J Pediatr Endocrinol Metab* 2007; 20 (7): 817-23.
40. Molnar D, Decsi T, Koletzko B. Reduced antioxidant status in obese children with multimetabolic syndrome. *Int J Obes Relat Metab Disord* 2004; 28 (10): 1197-202.
41. Winklhofer-Roob BM, van't Hof MA, Shmerling DH. Reference values for plasma concentrations of vitamin E and A and carotenoids in a Swiss population from infancy to adulthood, adjusted for seasonal influences. *Clin Chem* 1997; 43 (1): 146-53.
42. Aeberli I, Molinari L, Spinaz G, Lehmann R, l'Allemand D, Zimmermann MB. Dietary intakes of fat and antioxidant vitamins are predictors of subclinical inflammation in overweight Swiss children 1-3. *Am J Clin Nutr* 2006; 84: 748-55.
43. Herbeth B, Spycykerelle Y, Deschamps JP. Determinants of plasma retinol, beta-carotene, and alpha-tocopherol during adolescence. *Am J Clin Nutr* 1991; 54 (5): 884-9.
44. World Health Organisation. Obesity, preventing and managing the global epidemic: Report of the WHO consultation of obesity. In: Report of the WHO consultation of obesity. Geneva: World Health Organisation 1997.
45. Biesalski H.K, Fürst P, Kasper H, Kluthe R, Pöler W, Puchstein C et al. Ernährungsmedizin. München, Jena: Elsevier GmbH, Urban Fischer Verlag 2004.
46. Board F.a.N. Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids. Washington DC 2000: *National Academy Press*.
47. Malvy JM, Mourey MS, Carlier C, Caces P, Dostalova L, Montagnon B et al. Retinol, beta-carotene and alpha-tocopherol status in a French population of healthy children. *Int J Vitam Nutr Res* 1989; 59 (1): 29-34.
48. Tanner J.M. Growth and maturation during adolescence. *Nutr Rev.* 1981; 39 (2): 43-55.
49. Buzina R, Grgi Z, Jusi M, Sapunar J, Milanovi N, Brubacher G. Nutritional status and physical working capacity. *Hum Nutr Clin Nutr* 1982; 36 (6): 429-38.
50. Hercberg S, Preziosi P, Galan P, Devanlay M, Keller H, Bourgeois C et al. Vitamin status of a healthy French population: dietary intakes and biochemical markers. *Int J Vitam Nutr Res* 1994; 64 (3): 220-32.
51. Drott P, Meurling S, Gebre-Medhin M. Interactions of vitamins A and E and retinol-binding protein in healthy Swedish children—evidence of thresholds of essentiality and toxicity. *Scand J Clin Lab Invest* 1993; 53 (3): 275-80.
52. Marktl W, Rudas B, Brubacher G. The vitamin status of Viennese school children aged 11-12 years. *Int J Vitam Nutr Res* 1982; 52 (2): 198-206.
53. Crawley HF, While D. The diet and body weight of British teenage smokers at 16-17 years. *Eur J Clin Nutr* 1995; 49 (12): 904-14.
54. Kraljovicová-Kudláčková M, Simoncic R, Béderová A, Grancicová E, Magálová T. Influence of vegetarian and mixed nutrition on selected haematological and biochemical parameters in children. *Nahrung* 1997; 41 (5): 311-4.
55. Hassapidou M, Kafatos A, Manoukas G. Dietary vitamin E intake and plasma tocopherol levels of a group of adolescents from Spili, Crete. *Int J Food Sci Nutr* 1996; 47 (5): 365-8.
56. Malvy DJ, Burtshy B, Dostalova L, Amedee-Manesme O. Serum retinol, beta-carotene, alpha-tocopherol and cholesterol in healthy French children. *Int J Epidemiol* 1993; 22 (2): 237-46.
57. Yetley EA. Assessing the vitamin D status of the US population. *Am J Clin Nutr* 2008; 88 (Suppl.): 558S-64S.
58. Schrijver, J. Biochemical markers for micronutrient status and their interpretation, in modern lifestyles, lower energy intake and micronutrient status. Springer-Verlag 1991.
59. Juttman JR, Visser TJ, Buurman C, de Kam E, Birkenhäger JC. Seasonal fluctuations in serum concentrations of vitamin D metabolites in normal subjects. *Br Med J (Clin Res Ed)* 1981; 25-282 (6273): 1349-52.
60. Tjellesen L, Christiansen C. Vitamin D metabolites in normal subjects during one year. A longitudinal study. *Scand J Clin Lab Invest* 1983; 43 (1): 85-9.
61. Woitge HW, Scheidt-Nave C, Kissling C, Leidig-Bruckner G, Meyer K, Grauer A et al. Seasonal variation of biochemical indexes of bone turnover: results of a population-based study. *J Clin Endocrinol Metab* 1998; 83 (1): 68-75.
62. Krall EA, Sahyoun N, Tannenbaum S, Dallal GE, Dawson-Hughes B. Effect of vitamin D intake on seasonal variations in parathyroid hormone secretion in postmenopausal women. *N Engl J Med* 1989; 321 (26): 1777-83.
63. Moreno LA, Mesana MI, Fleta J, Ruiz JR, González-Gross MM, Sarría A et al and the AVENA Study Group. Overweight, obesity and body fat composition in Spanish adolescents. The AVENA Study. *Ann Nutr Metab* 2005; 49: 71-76.
64. Lenders CM, Feldman HA, Von Scheven E, Merewood A, Sweeney C, Wilson DM et al. Relation of body fat indexes to vitamin D status and deficiency among obese adolescents. *Am J Clin Nutr* 2009; 90: 459-67.
65. Grant WB, Cross HS, Garland CF, Gorham ED, Moan J, Peterlik M et al. Estimated benefit of increased vitamin D status in reducing the economic burden of disease in western Europe. *Progress in Biophysics and Molecular Biology* 2009; 99: 104-113.
66. Serra-Majem, L. Vitamin and mineral intakes in European children. Is food fortification needed? *Public Health Nutrition* 2001; 4(1A): 101-107.
67. Frances A, Tylavsky, Sulin Cheng, Arja Lyytikäinen, Heli Viljakainen, Christel Lamberg-Allardt. Strategies to Improve Vitamin D Status in Northern European Children: Exploring the Merits of Vitamin D Fortification and Supplementation. *J Nutr* 2006; 136: 1130-1134.
68. Jain A, Agrawal BK, Varma M, Jadhav A. Antioxidant status and smoking habits: relationship with diet. *Singapore Med J* 2009; 50 (6):624.