



Original/Síndrome metabólico

Correlation of serum vitamin D level with type 1 diabetes mellitus in children: a meta-analysis

Changwei Liu¹, Mi Lu², Xiaona Xia¹, Jingwen Wang¹, Yuanyuan Wan¹, Lianping He³ and Mei Li¹

¹Nanjing Children's Hospital Affiliated to Nanjing Medical University (Nanjing), Jiangsu. ²School of Medical Imaging and laboratory, Wannan Medical College (Wuhu), Anhui. ³School of Public Health, Wannan Medical College (Wuhu), Anhui. China.

Abstract

Objective: to assess the relationship between serum vitamin D level and type 1 diabetes mellitus (T1DM) in children.

Methods: the following electronic databases were searched until Sep 2014 to identify relevant studies that assessed the relationship between serum vitamin D with T1DM: PubMed\EMbase\Medline\Central Register of Controlled Trials\CBM\Chinese National Knowledge Infrastructure (CNKI)\wangfang; The NOS scale was used to evaluate the quality of studies, and the statistical tests were performed by Stata 11.0 software.

Result: a total of 10 studies were included in this study. Our results showed that serum vitamin D was significantly lower in children with T1DM than in healthy controls (MD = -0.60, $P < 0.05$). No evidence support publication bias in present study.

Conclusion: the meta-analysis suggests that serum vitamin D level is associated with T1DM in children.

(Nutr Hosp. 2015;32:1591-1594)

DOI:10.3305/nh.2015.32.4.9198

Key words: Serum vitamin D. T1DM. Meta-analysis.

CORRELACIÓN DE LOS NIVELES DE VITAMINA D EN SUERO Y LA DIABETES MELLITUS TIPO 1: UN METAANÁLISIS

Resumen

Objetivo: evaluar la relación entre sueros de vitamina D y la diabetes mellitus tipo 1 (T1DM) en niños.

Métodos: fueron utilizadas las siguientes bases de datos electrónicas hasta septiembre de 2014 para identificar los estudios que evaluaron la relación entre la vitamina D y la T1DM: PubMed, EMBASE, MEDLINE, Central Register of Controlled Trials, CBM, China National Knowledge Infrastructure (CNKI), wangfang; las encuestas a la escala se utilizan para evaluar la calidad de los estudios, y los ensayos estadísticos fueron realizados mediante el software Stata 11.0.

Resultado: fueron incluidos un total de 10 estudios. Nuestros resultados mostraron que los niveles séricos de vitamina D fueron significativamente menores en los niños con T1DM que en los controles sanos (MD = -0,60, $P < 0,05$). No hay evidencia de sesgo de publicación en el presente estudio.

Conclusión: el metaanálisis sugiere que la vitamina D está asociada con la T1DM en niños.

(Nutr Hosp. 2015;32:1591-1594)

DOI:10.3305/nh.2015.32.4.9198

Palabras clave: Serum de vitamina D. T1DM. Metaanálisis.

Introduction

Diabetes mellitus is a group of metabolic disease characterized by hyperglycemia resulting from defects in insulin secretion, insulin action, or both¹. The most common classifications include type 1 diabetes mellitus, type 2 diabetes mellitus, and gestational diabetes². Of those, autoimmune destruction of the β -cells of the pancreas which lead to absolute insulin deficiency was defined as type 1 diabetes mellitus (T1DM)³. In Finland, the incidence of T1DM among children has an increasing trend with age and is highest among 14 years or under in the world⁴. The National Pediatric Diabetes Audit reported that 26 867 children and the

Correspondence: Mei Li.
Nanjing Children's Hospital Affiliated to Nanjing Medical University.
Nanjing (Jiangsu), 210008, China.
E-mail: limeilimeimei6868@126.com

Recibido: 4-V-2015.
Aceptado: 26-VI-2015.

youth (aged 0-25 years) suffered from diabetes, and most (95%) of them were T1DM⁵. Diabetic ketoacidosis (DKA) is an acute potentially life-threatening complication of diabetes, which remains the leading cause of hospitalization, morbidity and death in children with T1DM. Therefore, taking into account the incidence rate and fatality rate of T1DM and DKA among children, it is essential to prevent the complication of diabetes.

Vitamin D is a group of fat-soluble secosteroids and the synthesis of vitamin D begins in the skin, where cholesterol is exposed to ultraviolet B photons⁶. In addition, it can also be obtained from foods or supplements⁷. More and more studies found that the vitamin D not only associates with the level of calcium, phosphorus, but also with the blood glucose metabolism and insulin resistance. Relevant studies had shown an association between low ultraviolet B irradiance and high incidence rates of T1DM in children⁸. In recent years, several studies reported a high prevalence of vitamin D deficiency in patients with T1DM, which suggested that vitamin D deficiency correlated with the severity and frequency of T1DM⁹. However, subsequent studies reported inconsistent results¹⁰. Thus, the aim of our meta-analysis was to examine the association between serum vitamin D and T1DM in children, then provide a comprehensive reference to prevent T1DM in children.

Materials and Methods

Literature search

To identify relevant studies that regarding of the association between serum vitamin D level and T1DM, we searched literatures with the English and Chinese language restriction using terms “vitamin D”, “25(OH) D”, “T1DM” and “childhood” in database of PubMed, EMbase, MEDLINE, CENTRAL, CBM, Chinese National Knowledge Infrastructure (CNKI) and Wangfang until September 2014. Eligible full texts were retrieved from the above data.

Inclusion and exclusion criteria

Included criteria: (1) The study was case-control study; (2) The papers published on the relationship between serum vitamin D level and T1DM in children; (3) all T1DM subjects met the criteria of diagnosis. Excluded criteria: (1) Did not meet the inclusion criteria; (2) The paper was a review; (3) The data was not extracted.

Data extraction

According to the inclusion and exclusion criteria, we screened the literature, extracted the data and evaluated the literature quality independently. Data ex-

tracted includes basic information, research design, diagnostic methods, the possible bias, and the features of the objects, results and outcomes.

Quality Assessment

All studies included and their biases were independently assessed by two reviewers using Newcastle-Ottawa Scale (NOS). The NOS contains eight items, categorized into three dimensions including the selection, the comparability and the outcome or exposure of interest¹¹. A ‘star system’ was used to judge data quality: The scores ranged from 0 to 9 stars. Studies with scores of 7 stars or greater were considered to be of high quality¹². The scores were added up to compare the study quality in a semi-quantitative fashion. Studies with scores of 5 stars or greater can be included in our meta-analysis. Disagreements between the reviewers were resolved by open discussion.

Statistical analysis

All statistical tests were performed using Stata11.0 software. We choose appropriate indexes according to the data types. Q test was performed in count data. If heterogeneity was existing among studies ($I^2 \geq 50\%$), a random-effects model was applied to estimate the relationship between serums vitamin D level and T1DM; otherwise, a fixed effects model was applied. In addition, we also calculated the value of mean difference (MD) and 95% confidence interval (CI).

Results

Results of literature research

A total of 69 relevant publications were systematically identified through the previous listed database. Of them, 59 articles were excluded because they either did not meet the inclusion criteria or failed to provide adequate information to determine. Finally, 10 literatures were included in the meta-analysis which assesses the relationship between serum vitamin D and T1DM¹³⁻²². Characteristics of the selected studies are presented in table I. Quality of literature to assess the stability by deleting the studies with scores of 6 stars, the samples less than 50 cases and some studies in which the serum vitamin D levels are too low or too high. There was no significantly statistical change after remerging the results.

The correlation of serum vitamin D with T1DM

A total of 10 literatures were included and heterogeneity testing was carried out in our meta-analysis, due

Table I
General character of studies included

first author	year	n	T1DM	control	NOS score
Bruna Franchi	2014	224	58	166	8
Somia H. Abd-Allah	2014	240	120	120	8
Seham FA Azab	2013	120	80	40	7
Lingfei Wang	2013	95	52	43	6
Ristan M Greer	2012	102	56	46	8
Xiaoyan Huang	2012	50	26	24	6
Yan Zhu	2011	152	52	100	6
Jill H Simmons	2011	57	27	30	7
Bassam S. Bin-Abbas	2011	200	100	100	7
Abdulbari Bener	2008	340	170	170	8
Abdulbari Bener	2008	1580	741	839	8

to I² value was 80.1%, a random-effects model was applied for meta-analysis. The serum vitamin D level in T1DM was lower than in the control (Fig. 1). MD (95%CI) = -0.60(-0.70,-0.49), z=5.51, P<0.05.

Publication bias

Funnel plots and Egger's test were used to test possible bias among literatures, which showed a better symmetry (Egger's test :t=-1.82, P=0.107), suggesting a low likelihood of publication bias.

Discussion

The etiology and natural history of T1DM are still unknown, but both genetics and environmental factors certainly contribute to the development of T1DM²³. It is confirmed that immune factors play a decisive role in onset of T1DM²⁴. T1DM is a T cell-mediated disease that ultimately destroys the capacity of the body to produce and secrete insulin²⁵ and was characterized by the apoptotic destruction of pancreatic β -cell²⁶.

Vitamin D is a fat-soluble secosteroids and plays a central role in the metabolism of calcium and phosphorus. Vitamin D deficiency is a risk factor for type 1 diabetes. Previous studies showed that 1, 25-(OH) 2D3 influences insulin secretion in the pancreatic beta-cell. Indeed, 1,25 D3 prevent human pancreatic islet cells from destruction²⁷. evidence based on observational studies revealed that vitamin D supplementation in infancy might be protective against the development of type 1 diabetes²⁸. Compared with the normal group, this study showed a significantly statistical difference in serum vitamin D level between a child with T1DM and without, which suggested that children with T1DM have a vitamin D deficiency.

This present research indicates that serum vitamin D is significantly lower in children with T1DM than in healthy. Because only some case-control studies were included in this research, causal association was not sure. In addition, Age, illumination, BMI, dieting and physical activity are important determinants of the level of serum vitamin D. Thus, the relationship between serum vitamin D levels in children with T1DM should be further studied. Moreover, there is heterogeneity between the studies included. The possible reason may be that studies are from different area, diagnostic crite-




Fig. 1.—forest plot of the correlation of serum vitamin D with T1DM in children.

ria, and sample size. Sensitivity analysis was conducted to assess the stability. There was no significantly statistical change in conclusion after deleting relevant factors.

Conclusion

The meta-analysis suggests that serum vitamin D is associated with T1DM in children. And provide a comprehensive reference to prevent T1DM in children.

Acknowledgement

Key Project supported by Medical Science and technology development Foundation, Nanjing Department of Health (No.YKK14119), Wannan Medical college key scientific research projects Engagement Fund (WK2014Z05), National Training Programs of Innovation and Entrepreneurship for Under-graduates (No.AH201510368042) and Undergraduate Scientific Research Foundation of Wannan Medical College (No. WK2014S18) .

Conflict of interest

None declared.

References

1. Expert Committee on the, D. and M. Classification of Diabetes. Report of the expert committee on the diagnosis and classification of diabetes mellitus. *Diabetes Care*, 2003. 26 Suppl 1: S5-20.
2. Triplitt, C., C. Solis-Herrera, C. Reasner, R.A. DeFronzo and E. Cersosimo, *Classification of Diabetes Mellitus*, in *Endotext*, L.J. De Groot, et al., Editors. 2000: South Dartmouth (MA).
3. Kerner, W., J. Bruckel and A. German Diabetes, Definition, classification and diagnosis of diabetes mellitus. *Exp Clin Endocrinol Diabetes*, 2014. 122(7): 384-6.
4. Karvonen, M., Z. Cepaitis and J. Tuomilehto, Association between type 1 diabetes and Haemophilus influenzae type b vaccination: birth cohort study. *BMJ*, 1999. 318(7192): 1169-72.
5. Kmietowicz, Z., More children showing early signs of serious diabetes complications, audit finds. *BMJ*, 2015. 350: h1265.
6. Adamczak, D.M., J.K. Nowak, M. Frydrychowicz, M. Kaczmarek and J. Sikora, The role of Toll-like receptors and vitamin D in diabetes mellitus type 1--a review. *Scand J Immunol*, 2014. 80(2): 75-84.
7. Feng, R., Y. Li, G. Li, Z. Li, Y. Zhang, Q. Li, et al., Lower serum 25 (OH) D concentrations in type 1 diabetes: A meta-analysis. *Diabetes Res Clin Pract*, 2015.
8. Mohr, S.B., C.F. Garland, E.D. Gorham and F.C. Garland, The association between ultraviolet B irradiance, vitamin D status and incidence rates of type 1 diabetes in 51 regions worldwide. *Diabetologia*, 2008. 51(8): 1391-8.
9. Mutlu, A., G.Y. Mutlu, E. Ozsu, F.M. Cizmecioglu and S. Hattun, Vitamin D deficiency in children and adolescents with type 1 diabetes. *J Clin Res Pediatr Endocrinol*, 2011. 3(4): 179-83.
10. Simpson, M., H. Brady, X. Yin, J. Seifert, K. Barriga, M. Hoffman, et al., No association of vitamin D intake or 25-hydroxyvitamin D levels in childhood with risk of islet autoimmunity and type 1 diabetes: the Diabetes Autoimmunity Study in the Young (DAISY). *Diabetologia*, 2011. 54(11): 2779-88.
11. Stang, A., Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol*, 2010. 25(9): 603-5.
12. Qi, L., X. Qi, H. Xiong, Q. Liu, J. Li, Y. Zhang, et al., Type 2 diabetes mellitus and risk of malignant melanoma: a systematic review and meta-analysis of cohort studies. *Iran J Public Health*, 2014. 43(7): 857-66.
13. Franchi, B., M. Piazza, M. Sandri, F. Mazzei, C. Maffei and A.L. Boner, Vitamin D at the onset of type 1 diabetes in Italian children. *Eur J Pediatr*, 2014. 173(4): 477-82.
14. Abd-Allah, S.H., H.F. Pasha, H.A. Hagrass and A.A. Alghobashy, Vitamin D status and vitamin D receptor gene polymorphisms and susceptibility to type 1 diabetes in Egyptian children. *Gene*, 2014. 536(2): 430-4.
15. Azab, S.F., S.H. Saleh, W.F. Elsaed, S.M. Abdelsalam, A.A. Ali and A.M. Esh, Vitamin D status in diabetic Egyptian children and adolescents: a case-control study. *Ital J Pediatr*, 2013. 39: 73.
16. Wang, L., S. Luo. and Y. Chen, Changes of serum 25(OH) D levels in children with T1DM. *China Prac Med*, 2013. 8(4): 60.
17. Greer, R.M., S.L. Portelli, B.S. Hung, G.J. Cleghorn, S.K. McMahon, J.A. Batch, et al., Serum vitamin D levels are lower in Australian children and adolescents with type 1 diabetes than in children without diabetes. *Pediatr Diabetes*, 2013. 14(1): 31-41.
18. Ma, X.H.Z.C.Z., Detection of serum 25(OH) D levels in children with T1DM and clinical significance. *Journal of Hainan Medical college*, 2012. 18(4): 524-29.
19. Yan Zhu, J.Z., Hongzhou Ye, Study on the correlation of 25(OH) D and children with T1DM and diabetic ketoacidosis. *Journal of Clinical Pediatrics*, 2011. 29(12): 1118-20.
20. Simmons, J.H., M. Raines, K.D. Ness, R. Hall, T. Gebretsadik, S. Mohan, et al., Metabolic control and bone health in adolescents with type 1 diabetes. *Int J Pediatr Endocrinol*, 2011. 2011(1): 13.
21. Bin-Abbas, B.S., M.A. Jabari, S.D. Issa, A.H. Al-Fares and S. Al-Muhsen, Vitamin D levels in Saudi children with type 1 diabetes. *Saudi Med J*, 2011. 32(6): 589-92.
22. Bener, A., A. Alsaied, M. Al-Ali, A.S. Hassan, B. Basha, A. Al-Kubaisi, et al., Impact of lifestyle and dietary habits on hypovitaminosis D in type 1 diabetes mellitus and healthy children from Qatar, a sun-rich country. *Ann Nutr Metab*, 2008. 53(3-4): 215-22.
23. Banin, P., F. Rimondi, A. De Togni, S. Cantoni, G. Chiari, L. Iughetti, et al., Type 1 diabetes (T1DM) in children and adolescents of immigrated families in Emilia-Romagna (Italy). *Acta Biomed*, 2010. 81(1): 35-9.
24. Zhou, Q.L.H.X.Y., Research of the relationship between Coxsackie virus infection and T lymphocyte subsets and type 1 diabetes mellitus. *Journal of Southern Medical University*, 2010. 30(12): 2699-701.
25. American Diabetes, A., Diagnosis and classification of diabetes mellitus. *Diabetes Care*, 2014. 37 Suppl 1: S81-90.
26. Cnop, M., N. Welsh, J.C. Jonas, A. Jorns, S. Lenzen and D.L. Eizirik, Mechanisms of pancreatic beta-cell death in type 1 and type 2 diabetes: many differences, few similarities. *Diabetes*, 2005. 54 Suppl 2: S97-107.
27. Riachy, R., B. Vandewalle, S. Belaich, J. Kerr-Conte, V. Gmyr, F. Zerimech, et al., Beneficial effect of 1,25 dihydroxyvitamin D3 on cytokine-treated human pancreatic islets. *J Endocrinol*, 2001. 169(1): 161-8.
28. Zipitit, C.S. and A.K. Akobeng, Vitamin D supplementation in early childhood and risk of type 1 diabetes: a systematic review and meta-analysis. *Arch Dis Child*, 2008. 93(6): 512-7.