



## Trabajo Original

Epidemiología y dietética

### Relationship of long-term macronutrients intake on anabolic-catabolic hormones in female elite volleyball players

*Relación de la ingesta a largo plazo de macronutrientes en las hormonas anabólicas-catabólicas de jugadoras de voleibol elite*

Juan Mielgo-Ayuso<sup>1</sup>, Michael C. Zourdos<sup>2</sup>, Aritz Urdampilleta<sup>1</sup>, Julio Calleja-González<sup>3</sup>, Jesús Seco<sup>4</sup> and Alfredo Córdova<sup>5</sup>

<sup>1</sup>Elikaesport, Nutrition, Innovation & Sport. Donostia-San Sebastián, Spain. <sup>2</sup>Department of Exercise Science and Health Promotion. Muscle Physiology Laboratory. Florida Atlantic University. Boca Raton, Florida. <sup>3</sup>Department of Physical Education and Sports. University of Basque Country (UPV-EHU). Lejona, Vizcaya. Spain. <sup>4</sup>Department of Nursing and Physical Therapy. Institute of Biomedicine (BIOMED). University of Leon. Leon, Spain. <sup>5</sup>Department of Biochemistry and Physiology. School of Physical Therapy. University of Valladolid. Soria, Spain

#### Abstract

**Introduction:** Specific macronutrient distribution and training can alter acute and chronic hormone behavior and, subsequently, sport performance.

**Objective:** The main aim was to examine relationships between dietary intake and anabolic/catabolic hormone response in elite female volleyball players during a 29-week season.

**Methods:** Twenty-two elite female volleyballers (26.4 ± 5.6 years; 178 ± 9 cm; 67.1 ± 7.5 kg) had dietary intake (seven-day dietary recall and food frequency questionnaire), blood concentration of anabolic/catabolic hormones concentration, physical performance, and body composition assessed at four time points: a) T1: baseline/pre-testing; b) T2: eleven weeks after T1; c) T3: ten weeks after T2; and d) T4: eight weeks after T3. Hormones evaluated were: total testosterone (TT), free testosterone (FT) adrenocorticotrophic hormone (ACTH), and cortisol (C), along with hormone ratios.

**Results:** Positive correlations were observed between carbohydrate/protein ratio with  $\Delta FT$  ( $r = 0.955$ ;  $p < 0.001$ ),  $\Delta TT/C$  ratio ( $r = 0.638$ ;  $p = 0.047$ ), and  $\Delta FT/C$  ratio ( $r = 0.909$ ;  $p < 0.001$ ). Significant and negative correlations were found between protein intake with  $\Delta TT$  ( $r = -0.670$ ;  $p = 0.034$ ), and FT ( $r = -0.743$ ;  $p < 0.001$ ), carbohydrate intake and  $\Delta ACTH$  ( $r = -0.658$ ;  $p = 0.006$ ). No relationships were observed regarding  $\Delta$ cortisol. On the other hand, there was no change ( $p > 0.05$ ) in body mass or body mass index at any time point, and the sum of six skinfolds improved ( $p < 0.05$ ) from T1 (86.5 ± 6.9 mm) to T4 (75.2 ± 5.6 mm) as did muscle mass (T1: 28.9 ± 0.7 kg vs T4: 30.1 ± 0.8 kg). Vertical jump, spike-jump and speed improved ( $p < 0.05$ ) from T1 to T4.

**Conclusions:** A high carbohydrate/protein ratio was associated with positive changes in anabolism, while high protein and low carbohydrates (CHO) were associated with an attenuated anabolic response.

#### Key words:

Dietary intake.  
Testosterone. Cortisol.  
Macronutrients.  
Performance.

#### Resumen

**Introducción:** la distribución específica de macronutrientes y el entrenamiento pueden alterar el comportamiento agudo y crónico de las hormonas y, posteriormente, el rendimiento deportivo.

**Objetivo:** el objetivo principal del estudio fue examinar la relación entre la ingesta dietética y la respuesta anabólica/catabólica hormonal de jugadoras de élite de voleibol durante una temporada de 29 semanas.

**Métodos:** se evaluó en 22 jugadoras de élite (26,4 ± 5,6 años, 178 ± 9 cm, 67,1 ± 7,5 kg) la ingesta dietética (mediante un registro de siete días y un cuestionario de frecuencia de consumo de alimentos), la concentración de hormona anabólica/catabólica en sangre, el rendimiento físico y la composición corporal en cuatro puntos durante la temporada: a) T1: pre-pretemporada; b) T2: once semanas después de T1; c) T3: diez semanas después de T2; y d) T4: ocho semanas después de T3. Las hormonas evaluadas fueron: testosterona total (TT), testosterona libre (FT), hormona adrenocorticotropa (ACTH) y cortisol (C); se calcularon distintos ratios hormonales.

**Resultados:** se observaron correlaciones positivas entre la ratio carbohidratos/proteínas con  $\Delta FT$  ( $r = 0,955$ ;  $p < 0,001$ ),  $\Delta TT/C$  ( $r = 0,638$ ;  $p = 0,047$ ) y  $\Delta FT/C$  ( $r = 0,909$ ;  $p < 0,001$ ). Se encontraron correlaciones significativas y negativas entre la ingesta proteica con  $\Delta TT$  ( $r = -0,670$ ;  $p = 0,034$ ) y FT ( $r = -0,743$ ;  $p < 0,001$ ), la ingesta de carbohidratos y  $\Delta ACTH$  ( $r = -0,658$ ;  $p = 0,006$ ). No se observó ningún tipo de correlación con el  $\Delta$ cortisol. Por otro lado, no hubo cambios ( $p > 0,05$ ) en la masa corporal ni en el índice de masa corporal en ningún momento, mientras que el sumatorio de seis pliegues mejoró ( $p < 0,05$ ) de T1 (86,5 ± 6,9 mm) a T4 (75,2 ± 5,6 mm), así como la masa muscular (T1: 28,9 ± 0,7 kg frente a T4: 30,1 ± 0,8 kg). El salto vertical, el salto de remate y la velocidad mejoraron ( $p < 0,05$ ) de T1 a T4.

**Conclusión:** en conclusión, una alta ratio de carbohidratos/proteínas se asoció con cambios positivos en el anabolismo, mientras que una ingesta alta de proteína y baja de CHO se asoció con una respuesta anabólica atenuada.

#### Palabras clave:

Ingesta dietética.  
Testosterona. Cortisol.  
Macronutrientes.  
Rendimiento deportivo.

Received: 26/11/2016

Accepted: 01/03/2017

*Author contributions:* Juan Mielgo Ayuso conceived and designed the experiments, performed the experiments, analyzed the data, wrote and reviewed the paper. Michael C. Zourdos analyzed the data, wrote and reviewed the paper. Aritz Urdampilleta wrote and reviewed the paper. Jesús Seco conceived and designed the experiments, wrote and reviewed the paper. Alfredo Córdova conceived and designed the experiments, wrote and reviewed the paper. DOI: <http://dx.doi.org/10.20960/nh.763>

Mielgo Ayuso J, Zourdos MC, Urdampilleta A, Calleja González J, Seco J, Córdova A. Relationship of long-term macronutrients intake on anabolic-catabolic hormones in female elite volleyball players. Nutr Hosp 2017;34:1155-1162

DOI: <http://dx.doi.org/10.20960/nh.763>

#### Correspondence:

Juan Mielgo Ayuso. Elikaesport, Nutrition, Innovation & Sport. Astigarrako Bidea, 1. Local 4. 20180 Oiartzun Donostia-San Sebastián, Spain  
e-mail: [juankaya@msn.com](mailto:juankaya@msn.com),  
[j.mielgo@elikaesport.com](mailto:j.mielgo@elikaesport.com)

## INTRODUCTION

Volleyball is intermittent, and often has  $\leq 48$  hours between training and matches, thus recovery-status must be monitored to avoid diminished performance (1) during match play. Specifically, inadequate recovery or overreaching can lead to elevated cortisol (C) and decreased serum testosterone (T) (2-4). Importantly, enhanced resting testosterone/cortisol (T/C) ratio has been associated with increased strength (5). Therefore, monitoring T and C can provide insight into an athlete's recovery/readiness (6), and can be a tool to program daily volume/intensity of training (7).

Physiologically, T and C are secreted via the hypothalamic-pituitary-adrenal axis. Adrenocorticotrophic releasing hormone (ACTH) facilitates C release, which is exercise volume-dependent (8,9). The steroid hormone, T, has both anabolic and anti-catabolic effects, contributing to the growth/remodelling of tissues (10), and when elevated enhances muscle glycogen synthesis (11). Moreover, nutritional intake (10,12) and specific macronutrient intake can effect acute hormone levels (13-17), however, chronic hormone concentrations (5), not acute (18), are most suggestive of long-term performance changes. Indeed, chronic high protein (8) and fat consumption (19) have been associated with elevated resting T. Data have shown decreased resting T when the carbohydrate/protein (CHO/P) ratio favors protein (8,12), but this is mitigated with sufficient CHO (8), suggesting T/C ratio is CHO-dependent (20).

Despite the data, which exists regarding macronutrient intake and changes in T/C ratio, the majority of this data is related to short-term protocols and male athletes (21,22). Therefore, the primary aim of this study was to examine relationships between total energy and macronutrient intake in conjunction with controlled training on chronic anabolic/catabolic hormone changes in elite female volleyball players during a 29-week season. We hypothesized that a low CHO/P ratio would be associated with decreased T and T/C ratio, and increased C and ACTH. Further, we anticipated the opposite hormone response if a high CHO/P ratio was observed.

## MATERIALS AND METHODS

### ETHICS COMMITTEE

All participants were informed of the protocol and risks/benefits and signed a written consent form prior to participation. The study was designed in compliance with the recommendations for clinical research of the Declaration of Helsinki of the World Medical Association (2008). The protocol was reviewed and approved by the Ethics Committee of the University of Leon.

### PARTICIPANTS AND DAILY TRAINING

Twenty-two elite female volleyball players ( $26.4 \pm 5.6$  years;  $178 \pm 9$  cm;  $67.1 \pm 7.5$  kg) participate from two teams of ten

teams from the Spanish First National Professional League that the previous season, were the best two teams of the regular league, participated in this study. Initially, 24 players began the study; however, two participants left the study (one in each group) during the experimental protocol due to injury sustained during the season. Of the participants, 62% are currently or have previously represented a national team (Spain: ten players; Argentina: one player; Brazil: three players; and Serbia: one player). All participants performed the same training program and matches throughout the season supervised by the same physician (Fig. 1). The training program was in accordance with team sport periodization theory (23), and similar to previous programs in this population (24). A standard training day involved two sessions: a) a morning session with 30 minutes of jogging and 90 minutes of resistance training; and b) an afternoon session, with a 3-hour volleyball practice. The specifications of this training program were followed every day except on match days and days immediately following a match. On post-match days players engaged in recovery training (i.e., 20 minutes jogging and stretching).


All athletes completed a medical history questionnaire and electrocardiographic and cardiopulmonary examinations. No participants had any diseases, nor smoke, drank alcohol or took medications (including oral contraceptives), which would alter hormone response. All volleyballers had normal menstrual cycles of 28-31 interval days before the study.

## EXPERIMENTAL PROTOCOL AND ASSESSMENT PLAN

This study was designed to examine relationships between macronutrient consumption and hormonal changes throughout a 29-week competitive season. The 22 participants were evaluated for dietary intake, physical performance, anthropometrics and hormone levels at four time points: a) baseline/pre-testing (T1-before preseason); b) eleven weeks later (T2); c) ten weeks following T2 (T3); and d) eight weeks following T3 (T4).

## BLOOD COLLECTION AND BIOCHEMICAL ANALYSIS

Blood samples (10 ml) were collected from the antecubital vein from all players, at each time point in basal conditions, which followed an overnight fast and 36 hours removed from exercise. Further, for blood collection, the players arrived at the laboratory at 8:30 am, and upon arrival sat comfortably for 30 minutes. Additionally, blood samples were obtained during the early follicular phase to avoid coinciding with the menstruation or ovulatory phases. The blood sample was left in room temperature for ten minutes before 15 minutes of centrifugation at  $4^{\circ}\text{C}$  and 3,000 rpm. Next, serum was separated and stored in aliquots at  $-20^{\circ}\text{C}$  until analysis. All analyses were conducted in accordance with the manufacturer's instructions.



**Figure 1.**

Amount of time and type of training performed by players in each week of study.

### Serum total testosterone (TT)

Serum TT was measured using commercially available enzyme-linked immunosorbent assay (ELISA) kits (DRG testosterone ELISA kit®, DRG Instruments GmbH, Marburg/Lahn, Germany). Intra-assay coefficient of variation (CV) was 4.3% and inter-assay CV was 9.2%. FT was obtained by a validated formula (25).

### Sex hormone binding globulin (SHBG)

SHBG concentrations were measured by a chemoluminescence based on immunometric method (Immulite, 2000), which was from the same supplier as TT. Intra-assay CV was 3% and interassay CV was 7%.

### Cortisol (C)

Serum C levels were measured by an enzyme-linked fluorescent assay with the aid of a multiparametric analyser (Minividas®, Biomerieux, Marcy l'Etoile, France). The substrate, 4-methyl umbeliferone, was used and was capable of a fluorescent emission at 450 nm after stimulation at 370 nm. Intra-assay CV was 5.7% and interassay CV was 6.2%.

### Adrenocorticotrophic hormone (ACTH)

ACTH was determined with a radioimmunoassay (RIA). Intra-assay CV was 4.8% and inter-assay CV was 6.7%.

### Hormone ratio calculations

Total testosterone/cortisol ratio TT/C and FT/C ratios were calculated from TT, FT and C concentrations. Similarly, ACTH/C ratio was calculated from ACTH and C concentrations, and the TT/SHBG ratio was calculated from TT and SHBG concentrations.

### DIETARY ASSESSMENT

Dietitians taught participants to accurately track food intake. After blood collection, at each time point, participants completed a validated food frequency questionnaire (FFQ) (26), used in a similar population (27). This FFQ asked the athletes to recall average energy intake between each time point. The FFQ included 139 different foods and drinks, arranged by food type and meal pattern. The FFQ had frequency categories based upon the number of times items were consumed per day, week and month. Daily consumptions of total energy (kcal) and macronutrients (in grams) were determined by dividing the reported intake by frequency in terms of days (26). Additionally, the participants completed a seven-day dietary recall at all four assessment points and were asked to recall their average total energy and specific macronutrient intake over the previous seven days as a check on the FFQ. When participants weighed food, that data was used; however, when it was not possible to weigh food, an estimation of serving size consumption was determined from either product names, the place of food consumption, or via standard weights of food items or portion size, which participants indicated in a picture booklet

containing 500 photographs of various foods. Food values were converted into total kcal intake and specific macronutrient distribution using the online validated food software package (Easy diet® online version) (28).

Macronutrient intake was compared to the established recommendations for this population: a) total energy: 50-80 kcal/kg/BM/day (29); b) protein: 1.6-1.8 g/kg/BM/day (30); c) fat: 20-35% of total calories (31); and d) CHO 5-8 g/kg/BM/day (29). Further, CHO/P ratio was calculated as CHO (g)/protein (g).

## BODY COMPOSITION AND ANTHROPOMETRIC MEASURES

Anthropometric measurements (height, body mass [BM], and the sum of six skinfolds- $\Sigma$ 6SF) were taken following the protocol of the International Society for the Advancement of Kinanthropometry (32), and the same certified investigator (ISAK level 3) took all anthropometric measurements. The  $\Sigma$ 6SF (triceps, subscapular, supraspinale, abdominal, front thigh, medial calf) was obtained using a Harpenden® skinfold caliper, with a precision of 0.2 mm. Two measurements were taken at each site, and if two measurements at any site were greater than 2.6% technical error of measurement (TEM), then a third measurement was taken. Body mass index (BMI) was calculated using the standard formula: BM/height<sup>2</sup> (kg/m<sup>2</sup>).

Limb girths and muscle mass were also assessed. Limb girths (cm) (relaxed arm, chest, waist, mid-thigh and calf girth) were measured with a metallic Lufkin® measuring tape (W606PM), with a precision of 1 mm. All TEMs for girths assessed were less than 0.45%. Muscle mass (kg) was calculated using the Lee equation (33). All girths were corrected for the skinfold at the site using the following formula: [corrected girth = girth - ( $\pi$  x skinfold thickness at the site)].

## PHYSICAL PERFORMANCE TESTS

To evaluate changes in physical performance, a battery of tests representing volleyball-specific fitness were performed at each time point in an indoor sports hall with standard conditions (temperature: 21 °C and humidity: 60%). The tests were as follows: jump tests (vertical jump [VJ] and spike-jump [SPJ]), a 2 x 18 m sprint test, and the overhead medicine ball throw. These tests have been previously validated as a measure of volleyball-specific fitness and used in a similar population (34).

## STATISTICAL ANALYSES

Results are expressed as mean  $\pm$  standard error. The Shapiro-Wilk test was used to determine normality of data. A one-way repeated measures analysis of variance (ANOVA) was carried out by Greenhouse-Geisser test to check for significant differences between percentage change of hormone concentrations and

hormone ratios from T1-T2, T2-T3, and T3-T4, as well as for changes in physical performance, and body composition parameters at each time point. A Bonferroni post-hoc test was applied for pairwise comparisons. Further, to examine relationships between dietary intake (total kcal and macronutrients), bivariate correlations between changes in hormones and nutritional intake during the season were tested using Pearson or Spearman's rank order correlation test. The hormonal percentage change (% $\Delta$ ) in hormone concentrations was also calculated from T1-T4, T1-T2, T2-T3, and T3-T4. Mean dietary intake for the entire 29 weeks was calculated by summing the average intake of each period and then dividing by 3 ((T1-T2 mean + T2-T3 mean + T3-T4 mean) / 3). Statistical analyses were performed using the IBM Statistical Package (SPSS version 22) and Graphpad Prism (Graphpad Software version 6, San Diego, CA). Significance was set at  $p \leq 0.05$ .

## RESULTS

### RELIABILITY OF FFQ AND SEVEN-DAY DIETARY RECORDS

The results regarding total energy intake and specific macronutrient consumption obtained by the FFQ in comparison to the seven-day dietary records were not significantly different ( $p > 0.05$ ) (data not shown).

### HORMONE: PERCENTAGE CHANGE

Reporting percentage change can effectively illustrate the magnitude of alteration in an observational study. Therefore, table I displays hormone values at T1 and % $\Delta$  of each hormone and hormone ratio in the following periods: T1-T2, T2-T3, and T3-T4. From T1-T2 there were no significant ( $p > 0.05$ ) changes in any hormonal parameter. In T2-T3 there were significant increases ( $p < 0.05$ ) in TT ( $26.1 \pm 11.0\%$ ), SHBG ( $24.8 \pm 11.8\%$ ), and TT/C ( $19.0 \pm 12.1\%$ ), however, no change ( $p > 0.05$ ) was observed in any other hormone parameter during this period. From T3-T4 there were significant changes ( $p < 0.05$ ) in TT ( $+10.4 \pm 4.9\%$ ), ACTH ( $11.5 \pm 5.4\%$ ), and TT/C ( $7.0 \pm 3.8\%$ ). Finally, % $\Delta$  from T1-T4 is displayed in figure 2.


### TOTAL ENERGY AND MACRONUTRIENT-SPECIFIC CONSUMPTION

Table II shows daily total energy intake and specific macronutrient distribution during each period along with average total kcal and macronutrient consumption for the entire study. This table also displays the dietary established recommendations (ER) for this population. Total kcal and specific macronutrient intake were not different ( $p > 0.05$ ) between periods. Compared to ER, athletes consumed lower total energy ( $41.8 \pm 1.9$  kcal) and CHO ( $4.5 \pm 0.2$  g/kg/d) in each period and over the 29 weeks. Con-

**Table I.** Absolute hormone levels at T1 (baseline) and the percentage change during each period

	T1	$\Delta$ (T1-T2)	$\Delta$ (T2-T3)	$\Delta$ (T3-T4)
TT (ng·ml <sup>-1</sup> )	0.29 ± 0.04	-12.2 ± 5.4	26.1 ± 11.0 <sup>a</sup>	10.4 ± 4.9 <sup>a</sup>
FT (pg·ml <sup>-1</sup> )	3.11 ± 0.64	5.3 ± 6.1	8.4 ± 14.8	27.2 ± 11.5
SHBG (nMol·l <sup>-1</sup> )	85.4 ± 18.4	-14.5 ± 9.9	24.8 ± 11.8 <sup>a</sup>	0.9 ± 6.1
ACTH (pg·ml <sup>-1</sup> )	17.0 ± 2.0	49.1 ± 16.4	30.9 ± 10.7	11.5 ± 5.4 <sup>a</sup>
Cortisol (µg·dl <sup>-1</sup> )	20.9 ± 3.1	11.8 ± 7.8	11.0 ± 5.3	1.5 ± 1.2
TT/C	1.90 ± 0.37	-18.7 ± 5.0	n	7.0 ± 3.8 <sup>a</sup>
FT/C	0.022 ± 0.005	-3.4 ± 6.9	3.4 ± 15.9	24.1 ± 9.8

Data expressed as mean ± standard error of the mean. %  $\Delta$  calculated as  $(\frac{T_{final} - T_{initial}}{T_{initial}}) \times 100$ . T1: baseline/pre-testing; T2: eleven weeks after T1; T3: ten weeks after T2; T4: eight weeks after T3. TT: Total testosterone; FT: Free testosterone; SHBG: Sex-hormone binding globulin; ACTH: Adrenocorticotrophic hormone; TT/C: Total testosterone/cortisol ratio; FT/C: Free testosterone/cortisol ratio. <sup>a</sup>Significant magnitude of change in the specific period as determined by the Bonferroni post-hoc test ( $p < 0.05$ ).

**Figure 2.**

Percentage change of each hormone parameter from T1-T4. Data expressed as mean ± standard error of the mean. Percentage change (% $\Delta$ ) calculated as  $(\frac{T_{final} - T_{initial}}{T_{initial}}) \times 100$ .

trastingly, we observed greater average protein ( $2.1 \pm 0.1$  vs  $1.6-1.8$  g/kg/d) and fat consumption ( $35.3 \pm 1.0$  vs 20-35% of total kcal) as compared with ER. Additionally, observed CHO/P ratio was  $2.22 \pm 0.10$  g/day, which was lower than ER.

## RELATIONSHIP BETWEEN DIETARY INTAKE AND HORMONE CONCENTRATIONS

Table III displays specific values of significant bivariate correlations between changes in hormone concentrations and

dietary intake parameters from T1-T4. Significant ( $p < 0.05$ ) and positive correlations were present between: total energy-kcal/kg/day and  $\Delta$ SHBG; CHO-g/kg/day and  $\Delta$ SHBG; and between CHO/P with  $\Delta$ FT,  $\Delta$ TT/C ratio, and  $\Delta$ FT/C ratio. Significant ( $p < 0.05$ ) and negative correlations were present between: protein intake-% of total kcal and  $\Delta$ TT; protein-% of total kcal and FT; CHO-g/kg/day and  $\Delta$ ACTH; and between CHO/P and  $\Delta$ SHBG. No relationship was discovered between  $\Delta$ C and dietary intake or between fat intake and any hormone parameter (data not shown).

## BODY COMPOSITION AND PERFORMANCE ASSESSMENTS

Specific values for body composition changes can be seen in table IV. There was no significant difference ( $p > 0.05$ ) in BM and BMI between periods. There was a significant change ( $p = 0.011$ ) in  $\Sigma$ 6S from T1-T4 (-12.9%). Similarly, muscle mass decreased ( $p = 0.005$ ) from T1-T4 (+2.4%).

Regarding vertical jump power (VJP), values at T2, T3, and T4 were significantly greater ( $p < 0.001$ ) than at T1; however, T2, T3, and T4 were not different ( $p > 0.05$ ) from each other. Similar to VJP, SJP was greater ( $p = 0.003$ ) at T1, T2, and T3 in comparison with T4, and T2, T3, and T4 were not different ( $p > 0.05$ ) from each other. Absolute VJ height was significantly greater ( $p < 0.05$ ) at T3 and T4 as compared with T1 ( $0.31 \pm 0.01$  cm), but not at T2 ( $p > 0.05$ ) (data not shown). Absolute SPJ height did not increase ( $p > 0.05$ ) from T1 to T2, but was significantly greater ( $p < 0.05$ ) at T3 and T4 ( $0.54 \pm 0.02$  cm) vs T1 ( $0.48 \pm 0.01$  cm; +12.5%) (data not shown).

Speed (2 x 18 m sprint) significantly improved ( $p < 0.001$ ; +3.2%) from T1-T2, then plateaued until T4. Throwing distance for the overhead medicine ball throw (OMBT) approached significance from T1-T4 ( $p = 0.092$ ; +1.9%).

**Table II.** Mean total energy and specific macronutrient intake during each period. The 29-week (total season) mean total energy and mean specific macronutrient intake, and the previously established recommendations for this population

	T1-T2	T2-T3	T3-T4	Mean	ER
Energy (kcal/day)	2,890 ± 88	2,790 ± 50	2,810 ± 75	2,830 ± 50	
Energy (kcal/kg/day)	42.4 ± 2.0	40.5 ± 2.1	41.8 ± 2.0 <sup>b</sup>	41.8 ± 1.9	50-80
Protein (g/day)	143 ± 6.8	135 ± 6	138 ± 7	139 ± 6.9	
Protein (g/kg/day)	2.1 ± 0.1	2.0 ± 0.1	2.0 ± 0.1	2.1 ± 0.1	1.6-1.8
Protein (%)	20.0 ± 0.7	19.7 ± 0.6	19.6 ± 0.6	19.8 ± 0.6	
Fat (g/day)	115 ± 6	106 ± 4	108 ± 3.7	109 ± 3.2	
Fat (g/kg/day)	1.7 ± 0.1	1.6 ± 0.1	1.7 ± 0.1	1.6 ± 0.1	
Fat (%)	35.5 ± 1.6	34.9 ± 1.2	35.5 ± 1.1	35.3 ± 1.0	20 - 35
CHO (g/day)	305 ± 11	297 ± 6.1	305 ± 9	302 ± 6.5	
CHO (g/kg/day)	4.5 ± 0.2	4.4 ± 0.2	4.5 ± 0.2	4.5 ± 0.2	5-8
CHO (%)	42.7 ± 1.2	43.9 ± 1.1	43.7 ± 1.0	43.5 ± 0.9	
CHO/P ratio	2.16 ± 0.09	2.25 ± 0.11	2.27 ± 0.11	2.22 ± 0.10	

Data expressed as mean ± standard error of the mean. T1: Baseline/pre-testing; T2: Eleven weeks after T1; T3: Ten weeks after T2; T4: Eight weeks after T3. ER: Established recommendations; CHO: Carbohydrates; CHO/P: Carbohydrate/protein ratio. kcal/day: Kilocalories per day; kcal/kg/day: Kilocalories per kilogram per day; g/day: Grams per day; g/kg/day: Grams per kilogram per day; %: Percentage of macronutrient consumed of total calories.

**Table III.** Correlations between changes in hormones from Δ (T1-T4) (29 weeks) and energy and macronutrient intake (mean during season)

	Δ TT	Δ FT	Δ SHBG	Δ ACTH	Δ Cortisol	Δ TT/C	Δ FT/C
<i>Energy</i>							
kcal/kg/day	ns	ns	0.774**	ns	ns	ns	ns
<i>Protein</i>							
%	-0.670*	-0.746**	ns	ns	ns	ns	ns
<i>CHO</i>							
g/kg/day	ns	ns	0.681*	-0.658*	ns	ns	ns
<i>Ratios</i>							
CHO/P	ns	0.995**	-0.644*	ns	ns	0.638*	0.909**

TT: Total testosterone; FT: Free testosterone; SHBG: Sex-hormone binding globulin; ACTH: Adrenocorticotrophic hormone; TT/C: TT/cortisol ratio; FT/C: FT/cortisol ratio. CHO: Carbohydrates; CHO/P: Carbohydrate/protein ratio. \* $p < 0.05$ . \*\* $p < 0.001$ .

## DISCUSSION

The main finding supports our hypothesis that lower protein intake was significantly and inversely correlated with ΔTT ( $r = -0.670$ ) and ΔFT ( $r = -0.743$ ). Additionally, in support of our hypothesis we observed strong and direct relationships between increased CHO/P ratio and increased anabolism. Specifically, CHO/P ratio was directly correlated with ΔFT ( $r = 0.955$ ), ΔTT/C ( $r = 0.638$ ), and ΔFT/C ( $r = 0.909$ ) over the 29 weeks. Total energy ( $r = 0.774$ ) and CHO ( $r = 0.681$ ) intakes were positively associated with ΔSHBG, and an inverse relationship was found between CHO intake and ΔACTH ( $r = -0.658$ ). However, no relationship was observed between C and any dietary intake param-

eter. Finally, significant improvements were observed from T1-T4 in body composition (i.e., Σ6S and muscle mass).

Even though it is well established that adequate total kcal must be consumed for positive performance outcomes (29), the present study has achieved novelty as the first investigation to examine the seasonal relationship between dietary intake and temporal hormone response in elite female athletes. Interestingly, the average total energy consumption observed over the 29 weeks ( $41.8 \pm 1.9$  kcal/kg/day) was substantially lower than the established recommendations: 50-80 kcal/kg/day. However, the athletes in this study still significantly improved power/speed and body composition, and experienced positive hormonal changes (i.e., TT: +23.6%, FT: +40.1%, TT/C: +6.8%, FT/C: +25.1%) over

**Table IV. Anthropometric, body composition and physical performance results at each assessment period**

	T1	T2	T3	T4	p-value
<i>Anthropometric and body composition</i>					
Body mass (kg)	66.8 ± 2.4	67.3 ± 2.3	67.3 ± 2.4	67.1 ± 1.5	0.701
BMI	21.0 ± 0.4	21.1 ± 0.4	21.7 ± 0.4	21.3 ± 0.4	0.776
Σ6S (mm)	86.5 ± 6.9	76.5 ± 6.3	77.5 ± 6.3	75.2 ± 5.6 <sup>a</sup>	0.011
MM Lee (kg)	28.9 ± 0.7	29.4 ± 0.8	29.6 ± 0.8	30.1 ± 0.8 <sup>a</sup>	0.005
<i>Physical performance tests</i>					
Vertical-jump power (W)	1,635 ± 56	1,730 ± 55 <sup>a</sup>	1,791 ± 76 <sup>a</sup>	1,842 ± 75 <sup>a</sup>	0.001
Spike-jump power (W)	2,024 ± 73	2,114 ± 68 <sup>a</sup>	2,167 ± 89 <sup>a</sup>	2,223 ± 74 <sup>a</sup>	0.003
Speed (sec)	7.48 ± 0.08	7.39 ± 0.08 <sup>a</sup>	7.25 ± 0.08 <sup>a</sup>	7.23 ± 0.08 <sup>a</sup>	< 0.001
OMBT (m)	7.75 ± 0.25	7.73 ± 0.23	7.88 ± 0.26 <sup>a</sup>	7.90 ± 0.25 <sup>b</sup>	0.092

Data are expressed as mean ± standard error of the mean. T1: Baseline/pre-testing; T2: Eleven weeks after T1; T3: Ten weeks after T2; T4: Eight weeks after T3; BMI: Body mass index; Σ6S (mm): Sum of six skinfold sites (triceps, subscapular, supraspinale, abdominal, front thigh, medial calf) reported in millimeters; OMBT: Overhead medicine ball throw. MM Lee: Muscle mass according to the Lee equation; W: Watts. Speed (sec.): Speed results in seconds of the 2 x 18 m sprint test; m: Meters; p-value: Effects tests within subjects by Greenhouse-Geisser test. <sup>a</sup>Significantly different from T1 according to the Bonferroni post-hoc test ( $p < 0.05$ ). <sup>b</sup>Change approached significance with respect to T1. Bolded text denotes statistical significance.

the season, thus it seems that proper balance between training/recovery was achieved and the dietary intake was sufficient to elicit positive training/endocrine adaptations. A possible explanation for the perceived low energy intake, yet beneficial endocrine response, is that the total kcal recommendations for high intensity athletes lack sex-specificity (27). Nevertheless, it does seem that a minimum threshold for total energy exists to optimize the endocrine response, as currently, total energy intake was strongly associated with ΔSHBG ( $r = 0.774$ ). Therefore, in terms of total energy consumption the findings are two-fold: a) female athletes who engage in high-intensity exercise can achieve appropriate recovery while consuming lower total kcal/day than the established recommendations; and b) due to the positive relationship between total energy consumption and anabolic hormone concentrations, a minimum energy intake threshold likely exists to elicit conditions for proper training/recovery balance.

Adequate protein intake in conjunction with training can increase rate of skeletal muscle repair (35). Similarly, T contributes to tissue remodelling (10) and muscle glycogen synthesis (11). However, previous data has demonstrated a negative relationship between protein intake and TT (8). Similarly, the present data revealed a strong inverse relationship between protein intake (% of total kcal) with ΔTT ( $r = -0.670$ ) and ΔFT ( $r = -0.743$ ), but demonstrated a positive relationship between CHO/P with ΔFT ( $r = 0.955$ ), ΔTT/C ( $r = 0.638$ ), and ΔTF/C ( $r = 0.909$ ). Therefore, the ratio of CHO/P seems to be of paramount importance to speed muscle tissue recovery. Interestingly, in the present study, mean protein intake over the entire season comprised  $19.8 \pm 0.6\%$  of total kcals ( $2.1 \pm 0.1$  g/kg/d), which was greater than the ER (1.6-1.8 g/kg/d). However, despite the elevated protein intake we observed positive changes in TT, FT, TT/C, and FT/C, which can be explained by the significant relationships between these changes and CHO/P. Thus, as long as an appropriate CHO/P ratio

is maintained, T will induce muscle glycogen synthesis (11), independent of absolute protein consumption. Additionally, a high-fat diet has increased endogenous T production in males, however, even though elevated fat intake (i.e.,  $35.3 \pm 1.0\%$  of total energy) in comparison to ER (i.e., 20-35%) was observed, there was no relationship between fat consumption and anabolism.

Typically, nutritional strategies have been more important for directly enhancing anabolism rather than blunting the catabolism (35), as various studies have demonstrated protein intake to yield no change in resting C concentrations (8,21,36). Changes in C due to nutrition have mostly been acute (16,37,38), however, it is important to note that an acute C increase is simply a regulatory response to maintain homeostatic blood glucose (39), and not catabolic. In agreement, we did not observe an association between resting C and any dietary parameter. Therefore, alterations in resting C are likely due to training and may take long periods of time (i.e., two years) to manifest (5).

The significant limitation of this study is the absence of a control group who consumed kcals/macronutrients consistent with ER. The control group would provide a basis to examine a cause-effect relationship between dietary intake and hormone response, or if hormone fluctuations are primarily training-dependent. Therefore, it must be highlighted that our results are simply correlational and causation cannot be known; thus, it is necessary for future research to include a control group to determine if a causative relationship exists in this population between CHO/P and increased resting testosterone. Further, the present study lacked measures examining muscle protein turnover, which could directly speak to the anabolic response. However, novelty has been achieved, as no study has examined the relationship between nutritional intake and anabolic/catabolic hormones over an entire volleyball season in elite female players and, in general, little data exist in regards to chronic mechanistic adaptations in elite female athletes.

## CONCLUSIONS

In conclusion, dietary tracking over an entire 29-week season in elite female volleyball players revealed dietary intake which varied from ER in that: lower total energy and CHO were consumed, while greater protein and fat were consumed. Importantly, this study is novel as it is the first to show that CHO/P ratio is positively related to  $\Delta$ FT,  $\Delta$ TT/C, and  $\Delta$ F/T/C over an entire season. Practically speaking, elite female volleyballers should aim to maintain a positive CHO/P ratio to aid in maintaining an appropriate balance of the training/recovery paradigm during the season. Finally, further research should be performed with various nutritional strategies in order to determine any cause-effect relationships between nutrient intake anabolic/catabolic hormone concentrations.

## ACKNOWLEDGEMENTS

The authors thank the players, coaches, and research assistants involved in this investigation for their participation, enthusiasm and cooperation.

*Declaration:* all authors declare that all experiments comply with the current laws of Spain (country in which the experiments were performed).

## REFERENCES

- Rietjens G, Kuipers H, Adam J, Saris W, Van Breda E, Van Hamont Da, et al. Physiological, biochemical and psychological markers of strenuous training-induced fatigue. *Int J Sports Med* 2005;26(01):16-26.
- Córdova A, Martín JF, Reyes E, Álvarez-Mon M. Protection against muscle damage in competitive sports players: The effect of the immunomodulator AM3. *J Sports Sci* 2004;22(9):827-33.
- Engelmann M, Landgraf R, Wotjak CT. The hypothalamic-neurohypophysial system regulates the hypothalamic-pituitary-adrenal axis under stress: An old concept revisited. *Front Neuroendocrinol* 2004;25(3-4):132-49.
- Oltres CM, Mora F, Vives F. Beta-endorphin and ACTH in plasma: Effects of physical and psychological stress. *Life Sci* 1987;40(17):1683-6.
- Hakkinen K, Pakarinen A, Alen M, Kauhanen H, Komi PV. Neuromuscular and hormonal adaptations in athletes to strength training in two years. *J Appl Physiol* (1985) 1988;65(6):2406-12.
- Martinez AC, Seco Calvo J, Tur Mari JA, Abecia Inchaurregui LC, Orella EE, Biescas AP. Testosterone and cortisol changes in professional basketball players through a season competition. *J Strength Cond Res* 2010;24(4):1102-8.
- Schelling X, Calleja-González J, Terrados N. Testosterone and cortisol with relation to training volume and playing time in professional Spanish basketballers. *Med Sci Sport Exer* 2011;43(5 Suppl).
- Hoffman JR, Ratamess NA, Kang J, Falvo MJ, Faigenbaum AD. Effects of protein supplementation on muscular performance and resting hormonal changes in college football players. *J Sports Sci Med* 2007;6(1):85.
- Viru A, Viru M. Cortisol-essential adaptation hormone in exercise. *Int J Sports Med* 2004;25(06):461-4.
- Kraemer WJ, Ratamess NA. Hormonal responses and adaptations to resistance exercise and training. *Sports Med* 2005;35(4):339-61.
- Griggs RC, Kingston W, Jozefowicz RF, Herr BE, Forbes G, Halliday D. Effect of testosterone on muscle mass and muscle protein synthesis. *J Appl Physiol* 1989;66(1):498-503.
- Volek JS, Kraemer WJ, Bush JA, Incledon T, Boetes M. Testosterone and cortisol in relationship to dietary nutrients and resistance exercise. *J Appl Physiol* (1985) 1997;82(1):49-54.
- Hjalmarson A, Aasebø U, Aakvaag A, Jorde R. Sex hormone responses in healthy men and male patients with chronic obstructive pulmonary disease during an oral glucose load. *Scan J Clin Lab Invest* 1996;56(7):635-40.
- Ivandić A, Prpić-Križevac I, Jakić M, Bačun T. Changes in sex hormones during an oral glucose tolerance test in healthy premenopausal women. *Fertil Steril* 1999;71(2):268-73.
- Volek JS. Influence of nutrition on responses to resistance training. *Med Sci Sports Exerc* 2004;36(4):689-96.
- Volek JS, Love DM, Avery NG, Sharman M, Kraemer WJ. Effects of a high-fat diet on postabsorptive and postprandial testosterone responses to a fat-rich meal. *Metab Clin Exp* 2001;50(11):1351-5.
- Venkatraman JT, Feng X, Pendergast D. Effects of dietary fat and endurance exercise on plasma cortisol, prostaglandin E2, interferon-gamma and lipid peroxides in runners. *J Am Coll Nutr* 2001;20(5):529-36.
- Schroeder ET, Villanueva M, West DD, Phillips SM. Are acute post-resistance exercise increases in testosterone, growth hormone, and IGF-1 necessary to stimulate skeletal muscle anabolism and hypertrophy? *Med Sci Sports Exerc* 2013;45(11):2044-51.
- Dorgan JF, Judd JT, Longcope C, Brown C, Schatzkin A, Clevidence BA, et al. Effects of dietary fat and fiber on plasma and urine androgens and estrogens in men: A controlled feeding study. *Am J Clin Nutr* 1996;64(6):850-5.
- Lane AR, Duke JW, Hackney AC. Influence of dietary carbohydrate intake on the free testosterone: Cortisol ratio responses to short-term intensive exercise training. *Eur J Appl Physiol* 2010;108(6):1125-31.
- Hoffman JR, Ratamess NA, Kang J, Falvo MJ, Faigenbaum AD. Effect of protein intake on strength, body composition and endocrine changes in strength/power athletes. *J Int Soc Sports Nutr* 2006;3(2):12-8.
- Lambert CP, Frank LL, Evans WJ. Macronutrient considerations for the sport of bodybuilding. *Sports Med* 2004;34(5):317-27.
- Gamble P. Periodization of training for team sports athletes. *Strength Condition J* 2006;28(5):56-66.
- González-Rave JM, Arijia A, Clemente-Suárez V. Seasonal changes in jump performance and body composition in women volleyball players. *J Strength Cond Res* 2011;25(6):1492-501.
- Vermeulen A, Verdonck L, Kaufman JM. A critical evaluation of simple methods for the estimation of free testosterone in serum. *J Clin Endocrinol Metab* 1999;84(10):3666-72.
- Martín-Moreno JM, Boyle P, Gorgojo L, Maisonneuve P, Fernández-Rodríguez JC, Salvini S, et al. Development and validation of a food frequency questionnaire in Spain. *Int J Epidemiol* 1993;22(3):512-9.
- Mielgo-Ayuso J, Zourdos MC, Calleja-González J, Urdampilleta A, Ostojic S. Dietary intake habits and controlled training on body composition and strength in elite female volleyball players during the season. *Appl Physiol Nutr Metab* 2015;40(8):827-34.
- Farrán A, Zamora R, Cervera P. Tablas de composición de alimentos del Centre d'Ensenyament Superior de Nutrició i Dietètica (CESNID). Barcelona: Universitat de Barcelona; 2004.
- Kreider RB, Wilborn CD, Taylor L, Campbell B, Almada AL, Collins R, et al. ISSN exercise & sport nutrition review: Research & recommendations. *J Int Soc Sports Nutr* 2010;7(7):2-43.
- Phillips SM, Van Loon LJ. Dietary protein for athletes: From requirements to optimum adaptation. *J Sports Sci* 2011;29(Suppl 1):S29-S38.
- Rodríguez NR, Di Marco NM, Langley S. American College of Sports Medicine position stand. Nutrition and athletic performance. *Med Sci Sports Exerc* 2009;41(3):709-31.
- Stewart A, Marfell-Jones M, Olds T, De Ridder H. International standards for anthropometric assessment. Lower Hutt, New Zealand: ISAK; 2011.
- Lee RC, Wang Z, Heo M, Ross R, Janssen I, Heymsfield SB. Total-body skeletal muscle mass: Development and cross-validation of anthropometric prediction models. *Am J Clin Nutr* 2000;72(3):796-803.
- Mielgo-Ayuso J, Calleja-González J, Clemente-Suárez VJ, Zourdos MC. Influence of anthropometric profile on physical performance in elite female volleyballers in relation to playing position. *Nutr Hosp* 2014;31(2):849-57.
- Crewther B, Keogh J, Cronin J, Cook C. Possible stimuli for strength and power adaptation. *Sports Med* 2006;36(3):215-38.
- Bird SP, Tarpenning KM, Marino FE. Independent and combined effects of liquid carbohydrate/essential amino acid ingestion on hormonal and muscular adaptations following resistance training in untrained men. *Eur J Appl Physiol* 2006;97(2):225-38.
- Alleman Jr RJ, Bloomer RJ. Hormonal response to lipid and carbohydrate meals during the acute postprandial period. *J Int Soc Sports Nutr* 2011;8(1):1-8.
- Martens MJ, Rutters F, Lemmens SG, Born JM, Westerterp-Plantenga MS. Effects of single macronutrients on serum cortisol concentrations in normal weight men. *Physiol Behav* 2010;101(5):563-7.
- Gerich JE. Control of glycaemia. *Baillieres Clin Endocrinol Metab* 1993;7(3):551-86.