



Trabajo Original

Valoración nutricional

The status of clinical nutrition in hospitals of Guilan province, Iran – A mixed-method study

*El estado de la nutrición clínica en los hospitales de la provincia de Guilan, Irán:
estudio de método mixto*

Samaneh Mirzaei Dahka¹, Soroor Fathi², Zahra Salimi³, Zahra Roumi⁴, Bahareh Aminnejad⁵, Neda Valisoltani⁶, Hanieh Shafaei⁷, Maryam Gholamalizadeh⁸, Mahdi Shadnough⁹, Saeid Doaei¹⁰, Marjan Ajami¹¹

¹Student Research Committee. Guilan University of Medical Sciences. Rasht, Iran. ²Department of Nutrition. Faculty of Allied Medical Sciences. Ahvaz Jundishapur University of Medical Sciences. Isfahan, Iran. ³Nutrition and Metabolic Diseases Research Center. Ahvaz Jundishapur University of Medical Sciences. Ahvaz, Iran. ⁴Department of Nutrition, Electronic Health and Statistics Surveillance Research Center. Science and Research Branch. Islamic Azad University. Tehran, Iran. ⁵Department of Nutrition. Science and Research Branch. Islamic Azad University. Tehran, Iran. ⁶MSC student. Department of Clinical Nutrition. School of Nutrition Science and Dietetics. Tehran University of Medical Sciences. Tehran, Iran. ⁷Nursing and Midwifery School. Guilan University of Medical Sciences. Rasht, Iran. ⁸Cancer Research Center. Shahid Beheshti University of Medical Sciences. Tehran, Iran. ⁹Department of Clinical Nutrition and Dietetics. Faculty of Nutrition and Food Technology. National Nutrition and Food Technology Research Institute. Shahid Beheshti University of Medical Sciences. Tehran, Iran. ¹⁰Department of Community Nutrition. Faculty of Nutrition and Food Technology. National Nutrition and Food Technology Research Institute. Shahid Beheshti University of Medical Sciences. Tehran, Iran. ¹¹Department of Food and Nutrition Policy and Planning. National Nutrition and Food Technology Research Institute. School of Nutrition Sciences and Food Technology. Shahid Beheshti University of Medical Sciences. Tehran, Iran

Abstract

Background: nutritional therapy has an important role in the development of medical care services, and quantitative and qualitative assessments of the status of clinical nutrition in hospitals is crucial. This study aimed to explore the current status of clinical nutrition in hospitals of Guilan province, Iran.

Methods: this quantitative-qualitative (mixed method) study was performed on public hospitals in Guilan province, Iran (n = 26). The required information was collected by visiting the hospitals and interviewing with hospital dietitians. The data collection tool had two quantitative and qualitative subscales. The quantitative subscale included a 21-item checklist containing information about the referral system, cooperation of hospital staff in clinical nutrition, clinical nutrition staff status, enteral nutrition, and parenteral nutrition.

Results: in more than 55 % of hospitals the patients were not adequately referred to a dietitian. In about 31.2 % of hospitals, cooperation of other departments in the field of clinical nutrition was insufficient. In 37 % of hospitals with intensive care unit (ICU), enteral nutrition was not provided properly. Only 27 % of the hospital properly provided parenteral nutrition for their patients. The most important problems mentioned by hospital dietitians included noncompliance of the food services employers with recommended diets, insufficient number of hospital dietitians, and lack of per case payment method for dietitians. Coverage of diet services by health insurance was the most common suggestion of dietitians to promote diet therapy in public hospitals.

Conclusion: the situation of providing nutritional services to patients in public hospitals is not favorable in terms of quantity and quality in Guilan province, Iran. Designing the necessary policies and reorient the clinical nutrition system in hospitals to promote patients' health and accelerate patient recovery is warranted.

Keywords:

Diet therapy. Hospital. Dietitian. Nutrition care service.

Received: 17/05/2022 • Accepted: 19/09/2022

Funding: funding for this study was provided by cancer research center, Shahid-Beheshti University of Medical Sciences, Tehran, Iran.

Authors' contributions: SM, SD, MSH, ZR and MGH designed the study, and carried out the data collection. HSH, MA and SD were involved in the design of the study, analysis of the data, and critically reviewed the manuscript. This study was conducted at the Shahid-Beheshti University of Medical Sciences, Tehran, Iran.

Consent for publication: institutional consent forms were used in this study.

Conflict of interest statement: the authors declare that they have no competing interests.

Mirzaei Dahka S, Fathi S, Salimi Z, Roumi Z, Aminnejad B, Valisoltani N, Shafaei H, Gholamalizadeh M, Shadnough M, Doaei S, Ajami M. The status of clinical nutrition in hospitals of Guilan province, Iran – A mixed-method study. *Nutr Hosp* 2023;40(1):88-95

DOI: <http://dx.doi.org/10.20960/nh.04239>

Correspondence:

Saeid Doaei and Marjan Ajami. National Nutrition and Food Technology Research Institute. Shahid Beheshti University of Medical Sciences. Tehran, Iran
e-mails: doaei@gums.ac.ir; Ajami789@yahoo.com

Resumen

Antecedentes: la terapia nutricional tiene un papel importante en el desarrollo de los servicios de atención médica, y las evaluaciones cuantitativas y cualitativas del estado de la nutrición clínica en los hospitales son cruciales. Este estudio tuvo como objetivo explorar el estado actual de la nutrición clínica en los hospitales de la provincia de Guilan, Irán.

Métodos: este estudio cuantitativo-cualitativo (método mixto) se realizó en hospitales públicos de la provincia de Guilan, Irán (n = 26). La información requerida se recopiló visitando los hospitales y entrevistando a los dietistas de los hospitales. La herramienta de recolección de datos tenía dos subescalas, cuantitativa y cualitativa. La subescala cuantitativa incluía una lista de verificación de 21 ítems que contenía información sobre el sistema de derivación, la cooperación del personal del hospital en nutrición clínica, el estado del personal de nutrición clínica, la nutrición enteral y la nutrición parenteral.

Resultados: en más del 40 % de los hospitales, los pacientes no fueron derivados adecuadamente a un dietista. En el 35 % de los hospitales, la cooperación de otros departamentos en el campo de la nutrición clínica fue insuficiente. En el 55 % de los hospitales con unidad de cuidados intensivos (UCI), la nutrición enteral no se brindó adecuadamente. Ninguno de los hospitales proporcionó nutrición parenteral adecuada para los pacientes. Los problemas más importantes mencionados por los dietistas hospitalarios incluyeron el incumplimiento de los programas de alimentos por parte del empleador del sector alimentario, un número insuficiente de dietistas hospitalarios y la falta de pago de honorarios de consulta a los dietistas. La cobertura de los servicios de dietética por parte del seguro de salud fue la sugerencia más común de los dietistas para promover la dietoterapia en los hospitales.

Conclusión: la situación de la prestación de servicios nutricionales a pacientes en hospitales públicos no es favorable en términos de cantidad y calidad en la provincia de Guilan, Irán. Se justifica diseñar las políticas necesarias para reorientar el sistema de nutrición clínica en los hospitales y, en definitiva, promover la salud y acelerar la recuperación de los pacientes.

Palabra clave:

Terapia dietética. Hospital.
Dietista. Servicio de
atención nutricional.

INTRODUCTION

Providing the nutritional needs of hospitalized patients is considered to be a fundamental approach to facilitate recovery from illness (1). Malnutrition, which can be considered as undernutrition in hospitalized patients (2), occurs at a frequency of about 37 % to 82 % in hospitals worldwide, varies according to the investigated population and the definitions for malnutrition used (3). Most studies conducted in Iran reported that the nutritional status of hospitalized patients is not favorable (4,5). Malnutrition is a costly disorder because of delayed recovery, increased length of hospital stay (LOS), and increased readmission rate (6). Poor food intake during hospitalization can lead to increased infections, poor wound healing, cardiac complications, and a higher mortality rate (3).

The most common nutritional risk factors in the hospitalized patients include lack of early detection and treatment of malnutrition, inadequate nutritional screening, prolonged nil by mouth status, missed meals due to medical procedures, poor appetite, food wasting, limited dietary options, and nutritional inadequacy of hospital meals (7). High quality nutritional practices in the hospital such as early nutritional screening, early diet therapy, and monitoring of nutritional status may improve the care processes and health outcomes (8). Assessment of the nutritional status of patients who are hospitalized should be considered as an integral part of the health assessment as soon as the patients are admitted to the hospital and should continue to be monitored throughout their hospitalization (1). Dietary practices need a multidisciplinary team to address a patients' nutritional needs and improve their health status and minimize hospital acquired risks associated with poor nutrition (9).

Dietitians are a professional group who are educated to provide individualized nutritional assessment and treatment in hospital settings. A physician referral is commonly required to involve nutrition services in the care of a patient. However, the dietitian's clinical roles are sometimes underestimated by

physicians in different situations, resulting in potentially inappropriate referrals and inadequate nutritional support (10). Many studies reported that early involvement of a dietitian can reduce LOS, decrease infection, and promote healing (11). For example, a recent study indicated that enteral nutrition, which is frequently used in critically ill patients, was more effective when the formula was prescribed by a dietitian compared by a physician. The result showed that dietitian-supervised patients received 10 % more energy and protein, and had an increased body weight and a shorter LOS (12). Another study reported that parenteral nutrition prescription provided by a dietitian compared with a surgeon resulted in a higher intake of vitamin B complex and lipids, and better patient clinical parameters such as fewer secondary infections, less weakness, rapid recovery, and reduced patient mortality (11). Therefore, considering the views of dietitians on the nutritional status of patients and clinical nutrition services provided in the hospital may promote the proper re-orientation of treatment policies. However, few studies have been conducted on the status of clinical nutrition services in hospitals (6). So, this study aimed to assess the status of nutrition care services and diet therapy in public hospitals of Guilan province, Iran.

METHODS

STUDY DESIGN AND DATA COLLECTION

This cross-sectional study was carried out from 1th October 2019 to 1th December 2020 in public hospitals of Guilan province, Iran (n = 26). The public hospitals of five cities including Siahkal, Astara, Talesh, Rudsar, and Fouman were excluded from the study due to lack of data, and three hospitals –Manjil, Rezvanshahr, and Rostamabad– were excluded from the study due to lack of a resident dietitian. Finally,

eighteen hospitals including eight hospitals of Rasht city (Razi, Velayat, Poursina, Heshmat, Alzahra, Shafa, Amiralмомenin, and Hefdah shahrivar), two hospitals of Lahijan city (22-Aban and seyedalshohada), and the hospitals of Amlash, Roudbar, Sowmesara, Bandare Anzali, Astaneh, Langeroud, Masaal, and Shaft cities were included in the study. Data was collected using a validated hospital nutrition assessment questionnaire (HNAQ) with 21 items, which was completed by visiting the hospitals and an in-depth interview with the hospitals' dietitians. The HNAQ has four domains including referral system, involvement of hospital staffs in clinical nutrition, the status of clinical nutrition staff, and enteral and parenteral nutrition. The referral system subscale means to refer to a dietitian for nutrition counseling any patient who has one or more of 18 special clinical conditions including the need for enteral and parenteral nutrition, diabetes, hypertension, cardiovascular disease, cancers, burns, stroke, renal failure, hemodialysis, cirrhosis, organ transplants, bed sores, malnutrition, neurologic diseases, abdomen surgery, chronic obstructive pulmonary disease, and head trauma: referral of a patient with malnutrition, and referral of new case patients to the ICU after 24 hours of hospitalization.

The involvement of the hospital staff in clinical nutrition was defined as compliance with the dietary plan, approval of a clinician to implement the diet provided by a dietitian, proper implementation of the diet by the ward nurse, filling specialized nutrition assessment forms, and record nutrition-related information in the hospital information systems (HIS).

Status of clinical nutrition staff describes an adequate number of dietitians as compared to the number of active hospital beds, the effect of regular and principled referral of patients for nutrition counseling in improving a patient's condition and reducing LOS, and improving the position of dietitian in the hospital. Furthermore, the status of parenteral and enteral nutrition was defined according to the items included separated gavage room, providing gavage feeding with a commercial formula, providing gavage feeding with a manual formula, providing supplements needed in enteral nutrition and vitamin and mineral vials in parenteral nutrition. The questionnaire is on a five-point Likert scale from never (0) to always (4). Generally, "never" and "few" were considered as unfavorable situation, and "always" and "high" were considered as desirable situation.

INTERVIEW GUIDE

A semi-structured interview was used to direct the interviews toward the dietitians' experiences about the nutritional status of their hospital by asking open-ended questions to clarify aspects that came up in the interviews. The interview contains two main topic areas included the participants' experience of problems and their helpful suggestions improving for nutritional services of the hospitals. The interview will be transcribed verbatim, deidentified, and integrated with written field notes following each interview.

STATISTICAL ANALYSIS

Descriptive statistics were used to describe the characteristics of the status of nutrition care services and diet therapy in the hospitals. The frequency of nutritional care service problems in four different domains was calculated. All statistical analyses of descriptive statistics were performed with IBM SPSS version 22.

Regarding qualitative data, we transcribed all the interviews, reviewed the transcriptions for accuracy, and anonymized them. We enacted the criterion for credibility through open-ended questioning, spending a prolonged amount of time with the material, and giving a detailed description of the methods. To meet the criterion for transferability, we present detailed and in-depth descriptive data and quote the participants in the text. To meet the criterion for dependability, each transcription was independently read, checked, and coded by two of the authors.

RESULTS

The situation of clinical nutrition in hospitals was assessed in four different domains (Table I). Regarding the referral of patients for receiving nutritional counseling, patients were not referred or were rarely referred to the nutrition specialist in more than 33 % of the hospitals. The average referral of specialists for nutrition counseling, referral of 18 special clinical conditions for nutrition counseling, and referral of patients with malnutrition for nutrition counseling were in a desirable situation only in 18.4 % of hospitals (Fig. 1).

The cooperation of personnel from other parts of the hospital with the hospital dietitian was insufficient in 48.5 % of hospitals. The adherence of employers to a diet was reported as desirable just in 5 percent of the hospitals (Fig. 2).

Regarding artificial nutrition, enteral nutrition was properly provided in about 40 % of hospitals with ICU ward (Fig. 3). Only about 17.7 % of hospitals in Guilan provided proper parenteral nutrition for patients.

In terms of the situation of the nutrition staff at the hospital, the number of dietitians in about one third of hospitals was less than the desirable number (at least one person per 100 active beds) (13).

According to the qualitative interview and scoring of the collected comments, approximately all dietitians confirmed the effect of regular and methodical referral of patients for nutrition counseling in the improvement of patient health status. In none of the studied hospitals, the dietitians were paid with per case payment methods for their counseling sessions (Table II).

The most common problems reported by the dietitian in the hospitals were included non-adherence of the employer of the food services from the dietary plan as lack of payment of counseling fees as per case payment to dietitians (Table III).

On the other hand, suggestions by dietitians for the improvement of the existing situation are presented in table IV. Some major points were related to lack of clearly defined responsibilities, lack of co-operation among all staff groups, and lack of involvement of hospital management.

Table I. Situation of clinical nutrition indices in hospitals of Guilan province, Iran

Evaluation Index	Number (percent)				
	Never	Few	Moderate	High	Always
Referral system					
Referral of specialists for the nutrition counseling	0	6 (33.3 %)	9 (50 %)	1 (5.6 %)	2 (11.1 %)
Referral of 18 diseases for the nutrition counseling	0	6 (33.3 %)	8 (44.4 %)	2 (11.1 %)	2 (11.1 %)
Referral of patients with malnutrition for nutrition counseling	3 (16.7 %)	8 (44.4 %)	4 (22.2 %)	1 (5.6 %)	2 (11.1 %)
Referral of new case patients in ICU, 24 hours after hospitalization for nutrition counseling	0	4 (36.3 %)	3 (27.3 %)	1 (9.1 %)	3 (27.3 %)
Guiding patients after discharge to the clinic for receiving diet	3 (20 %)	7 (46.7 %)	5 (33.3 %)	0	0
Average	3 (16.3 %)	6 (38.8 %)	6 (35.4 %)	2 (6.3 %)	3 (12.1 %)
Cooperation of other wards of the hospital in the clinical nutrition					
Adherence of employer to the diet	2 (11.8 %)	1 (5.9 %)	2 (11.8 %)	5 (29.4 %)	7 (41.2 %)
Approval of dietician proposed diet by the physician	1 (6.7 %)	3 (20 %)	5 (33.3 %)	4 (26.7 %)	2 (13.3 %)
Proper implementation of the proposed diet by the ward nurse	1 (5.9 %)	3 (17.6 %)	4 (23.5 %)	5 (29.4 %)	4 (23.5 %)
Nutrition specialized evaluation forms in patient's file	1 (5.9 %)	1 (5.9 %)	2 (11.8 %)	3 (17.6 %)	10 (58.8 %)
Registration of diet information in hospital HIS system	11 (64.7 %)	2 (11.8 %)	1 (5.9 %)	1 (5.9 %)	2 (11.8 %)
Average	3 (19 %)	2 (12.2 %)	3 (17.3 %)	4 (21.8 %)	5 (29.7 %)
Enteral nutrition					
Availability and utilization from a separate gavage room	16 (88.9 %)	2 (9.1 %)	0	0	0
Methodical presentation of gavage nutrition with commercial formula	4 (22.2 %)	5 (27.8 %)	3 (16.7 %)	1 (5.6 %)	5 (27.8 %)
Methodical presentation of gavage nutrition with manual formula	0	0	5 (33.3 %)	8 (53.3 %)	2 (13.3 %)
Presentation of the required supplements in Enteral nutrition	0	1 (6.3 %)	4 (25 %)	4 (25 %)	7 (43.8 %)
Average	5 (27.8 %)	2 (9.2 %)	3 (18.7 %)	4 (21.2 %)	4 (21.2 %)
Parenteral nutrition					
Availability and utilization from a separate TPN room (clean room)	7 (63.6 %)	0	2 (18.2 %)	0	2 (18.2 %)
Methodical presentation of parenteral nutrition	6 (60 %)	0	3 (30 %)	0	1 (10 %)
Presentation of vitamin and mineral vial in parenteral nutrition	1 (8.3 %)	4 (33.3 %)	2 (16.7 %)	2 (16.7 %)	3 (25 %)
Average	5 (44 %)	1 (9 %)	2 (18 %)	1 (9 %)	2 (18 %)




Figure 1. The status of the referral of patients for nutrition counseling in the hospitals of Guilan, Iran.




Figure 2.
The status of cooperation of other hospital sections with dietitians in the hospitals of Guilan, Iran.




Figure 3.
The status of proper preparation of enteral feeding at the hospitals of Guilan, Iran.

Table II. Situation of nutrition personnel in Guilan province hospitals, Iran

Evaluation Index	Number (percent)						
	Never	Few	Moderate	High	Always	Not applicable	Unclear
Does the counseling given at the wards affect your received salary?	17 (95 %)	1 (5 %)	0	0	0	0	0
A sufficient number of dietitians compared to the number of active beds in the hospital	2 (11 %)	4 (22 %)	4 (22 %)	1 (5 %)	6 (34 %)	0	1 (5 %)
Effect of regular and methodical referral of patients for nutrition counseling on the improvement of patient status and a decrease in hospitalization period	0	0	1 (5 %)	12 (67 %)	5 (28 %)	0	0 (0 %)
Effect of regular and methodical referral of patients for nutrition counseling on the improvement of the position of dietitian in a hospital	0	0	0	9 (50 %)	7 (39 %)	0	2 (11 %)

Table III. Hospital dietitians' perceptions of the problems existing in the clinical nutrition department of public hospitals in Guilan province, Iran

Theme	Subtheme
Organizational support	a) Lack of support of insurance companies from diet therapy services b) Since the dietitian does not receive any fee for the counseling and preparation of diet, there is not sufficient incentive for more activity in this field c) Some physicians assume low priority for the service of a dietitian
Educational role	a) Lack of identification of patients with malnutrition by the ward personnel to inform the matter to the dietitian b) Sufficient training for the personnel of hospitalization wards for the proper identification of the patients suffering from malnutrition
Management	a) Utilization of manual gavage for the high cost of the formula of commercial gavage b) The company in charge of food distribution in this center, due to claims from the University of Medical Sciences, could not observe the contract c) Unavailability of nutrition clinic for the visit of outpatients d) Imposing non-nutritional tasks to dietitians (medical documents, staff department, IT...)
Interdisciplinary interaction	a) There is an active HIS system, but the ward nurse does not use it

Table IV. Suggestions proposed in clinical nutrition by dietitians in hospitals of Guilan province, Iran

1. Coverage of diet services by health insurance
2. Holding meetings on the importance of nutrition counseling on the improvement process with attendance of hospital physicians, dietitians, and authorities in the healthcare center as well as the province, notification of instructions and related books to the physicians
3. There should be an order about nutrition for each patient for registration of the quality and quantity of food taken by the patient at each meal
4. The emphasis of supervisors of different wards on the nutritional referral of patients
5. A separate dietitian is better for the clinical nutrition department
6. Amendment of the contract between hospital and food distribution contractor
7. Conformity of patient's received food with the diet advised by the therapist
8. Provision of required space and appropriate anthropometry equipment for the evaluation of nutrition status and preparation of a suitable diet
9. Electronic referral by the physician and registration of requested counseling
10. Informing specialist physicians through the director, and communication with the concerned physicians
11. Communication with physicians and identification of 18 diseases requiring nutrition counseling
12. Activation of an automation system (kitchen of nutrition office and hospitalization wards) for food registration
13. Catering manufacture and design according to HACCP rules
14. Making obligatory the design of gavage regimes for patients by dietitians and informing the physicians

DISCUSSION

The main aim of the present study was to explore the status of nutrition care services and diet therapy in public hospitals of Guilan province, Iran. The results of this study indicated that almost 50 % of patients were not referred for further assessment by dietitians in public hospitals in Guilan, Iran. Some studies reported that hospitalized patients in developing countries are at a high risk of malnutrition (14), and this situation can be associated with higher rates of some complications such as increased nosocomial infection and mortality (15).

The patients at risk of malnutrition are supposed to be referred to specialists for nutritional assessment and intervention. Nu-

tritional assessment usually includes anthropometric, dietary, and biochemical measurements and medical history (16). A cross-sectional description showed that nutrition counseling and diet therapy in patients reduced the prevalence of malnutrition and improved weight gain (17). So, comprehensive nutritional assessment by a dietitian should be performed in those patients identified as at risk of malnutrition or who are malnourished using many items for assessing clinical status such as clinical history, physical examination, body composition measurements, and functional and laboratory assessments. The most common reasons given for lack of dietary practices were reported to be insufficient knowledge, low priority, unclear assignment of responsibility, and lack of procedures or guidelines (11).

The results of the present study showed that the shortage of facilities for managing food service operations, such as parenteral nutrition and enteral nutrition (PN/EN) preparation, is a common problem in hospitals of Guilan province, Iran. One of the main roles of the nutrition department in hospitals is involving in PN/EN preparation (18). PN/EN assessments should involve medical, nutritional, and pharmacological considerations as well as physical examination, anthropometric measurement, and laboratory data (11), which are mainly dependent on the specialized skills associated with dietitians (18).

According to the results of this study, in 31.2 % of the hospitals, the cooperation of other departments in the field of clinical nutrition was insufficient. Clinical dietitians help hospital patients to determine their nutritional needs in interaction with another healthcare worker such as physicians and nurses (19). This interaction helps to develop appropriate nutritional programming for patients and track their progress (20). Clinical dietitians may also review laboratory tests ordered by physicians (11). Therefore, inter-departmental cooperation is crucial for the optimal performance of the duties of a dietitian.

In this study, the number of dietitians in the hospitals was below international standards. Pan et al. investigated the status of the clinical department in Jiangsu, China (21) and reported that the nutrition departments of Jiangsu had some problems in different aspects. Nutrition departments did not have a clear department definition and the management and development of nutrition departments were confusing. Also, the ratio of dietitians per hospital bed was far from the standard levels of the hospitals in Jiangsu, China (18). The departments of nutrition in hospitals need enough human resources to improve the provision of nutrition services in hospitals. This study indicated that, compared to a similar study in 2018, the number of total human resources in clinical nutrition departments was increased. However, the most growth of human resources was related to an increased number of clinicians and nurses instead of dietitians (22). Um et al. indicated that the hospitals of Korea had a low performance rate regarding nutritional care and there were large variations in the quality of nutritional service between hospitals (23). Rasmussen et al. found that nutrition departments in Danish hospitals did not follow the standards for nutrition support according to the established criteria (24).

In the present study, lack of organizational support was considered as the most important problem of nutritional services by hospital dietitians.

In Iran, the hospital dietitian is responsible for several duties, such as monitoring the process of receiving and buying foods, monitoring storage, preparing and transporting raw or processed foods, monitoring the preparation of gavage solution, monitoring food cooking, nutritional assessment of the patients and designing the food plan of the patients and staff, and follow up of nutritional status of hospitalized patients. Due to lack of sufficient staff for the number of active beds in the hospital, the multiplicity of responsibilities of dietitians, and the lack of per case payment method (dietitians have a fixed salary), providing a specialized diet for the patients may not be done properly in many hospitals.

Overall, the results of the present study showed that the nutritional status of public hospitals in Guilan province of Iran was not favorable. However, the present study had some limitations. First, this study was conducted only in public hospitals and cannot be generalized to private hospitals. Secondly, some hospitals in different cities of Guilan province were excluded from the study for various reasons, such as lack of access to information and lack of a resident dietitian in the hospital, and this may affect the accuracy and generalizability of the results.

Also, this study focused on the status of providing clinical nutrition services in hospitals, and the nutritional status of the patients has not been studied. Future more comprehensive studies may be effective for better policy-making and order to improve clinical nutrition in hospitals.

CONCLUSION

The status of clinical nutrition in the Guilan public hospitals of Guilan, Iran was inadequate in terms of most of the evaluated indicators. Providing appropriate training to hospital staff, specialists, and nurses, as well as providing a sufficient number of dietitians to hospital, per case payment method for nutrition counseling, of the hospital catering department may improve the situation of clinical nutrition in hospitals.

Ethics approval and consent to participate.

The research was performed in accordance with institutional regulations after protocol review and approval by the Medical Ethics Committee of the Shahid-Beheshti University of Medical Sciences, Tehran, Iran (IR.SBMU.NNFTRI.REC.1400.033).

REFERENCES

- Smith L, Chapman A, Flowers K, Wright K, Chen T, O'connor C, et al. Nutritional screening, assessment and implementation strategies for adults in an Australian acute tertiary hospital: a best practice implementation report. *JBI Database System Rev Implement Rep* 2018;16(1):233-46. DOI: 10.11124/JBISRIIR-2016-003071
- Beavan S, Baker R, Sadler H, Collinson A. Improving the nutritional intake of hospital patients: how far have we come? A re-audit. *J Hum Nutr Diet* 2019;32(3):372-84. DOI: 10.1111/jhn.12607
- Leandro-Merhi VA, Costa CL, Saragiotto L, Aquino JLBd. Nutritional indicators of malnutrition in hospitalized patients. *Arq Gastroenterol* 2019;56:447-50. DOI: 10.1590/S0004-2803.201900000-74
- Amirkalali B, Sharifi F, Fakhrzadeh H, Mirarefin M, Ghaderpanahi M, Larjani B. Evaluation of the mini nutritional assessment in the elderly, Tehran, Iran. *Public Health Nutr* 2010;13(9):1373-9. DOI: 10.1017/S1368980010000303
- Razavi S, Ibrahimpoor M, Kashani A, Jafarian A. Abdominal surgical site infections: incidence and risk factors at an Iranian teaching hospital. *BMC Surgery* 2005;5(1):2. DOI: 10.1186/1471-2482-5-2
- Keller HH, Valaitis R, Laur CV, McNicholl T, Xu Y, Dubin JA, et al. Multi-site implementation of nutrition screening and diagnosis in medical care units: Success of the More-2-Eat project. *Clin Nutr* 2019;38(2):897-905. DOI: 10.1016/j.clnu.2018.02.009
- Reber E, Gomes F, Vasiloglou MF, Schuetz P, Stanga Z. Nutritional risk screening and assessment. *J Clin Med* 2019;8(7):1065. DOI: 10.3390/jcm8071065
- Ruiz AJ, Buitrago G, Rodríguez N, Gómez G, Sulo S, Gómez C, et al. Clinical and economic outcomes associated with malnutrition in hospitalized patients. *Clin Nutr* 2019;38(3):1310-6. DOI: 10.1016/j.clnu.2018.05.016
- Liu G, Zhang S, Mao Z, Wang W, Hu H. Clinical significance of nutritional risk screening for older adult patients with COVID-19. *Eur J Clin Nutr* 2020;74(6):876-83. DOI: 10.1038/s41430-020-0659-7

10. Haile A, Hailu M, Tesfaye E. Prevalence and associated factors of malnutrition among adult hospitalized patients at Amhara National Regional State Referral Hospitals, Ethiopia. *Integr Obes Diabetes* 2015;1(3):80-3. DOI: 10.15761/IOD.1000117
11. Cederholm T, Barazzoni R, Austin P, Ballmer P, Biolo G, Bischoff SC, et al. ESPEN guidelines on definitions and terminology of clinical nutrition. *Clin Nutr* 2017;36(1):49-64.
12. Braga JM, Hunt A, Pope J, Molaison E. Implementation of Dietitian Recommendations for Enteral Nutrition Results in Improved Outcomes. *J Am Diet Assoc* 2006;106(2):281-4. DOI: 10.1016/j.jada.2005.10.039
13. Treatment OoHMaCSEaDo. Rules of the nutrition department in the country's hospitals in order to promote hospital nutrition hotels. Tehran, Iran; 2016. p. 1-113.
14. Edington J, Boorman J, Durrant ER, Perkins A, Giffin CV, James R, et al. Prevalence of malnutrition on admission to four hospitals in England. *Clin Nutr* 2000;19(3):191-5. DOI: 10.1054/clnu.1999.0121
15. Linn BS. Outcomes of older and younger malnourished and well-nourished patients one year after hospitalization. *Am J Clin Nutr* 1984;39(1):66-73. DOI: 10.1093/ajcn/39.1.66
16. Kyle UG, Kossovsky MP, Karsegard VL, Pichard C. Comparison of tools for nutritional assessment and screening at hospital admission: A population study. *Clin Nutr* 2006;25(3):409-17. DOI: 10.1016/j.clnu.2005.11.001
17. O'Flynn J, Peake H, Hickson M, Foster D, Frost G. The prevalence of malnutrition in hospitals can be reduced: Results from three consecutive cross-sectional studies. *Clin Nutr* 2005;24(6):1078-88. DOI: 10.1016/j.clnu.2005.08.012
18. Pan C, Wang J. Current status of clinical nutrition department in tertiary hospitals in Jiangsu. *Ethics, Med Public Health* 2021;16:100620. DOI: 10.1016/j.jemep.2020.100620
19. Ryan AS, Foltz MB, Finn SC. The role of the clinical dietitian: II. Staffing patterns and job functions. *J Am Diet Assoc* 1988;88(6):679-83. DOI: 10.1016/S0002-8223(21)02034-4
20. Mustapha M, Callan J, Champion H, Radbone L. Making the best use of a hospital Dietitian. *Paediatr Child Health* 2013;23(8):342-5. DOI: 10.1016/j.paed.2013.04.010
21. Skipper A, Young M, Rotman N, Nagl H. Physicians' implementation of dietitians' recommendations: a study of the effectiveness of dietitians. *J Am Diet Assoc* 1994;94(1):45-9. DOI: 10.1016/0002-8223(94)92039-7
22. Wang J, Pan C, Ma X. Assessment of the Quality Management System for Clinical Nutrition in Jiangsu: Survey Study. *JMIR Form Res* 2021;5(9):e27285. DOI: 10.2196/27285
23. Um MH, Park YK, Lee SM, Lee SM, Lee E, Cha JA, et al. Clinical nutrition service in Korean tertiary hospitals and general hospitals: result of nationwide cross-sectional survey. *Journal of the Korean Dietetic Association* 2014;20(3):183-98. DOI: 10.14373/JKDA.2014.20.3.183
24. Rasmussen HH, Kondrup J, Ladefoged K, Staun M. Clinical nutrition in Danish hospitals: a questionnaire-based nutrition among doctors and nurses. *Clin Nutr* 1999;18(3):153-8. DOI: 10.1016/S0261-5614(99)80005-3