

Revista Española de Nutrición Humana y Dietética

Spanish Journal of Human Nutrition and Dietetics



CrossMark
click for updates

www.renhyd.org



RESEARCH ARTICLE

Peak height velocity and muscle mass in young soccer players

Juan Diego Hernández Camacho^{a,b}, Ana Belen Huelva Leal^b, José Miguel Martínez-Sanz^{c,d,*},
María Dolores Lahoz Ruano^b, Javier Vázquez Carrión^b

^a Universidad Pablo de Olavide, Sevilla, España.

^b Unidad de Optimización del Rendimiento, Prevención y Recuperación de Lesiones, C.D. Agrupación Deportiva Nervión, Sevilla, España.

^c Departamento de Enfermería, Facultad de Ciencias de la Salud, Universidad de Alicante, San Vicente del Raspeig, España.

^d Gabinete de Alimentación y Nutrición, Universidad de Alicante (ALINUA), San Vicente del Raspeig, España.

*josemiguel.ms@ua.es

Assigned Editors: Fernanda de Souza Teixeira, Eduard Baladia. Comité Editorial de la Revista Española de Nutrición Humana y Dietética. Pamplona, España.

Received: 28/09/2017; accepted: 30/08/2018; published: 28/09/2018.

Peak height velocity and muscle mass in young soccer players

KEYWORDS

Growth and Development;
Adolescent;
Sports;
Soccer;
Anthropometry.

ABSTRACT

Introduction: It has been reported that peak height velocity could be an important period in body development in athletes. The objective was to examine the peak height velocity and maturity offset in young soccer players and analyze anthropometrical measures relate with body development.

Material and Methods: Fifty-eight male young soccer players were studied. They were categorized in three diverse age categories, Under-19 (post-peak height velocity), Under-16 (on peak height velocity) and Under-14 (pre-peak height velocity) years old. Height, sitting height, weight, girths and skinfolds were measured to determine peak height velocity, maturity offset and body components. Descriptive statistics means (Standard Deviation) were calculated. One-way ANOVA analyses and Pearson correlations were determined. The level of significance was set at $p < 0.05$.

Results: The mean height and weight values were 167.63 (10.52) cm and 60.12 (12.43) kg, the medium fat mass was 13.49 (3.65)%. The average muscular transverse areas for arm, thigh and calf were 41.70 (10.82) cm², 154.87 (38.02) cm² and 85.76 (17.67) cm² respectively. The average PHV was 13.97 (0.53) years and the median maturity offset was 1.00 (1.92) years. Significant differences were found among the three age categories analyzed for the anthropometrical elements and for the maturity offset. Significant correlations between maturity offset and anthropometrical components were obtained.

Conclusions: In agreement with previous studies, increases on anthropometrical components and muscle areas were reported after the peak velocity. Consequently, peak height velocity and maturity offset should be considered in young soccer players' management.



PALABRAS CLAVE

Crecimiento y Desarrollo;
Adolescente;
Deportes;
Fútbol;
Antropometría.

Pico de crecimiento y masa muscular en jugadores jóvenes de fútbol

RESUMEN

Introducción: Se ha observado que el pico de crecimiento podría ser un período importante en el desarrollo de deportistas. Objetivo: Determinar el pico de crecimiento y el estado madurativo en jugadores jóvenes de fútbol y analizar medidas antropométricas relacionadas con el desarrollo corporal.

Material y Métodos: Cincuenta y ocho jugadores se clasificaron en 3 categorías de edad diferentes, Sub-19 (post-pico de crecimiento), Sub-16 (en el pico) y Sub-14 (pre-pico). Se midieron la altura, la altura sentada, el peso, perímetros y pliegues para determinar el pico de crecimiento, el estado madurativo y los componentes corporales. Métodos estadísticos descriptivos (Desviación Estándar), análisis ANOVA de una vía y correlaciones de Pearson calculados. El nivel de significación se fijó en $p < 0,05$.

Resultados: La altura y el peso medios fueron de 167,63 (10,52) cm y 60,12 (12,43) kg, la grasa corporal media fue 13,49 (3,65)%. La masa muscular transversal del brazo, muslo y pantorrilla fueron 41,70 (10,82) cm², 154,87 (38,02) cm² y 85,76 (17,67) cm². El pico de crecimiento y el estado madurativo medios fueron 13,97 (0,53) y 1,00 (1,92) años. Se encontraron diferencias significativas entre las categorías analizadas para los parámetros antropométricos y el estado madurativo. Se obtuvieron correlaciones entre el estado madurativo y los componentes antropométricos.

Conclusiones: Se reportaron incrementos en componentes antropométricos y en las áreas musculares tras el pico de crecimiento. Consecuentemente, el pico de crecimiento y el estado madurativos deberían ser tenidos en cuenta en el manejo de deportistas jóvenes.

CITATION

Hernández Camacho JD, Huelva Leal AB, Martínez-Sanz JM, Lahoz Ruano MD, Vázquez Carrión J. Peak height velocity and muscle mass in young soccer players. Rev Esp Nutr Hum Diet. 2018; 22(3): 219-26. doi: 10.14306/renhyd.22.3.468

INTRODUCTION

From long time ago, it is well known that development state could be different in teenagers with same chronological age. Consequently maturity assessment is needed to be taken in consideration in adolescent sport stratification in order to equalize competition and in the management of body development in athletes. The peak height velocity (PHV) is a common method used to examine the maturational state in adolescent body development¹. In order to determine this PHV an equation based on gender, age and anthropometric components has been previously validated². Maturity offset equation was a noninvasive and reliable method to determine the biological age in teenagers. Furthermore, these equations have presented a correlation with skeletal age offset, showing a maturational commonality between

these two methods. Chae *et al.*³ described that the age at PHV differs in function of gender and the height velocity growth was higher in boys than in girls. Additionally, it has been mentioned⁴ that men reach their PHV later than women.

A previous study⁵ investigated the influence of several factors (relative age, anthropometry and fitness) in the career attainment outcomes in 580 junior league rugby players. These elements could have a real impact in the career attainment of these rugby players. Consequently, talent identification programs should pay attention to biological age. Philippaerts *et al.*⁶ analyzed changes in PHV and physical performance in male youth soccer players. Interestingly, many physical sport performances such as balance, speed, muscular endurance, explosive strength, agility, cardiorespiratory capacity and anaerobic endurance

presented a peak evolution during the peak height velocity. Perhaps, these improvements could be connected with changes in body development and composition. Additionally, Buchheit and Mendez-Villanueva⁷ examined the role of maturity status in match running performance, focusing on maximal sprinting and aerobic speeds. More mature soccer players exhibited higher values on match running performance. However, another research⁸ discovered that adolescent growth could affect to vulnerability for injuries in soccer players. Athletes exhibited a higher significant probability to suffer traumatic and overuse injuries during the year of PHV than the year before. Moreover, days missed as consequence of injuries, match and training injury incidences were higher during the PHV year. Maturation and growth are potentiality risk factors for injuries in youth soccer players.

Bidaurrazaga-Letona *et al.*⁹ studied the elements involved in identification and selection of young soccer players. Body size and maturation are important factors to pay attention. Highlighting that identification and promotion of young soccer players by coaches are based on factors which are age-dependent. Another research¹⁰ studied maturity-specific relationships of static/dynamic balance with power and strength analyses in young male soccer players they examine athletes pre- (-3 years to >-1 year from PHV), circa PHV (-1 to +1 year from PHV) and post- (>1 to +3 years from PHV) PHV state. Relations between balance and muscle power/strength were described, these associations were increased with maturity, indicating a possible enhancement in body composition. Supplementary, another study¹¹ analyzed the effect of maturation in selection procedures in basketball. Basketball players reached their PHV at an earlier age than general population, suggesting a possible connection between maturity timing and selection procedures in basketball. Therefore, it seems that PHV could play an essential role in young soccer player's selection, sport performance and injury risk.

The purposes of the present research were to examine the PHV and maturity offset in youth soccer players and analyzed relations between maturity offset values and anthropometric components.

MATERIAL AND METHODS

Subjects

Fifty eight male young soccer players voluntary participated in the current study. Eighteen players were in the Under-14 team (U-14, age 12.54 years), seventeen in Under-16

category (U-16, age 14.49 years) and twenty-three in the Under-19 category (U-19, age 17.24 years). Thus, adolescent athletes at pre-PHV, on PHV and post-PHV were included in the current study. This classification was previously used to study PHV in young soccer players¹². They had trained for ~1.5 hour/day, 4 day/week (including a weekly competitive match) during the previous year. All players were regularly involved in competitive seasons. They delivered informed written consents which had been signed by their parents.

The experimental protocol was written following the ethics rules from Helsinki Declaration. All experimental procedures were in accordance with the Pablo de Olavide University Ethical Committee rules.

Procedures

Anthropometric data were collected by highly trained and experienced technicians certified by the International Society for the Advancement of Kinanthropometry (ISAK) level 1. Height and sitting height were measured with a stadiometer (Seca), to the nearest 0.1cm, and an anthropometrical bench. Weight was determined using an electronic weighing machine (Tanita UM-076) to the nearest 0.1kg. Body perimeters were also collected with a metal tape measure, narrow and inextensible (Cescorf) to the nearest 0.1cm. Skinfolds were taken with a slim guide skinfold caliper, to the nearest 0.5mm. The anthropometric characteristics measured were height, sitting height, weight, three girths (arm relaxed, thigh and calf) and three skin folds (triceps, thigh and calf). Leg length was calculated as height minus sitting height. Data were obtained following the standard techniques from the ISAK¹³⁻¹⁵, given the technical error of measurement (TEM) intra-observer (7.5% to skinfolds and 1.5% to basic measurements, girths and lengths).

Experimental Approach to the Problem

The equation used was:

$$\text{Maturity offset} = -9.236 + 0.0002708 \cdot \text{Leg Length} \\ \text{and Sitting Height interaction} - 0.001663 \cdot \text{Age} \\ \text{and Leg Length interaction} + 0.007216 \cdot \text{Age} \\ \text{and Sitting Height interaction} + 0.02292 \cdot \text{Weight by} \\ \text{Height ratio.}$$

A previous study² used this equation in male teenagers obtaining $R=0.94$, $R^2=0.891$ and $SEE=0.592$. PHV was calculated as the result of the chronological age less the value of maturity offset. Although, anthropometric equations are commonly accepted to predict PHV in adolescents, it has been recently proposed that radiographic-based methods are more accurate and reliable¹⁶. Authors decided

to use anthropometric equations because they did not have access to radiographic technology.

Moreover, fat mass was determined with Slaughter's equation that considers sex, triceps skinfold and calf skinfold¹⁷. It has been recently validated in teenagers¹⁸. Muscle mass was estimated in a cross-sectional way for the arm, the thigh and the calf. Triceps, thigh and calf skinfolds, and arm relaxed, calf and thigh girths were measured to determine muscle transverse areas. This method has been previously used to evaluate muscle mass^{19,20}.

Statistical Analyses

SigmaPlot 12.5 version (Systat software) was used for Statistical Analysis. Descriptive statics (mean [Standard Deviation]) were reported for the different parameters analyzed. Variables were normally distributed. One-way ANOVA analyses were used in order to determine significant differences. Correlations between different variables were examined through the Pearson correlation analysis. The level of significance was set at $p < 0.05$.

RESULTS

The mean height was 167.63 (10.52) cm, the sitting height was 87.36 (6.67) cm, the global value for leg length was 80.27 (4.78). The medium weight in the sample examined was 60.12 (12.43) kg, the mean values for triceps, thigh and calf skinfolds were 9.17 (2.81) mm, 11.70 (4.30) mm and 7.83 (2.74) mm. The values for arm relaxed, thigh and calf girths were 25.57 (3.16) cm, 47.45 (5.07) cm and 35.11 (3.21) cm respectively. In Table 1 can be observed these anthropometrical characteristics for the three different teams examined.

The mean age in the athletes analyzed was 14.97 (2.08) years, the average PHV was 13.97 (0.53) years and the median maturity offset was 1.00 (1.92) years. In Table 1 can be viewed the age, and the maturity for the three age categories analyzed.

The mean fat mass was 13.49 (3.65) %, the average muscular transverse areas for arm, thigh and calf were 41.70 (10.82) cm², 154.87 (38.02) cm² and 85.76 (17.67) cm² respectively.

Table 1. Anthropometrics and maturational characteristics of soccer players analyzed.

Data / Soccer category	U-19 (n=23)	U16 (n=17)	U14 (n=18)	P value
Height (cm)	175.88 ± 5.49	169.83 ± 4.89	155.02 ± 6.64	<0.001
Sitting height (cm)	93.25 ± 3.00	87.94 ± 2.13	79.28 ± 4.13	<0.001
Leg length (cm)	82.62 ± 3.86	81.89 ± 3.95	75.73 ± 3.29*	<0.001
Weight (kg)	71.18 ± 7.09	60.76 ± 4.71	45.37 ± 6.02	<0.001
Triceps skinfold (mm)	9.37 ± 2.76	8.26 ± 2.20	9.77 ± 3.28	0.370
Thigh skinfold (mm)	10.02 ± 3.70	10.88 ± 3.51	14.63 ± 4.35*	0.001
Calf skinfold (mm)	6.87 ± 2.53*	7.60 ± 2.16	9.27 ± 3.00*	0.014
Arm relaxed circumference (cm)	27.93 ± 2.21	26.05 ± 1.93	22.09 ± 1.74	<0.001
Thigh circumference (cm)	51.10 ± 3.78	48.13 ± 3.05	42.14 ± 3.26	<0.001
Calf circumference (cm)	37.55 ± 2.54	35.44 ± 1.53	31.68 ± 1.86	<0.001
Troch-t. laterale (cm)	41.91 ± 3.58	41.79 ± 4.70	36.21 ± 2.25*	<0.001
Tibiale laterale (cm)	36.34 ± 1.70	36.05 ± 1.53	32.86 ± 1.09*	<0.001
Maturity offset (years)	3.01 ± 0.66*	0.76 ± 0.31*	-1.34 ± 0.59*	<0.001
Age (years)	17.24 ± 0.79*	14.49 ± 0.28*	12.54 ± 0.54*	<0.001
PHV (years)	14.22 ± 0.54*	13.73 ± 0.32	13.88 ± 0.56	0.008

Note: Data is reported as mean (standard deviation); **U-14, U-16; U-19:** Under-14, 16, 19 category, respectively; **n:** sample size.

* value was significant different from the rest of the groups at $p < 0.05$.

In Table 2 can be observed these values in the three soccer categories examined.

In Table 3 can be observed the correlations found between the different variables analyzed. It has been obtained significant correlations between the three muscle areas examined and the maturity offset, indicating a positive relation between them. Other correlations between maturity offset and body development points have been reported.

In Figure 1 can be observed increases on the three muscle areas in the three groups analyzed. A non-linear growth in muscular masses can be noticed.

DISCUSSION

The objective of this study was to determine maturity offset values in youth soccer and correlations between maturity offset values, anthropometrical components and muscle transverse areas. Maturity offset values differ in functions of the age category while the PHV tend to be similar in the three teams studied, indicating that one group was in a previous developmental state to the PHV, another team was immediately after the PHV and the another category had reached the PHV some time ago. In accordance with the results of a previous study⁸, players from U-16 category

Table 2. Fat mass and muscle transverse areas of soccer players studied.


Soccer team	Fat mass (%)	Arm muscle transverse area (cm ²)	Thigh muscle transverse area (cm ²)	Calf muscle transverse area (cm ²)
U-14 (n=18)	15.00 (4.22)	28.98 (4.90)*	113.07 (21.46)*	66.16 (8.89)*
U-16 (n=17)	12.66 (2.95)	44.00 (6.03)*	159.74 (20.05)*	87.10 (8.10)*
U-19 (n=23)	12.93 (3.42)	49.96 (7.12)*	183.99 (27.64)*	100.11 (13.02)*
P value	0.143	<0.001	<0.001	<0.001

Note: Data is reported as mean (standard deviation); **U-14, U-16; U-19:** Under-14, 16, 19 category, respectively; **n:** sample size.

* value was significant different from the rest of the groups at p<0.05.

Table 3. Correlations determined.

Variables related	r	P value
Maturity offset (years) / thigh muscle transverse area (cm ²)	0.72	<0.001
Maturity offset (years) / arm muscle transverse area (cm ²)	0.82	<0.001
Maturity offset (years) / calf muscle transverse area (cm ²)	0.79	<0.001
Maturity offset (years) / calf skinfold (mm)	-0.39	0.001
Maturity offset (years) / arm relaxed circumference (cm)	0.79	<0.001
Maturity offset (years) / thigh circumference (cm)	0.56	<0.001
Maturity offset (years) / calf circumference (cm)	0.80	<0.001
Maturity offset (years) / height (cm)	0.89	<0.001
Maturity offset (years) / sitting height (cm)	0.94	<0.001
Maturity offset (years) / leg length (cm)	0.64	<0.001
Maturity offset (years) / weight (kg)	0.92	<0.001
Maturity offset (years) / age (years)	0.96	<0.001

Figure 1. Increases on muscle transverse areas studied.

*value was significant different from the rest of the groups at $p < 0.05$.

could present a potential higher risk of suffering traumatic and overuse injuries than younger athletes because they were in the year of the PHV. Based on the results of this study, the players from this team could possibly benefit of preventive injury training.

Significant differences were observed in almost all anthropometrical elements analyzed in the three categories analyzed. In height, sitting height, leg length, weight, arm, thigh and calf girths the same tendency was found, increasing when the players were older. However, the opposite result was observed in thigh and calf skinfolds, where the younger players exhibited the highest values. Taking all these data in consideration, it seems that these anthropometrical factors could be incremented or decreased in function of having reached the PHV or not. This idea could be supported by the different correlations found between the maturity offset value and some anthropometrical components such as arm, thigh and calf girths, height, sitting height, leg length and weight. Jorquera *et al.*²¹ observed differences in height and weight between U-16 and U-17 categories in players from several soccer teams. In affinity, another study²² investigated anthropometry and performance in soccer players were born during the same year in order to look for differences between older and younger athletes. Older players were taller, had longer legs and presented a higher fat-free mass. The older athletes exhibited a better score performance. Figueiredo *et al.*²³ investigated a possible variation in size connected with changes in biological maturation in two age soccer teams. Athletes developed in maturity were heavier and taller than players retarded in maturation. Another research²⁴ reported that in a sample of soccer players from

two age groups, the most mature athletes were the taller and heavier and also presented the longest legs length. As it was previously proposed²⁵ growth acceleration should be taken into account by coaches when they work with young athletes because more mature soccer players tend to present anthropometrical parameters more developed.

Interestingly, correlations between the maturity offset and the transverse muscle areas of the arm, thigh and calf have been found, indicating that a higher maturity status is possibly connected with a muscle development in the three areas analyzed. It can be noted that muscle transverse areas increase with the age of the soccer players studied. Particularly, this muscle development specially takes place after having reached the PHV, as can be visualized in the significant difference between the U-14 category and the U-16 category in the three transverse muscle areas examined. Interestingly, increases on muscle development did not follow a lineal tendency as can be observed in Figure 1, U-16 players were in the same year of mean PHV showed a peak in the muscle development in three muscular areas examined. Consequently, it could be speculate that PHV is a crucial period in muscle growth in soccer players. In accordance, Ahmad *et al.*²⁶ examined the effect of maturity on quadriceps-to-hamstring strength ratio. They observed that with the advanced of maturity, increases in quadriceps and hamstring muscle strength were reported in adolescents of both genders. Perhaps, these increases on muscular strength could be related with a higher muscle development in mature adolescents. Another recent study¹⁰ determined maturity connections with strength and power measures in young soccer players. They performed strength/power

measures such as back extensor muscle strength, standing long jump, countermovement jump and 3-hop jump tests. They found that post-PHV athletes exhibited better significant results in these tests. It could be possible that post-PHV who presented higher strength/power measures exhibited a greater muscle development in relation with the result obtained in the current study.

It has been showed that nutrition²⁷ and training programs²⁸ play a crucial role in skeletal muscle synthesis. We have observed a potential increase on muscle areas on PHV state, consequently, it would be essential assure to fulfill nutritional requirements for soccer players on PHV period²⁹ in order to maximize this muscle development. Another study³⁰ found that arm muscle area and PHV were related with a performance parameter in rowing. Even the authors proposed that body components that influence time in rowing could be affected by maturity state in adolescent athletes. This idea would be in agreement with the results obtained in this study. The current study constitutes one of the first studies that analyze the PHV, anthropometric parameters and muscle development in young Spanish soccer players.

Previously, it has been proposed that methods to analyze PHV, maturation or age estimation present internal limitations^{31,32}. Malina *et al.* report that the equations are really useful with teenagers close to the time of PHV, although it has limitations in early and late matured teenagers³³⁻³⁵. Their studies show that PHV could be influenced by chronological age, early and late maturation and individual differences^{34,35}.

In addition, a review¹⁶ suggested that radiographic technology would be the most adequate methods to assess PHV and anthropometry methods could overestimated the timing of PHV. Nevertheless, anthropometry methods are more used by clinicians due to their open approachability and easy application. Therefore, PHV evaluation with anthropometric equations could over- or underestimate PHV timing. Consequently, PHV measurements should be contrast with radiographic methods¹⁶.

The studies of Malina *et al.* conclude that equations are useful for average maturing childrens close to the time of peak height velocity, but have limitations with early and late maturing boys and girls³²⁻³⁵.

The present study has limitations such as the range of sex we only examined male soccer players or that all of them played in the same soccer team. Moreover, they were not probabilistic selected. The PHV estimate through anthropometric equation-based methods presents internal limitations as an indicator of maturity timing. Radiographic

technology could reduce the internal limitations to predict PHV.

CONCLUSIONS

Peak height velocity is a notable point in young soccer player's body development when relevant increases take part in anthropometrical components and muscle areas. Consequently, it would be valuable for physical coaches, trainers and medical services of soccer teams to consider PHV and maturity offset value when they work with young athletes. Anthropometric equations are a quick, simple and accessible way to estimate PHV, although this method could present limitations.

COMPETING INTERESTS

Authors state that there are no conflicts of interest in preparing the manuscript.

REFERENCES

- (1) Malina RM, Bouchard C. Growth, maturation, and physical activity. Champaign, Ill, USA: Human Kinetics Books; 1991.
- (2) Mirwald RL, Baxter-Jones ADG, Bailey DA, Beunen GP. An assessment of maturity from anthropometric measurements. *Med Sci Sports Exerc.* 2002; 34(4): 689-94.
- (3) Chae HW, Suh I, Kwon AR, Kim YJ, Kim YH, Kang DR, et al. Longitudinal standards for height and height velocity in Korean children and adolescents: the Kangwha study. [corrected]. *J Korean Med Sci.* 2013; 28(10): 1512-7.
- (4) Iuliano-Burns S, Mirwald RL, Bailey DA. Timing and magnitude of peak height velocity and peak tissue velocities for early, average, and late maturing boys and girls. *Am J Hum Biol.* 2001; 13(1): 1-8.
- (5) Till K, Copley S, Morley D, O'hara J, Chapman C, Cooke C. The influence of age, playing position, anthropometry and fitness on career attainment outcomes in rugby league. *J Sports Sci.* 2016; 34(13): 1240-5.
- (6) Philippaerts RM, Vaeyens R, Janssens M, Van Renterghem B, Matthys D, Craen R, et al. The relationship between peak height velocity and physical performance in youth soccer players. *J Sports Sci.* 2006; 24(3): 221-30.
- (7) Buchheit M, Mendez-Villanueva A. Effects of age, maturity and body dimensions on match running performance in highly trained under-15 soccer players. *J Sports Sci.* 2014; 32(13): 1271-8.

- (8) van der Sluis A, Elferink-Gemser MT, Coelho-e-Silva MJ, Nijboer JA, Brink MS, Visscher C. Sport injuries aligned to peak height velocity in talented pubertal soccer players. *Int J Sports Med.* 2014; 35(4): 351-5.
- (9) Bidaurrazaga-Letona I, Lekue JA, Amado M, Gil SM. Progression in youth soccer: selection and identification in youth soccer players aged 13-15 years. *J Strength Cond Res.* 2017.
- (10) Hammami R, Chaouachi A, Makhlof I, Granacher U, Behm DG. Associations Between Balance and Muscle Strength, Power Performance in Male Youth Athletes of Different Maturity Status. *Pediatr Exerc Sci.* 2016; 28(4): 521-34.
- (11) te Wierike SCM, Elferink-Gemser MT, Tromp EJJ, Vaeyens R, Visscher C. Role of maturity timing in selection procedures and in the specialisation of playing positions in youth basketball. *J Sports Sci.* 2015; 33(4): 337-45.
- (12) Mendez-Villanueva A, Buchheit M, Kuitunen S, Douglas A, Peltola E, Bourdon P. Age-related differences in acceleration, maximum running speed, and repeated-sprint performance in young soccer players. *J Sports Sci.* 2011; 29(5): 477-84.
- (13) Ross WD, Marfell-Jones MJ. *Kinanthropometry*. En: *Physiological testing of the high-performance athlete*. Ottawa: CASS; 1983. p. 75-115.
- (14) Ross WD, De Rose EH, Ward R. *Anthropometry applied to sport medicine*. In: Dirix A, Knuttgen HG, Tittel K, editors. *The Olympic Book of Sports Medicine*. Oxford: Blackwell Scientific Publication, 1988; 233-65.
- (15) Ross WD, Marfell-Jones MJ. *Kinanthropometry*. En: *Physiological testing of the high-performance athlete*. Champaign, Ill: Human Kinetic; 1991. p. 223-308.
- (16) Mills K, Baker D, Pacey V, Wollin M, Drew MK. What is the most accurate and reliable methodological approach for predicting peak height velocity in adolescents? A systematic review. *J Sci Med Sport.* 2017; 20(6): 572-7.
- (17) Wedderkopp N, Froberg K, Hansen HS, Andersen LB. Secular trends in physical fitness and obesity in Danish 9-year-old girls and boys: Odense School Child Study and Danish substudy of the European Youth Heart Study. *Scand J Med Sci Sports.* 2004; 14(3): 150-5.
- (18) Gómez R, De Marco A, de Arruda M, Martínez C, Salazar CM, Valgas C, et al. Predicción de ecuaciones para el porcentaje de grasa a partir de circunferencias corporales en niños pre-púberes. *Nutr Hosp.* 2013; 28(3): 772-8.
- (19) Canda AS. Puntos de corte de diferentes parámetros antropométricos para el diagnóstico de sarcopenia. *Nutr Hosp.* 2015; 32(2): 765-70.
- (20) Anthropometric estimation of muscle mass in top class sportmen. Métodos de estudio de composición corporal en deportistas 8 CSD. Icd Nº 8.
- (21) Jorquera C, Rodríguez F, Torrealba MI, Campos J, Gracia N, Holway F. Características Antropométricas de Futbolistas Profesionales Chilenos. *Int J Morphol.* 2013; 31(2): 609-14.
- (22) Gil SM, Badiola A, Bidaurrazaga-Letona I, Zabala-Lili J, Gravina L, Santos-Concejero J, et al. Relationship between the relative age effect and anthropometry, maturity and performance in young soccer players. *J Sports Sci.* 2014; 32(5): 479-86.
- (23) Figueiredo AJ, Gonçalves CE, Coelho E Silva MJ, Malina RM. Youth soccer players, 11-14 years: maturity, size, function, skill and goal orientation. *Ann Hum Biol.* 2009; 36(1): 60-73.
- (24) Figueiredo AJ, Coelho E Silva MJ, Cumming SP, Malina RM. Size and maturity mismatch in youth soccer players 11- to 14-years-old. *Pediatr Exerc Sci.* 2010; 22(4): 596-612.
- (25) Rubajczyk K, Świerzko K, Rokita A. Doubly Disadvantaged? The Relative Age Effect in Poland's Basketball Players. *J Sports Sci Med.* 2017; 16(2): 280-5.
- (26) Ahmad CS, Clark AM, Heilmann N, Schoeb JS, Gardner TR, Levine WN. Effect of gender and maturity on quadriceps-to-hamstring strength ratio and anterior cruciate ligament laxity. *Am J Sports Med.* 2006; 34(3): 370-4.
- (27) Churchward-Venne TA, Burd NA, Phillips SM. Nutritional regulation of muscle protein synthesis with resistance exercise: strategies to enhance anabolism. *Nutr Metab (Lond).* 2012; 9(1): 40.
- (28) Coratella G, Schena F. Eccentric resistance training increases and retains maximal strength, muscle endurance, and hypertrophy in trained men. *Appl Physiol Nutr Metab.* 2016; 41(11): 1184-9.
- (29) Mujika I, Burke LM. Nutrition in team sports. *Ann Nutr Metab.* 2010; 57(Suppl 2): 26-35.
- (30) Holway FE, Guerci G. Capacidad predictiva de los parámetros antropométricos y de maduración sobre el rendimiento de adolescentes noveles en remo-ergómetro. *Apunts Med Esport.* 2012; 47(175): 99-104.
- (31) Walvoord EC. The timing of puberty: is it changing? Does it matter? *J Adolesc Health.* 2010; 47(5): 433-9.
- (32) Kozielec SM, Malina RM. Modified Maturity Offset Prediction Equations: Validation in Independent Longitudinal Samples of Boys and Girls. *Sports Med.* 2018; 48(1): 221-36.
- (33) Malina RM, Choh AC, Czerwinski SA, Chumlea WC. Validation of Maturity Offset in the Fels Longitudinal Study. *Pediatr Exerc Sci.* 2016; 28(3): 439-55.
- (34) Malina RM, Kozielec SM. Validation of maturity offset in a longitudinal sample of Polish girls. *J Sports Sci.* 2014; 32(14): 1374-82.
- (35) Malina RM, Kozielec SM. Validation of maturity offset in a longitudinal sample of Polish boys. *J Sports Sci.* 2014; 32(5): 424-37.