

## SHORT NOTE

# PROBIOTIC INCREASE THE ANTIRABIES HUMORAL IMMUNE RESPONSE IN BOVINE

## PROBIÓTICO AUMENTA A RESPOSTA IMMUNE HUMORAL ANTI-RÁBICA EM BOVINOS

Arenas, S.E.<sup>1</sup>, L.S.L.S. Reis<sup>2\*</sup>, N.M. Frazatti-Gallina<sup>3</sup>, S.H. Fujimura<sup>3</sup>, H. Bremer-Neto<sup>1</sup> and P.E. Pardo<sup>1</sup>

<sup>1</sup>Universidade do Oeste Paulista. UNOESTE. Presidente Prudente. SP, Brazil.

<sup>2</sup>Universidade Norte do Paraná. Pólo de Apoio Presencial de Assis. UNOPAR VIRTUAL. Assis. SP, Brazil.

<sup>3</sup>Laboratory of Rabies. Instituto Butantan. São Paulo. SP, Brazil.

\*Correspondence: Rodovia Raposo Tavares, km 572. Bairro Limoeiro. CEP 19067-175. Presidente Prudente. SP, Brazil. centraldepesquisaepublicacoes@yahoo.com.br

### ADDITIONAL KEYWORDS

Cattle. Probiotic. Rabies vaccine. Humoral response.

### PALABRAS CHAVE ADICIONAIS

Bovino. Probiótico. Vacina anti-rábica. Resposta humoral.

### SUMMARY

The effect of a probiotic on the immune response of cattle immunized with a dose of rabies vaccine was evaluated in this study. The Nelore bovines (N= 75) were divided randomly into 5 groups: supplemented with 0, 2, 3, 4 and 8 g the probiotic (Proenzime<sup>®</sup>)/animal/day, groups G<sub>0</sub>, G<sub>2</sub>, G<sub>3</sub>, G<sub>4</sub> and G<sub>8</sub> respectively, for 60 days. All the animals were immunized with a single dose of rabies vaccine on the first day of the experiment (day 0). The obtained results showed that there was significant increase (p<0.05) in the titers of antirabies antibodies and frequency of animals immunized in the groups G<sub>4</sub> and G<sub>8</sub> on 30 and 60. In conclusion, the probiotic (Proenzime<sup>®</sup>) increase the humoral immune response of bovine immunized with rabies vaccine.

### RESUMO

Foi avaliado o efeito do probiótico sobre a resposta imune de bovinos imunizados com uma dose de vacina anti-rábica. Utilizou-se bovinos Nelore (N= 75) divididos aleatoriamente em 5 grupos: suplementados com 0, 2, 3, 4 e 8 g probiótico (Proenzime<sup>®</sup>)/bovino/dia, grupos G<sub>0</sub>, G<sub>2</sub>, G<sub>3</sub>, G<sub>4</sub> e G<sub>8</sub>, respectivamente, por 60 dias. Todos os animais foram imunizados com uma dose de vacina anti-rábica no dia zero do experimento. Os resultados obtidos mostraram que houve aumento significativo (p<0,05) nos títulos de anticorpos anti-

rábicos e na frequência de animais imunizados nos grupos G<sub>4</sub> e G<sub>8</sub> no dia 30 e 60. Conclui-se que o probiótico (Proenzime<sup>®</sup>) aumentou a resposta imune humoral em bovinos imunizados com vacina anti-rábica.

### INTRODUCTION

Rabies is an infection caused by a virus of the genus *Lyssavirus*, order Rhabdoviridae. It's a major animal disease in the world because causes fatal encephalitis in mammals, including humans, has wide geographical distribution and constitutes a serious health problem (Consaes *et al.*, 2006). The vaccination is the half most effective and economical instrument for the control of disease (Oliveira *et al.*, 2000).

The use of probiotics is a strategy to increase the immunological response of animals vaccinated or infected by virus or bacteria (Hamilton-Miller, 2004). Havenaar and Huis In't Veld defined probiotics in 1992 as viable microorganisms, including lactic bacteria and yeast, like lyophilized cells or fermented products that have beneficial effects on host health after being ingested. These substances are used as growth pro-

Recibido: 25-2-07. Aceptado: 6-3-07.

Arch. Zootec. 58 (224): 733-736. 2009.

moters (Arenas *et al.*, 2007).

In the present study was evaluated the effect of different concentrations of probiotic added to the mineral diet, in the immunological response of bovine immunized with rabies vaccine.

MATERIALS AND METHODS

Nellore cattle (*Bos taurus indicus*) aged 18 months, from a farm in Presidente Bernardes, SP, Brazil was utilized in this study. These animals were divided randomly into 5 groups (N= 15), animals from the control group (Gc) fed pasture and received mineral salt (Premix R<sup>®</sup>) *ad libitum* and the other groups were fed in a similar system, but each one received 2, 3, 4 or 8 g probiotic (Proenzime<sup>®</sup>) added to 70 g of mineral salt, groups G<sub>2</sub>, G<sub>3</sub>, G<sub>4</sub> and G<sub>8</sub> respectively. This probiotic/salt ratio was chosen because in observations taken for 20 days before the experiments the daily average salt intake was estimated at 70 g per bovine. All the animals were immunized with 1 dose of rabies vaccine on the first day of the experiment (day 0).


The probiotic (Proenzime<sup>®</sup>) is produced by EMBRAUPEC, Paranavaí, PR, Brazil. It is composed of amylase, cellulase, protease, lipase, pectinase, *Lactobacillus acidophilus*, *Streptococcus faecium*, *Bifidobacterium thermophilum*, *Bifidobacterium longum* and zinc.

A commercial rabies vaccine with satisfactory antigenic values contains a suspension of fixed rabies PV (Pasteur virus) obtained in BHK-21 cells, inactivated by beta-propiolactone, adsorbed in aluminum hydroxide and preserved with thimerosal at 1:10000.

On days 0, 30 and 60, blood (10 ml) was collected from the jugular vein in vacuum tubes with no anticoagulant. After the blood samples were clotted and centrifuged 2500 rpm, 10 min, serum samples were stored at -20 °C for further determination of rabies-neutralizing antibodies in BHK-21 cells.

The neutralizing antibodies were determined by serum neutralization in BHK-21 clone 13 cells. This test is based on the Rapid fluorescent focus inhibition test (RFFIT) (Smith *et al.*, 1996) and the fluorescent inhibition microtest (FIMT RFFIT) (Zanlan *et al.*, 1979).

Data were compared by repeated measures analysis of variance; further multiple



**Figure 1.** Mean titers ( $\pm$  standard deviation) of rabies antibodies in first-time-vaccinated cattle receiving feed supplementation with 0 (Gc), 2 (G<sub>2</sub>), 3 (G<sub>3</sub>), 4 (G<sub>4</sub>) or 8 (G<sub>8</sub>) g of probiotic (Proenzime<sup>®</sup>)/animal/day. In a day means followed by the same letter indicate no significant difference among groups ( $p < 0.05$ ). \*indicates significant difference between days 30 and 60 within a same group ( $p < 0.05$ ). (Títulos médios ( $\pm$  desvio padrão) de anticorpos anti-rábicos de bovinos primovacinaados com suplementação alimentar de 0 (Gc), 2 (G<sub>2</sub>), 3 (G<sub>3</sub>), 4 (G<sub>4</sub>) ou 8 (G<sub>8</sub>) g de probiótico (Proenzime<sup>®</sup>)/animal/dia. Médias seguidas por a mesma letra indicam, dentro de um mesmo dia, diferença não significava entre os grupos ( $p < 0,05$ ). \*indica diferença significativa entre dias 30 e 60 dentro de um mesmo grupo ( $p < 0,05$ )).

## PROBIOTIC INCREASE THE ANTIRABIES HUMORAL IMMUNE RESPONSE IN BOVINE

comparisons were carried out by LSD test. In all the analyses, outliers were excluded. To compare the frequency of animals immunized against rabies among the experimental groups as a function of observation day, the Goodman test was used. In all the analyses, probability of error was set at 5% (Zar, 1999).

### RESULTS AND DISCUSSION


The World Health Organization (WHO) recommends rabies-neutralizing antibody titers of at least 0.50 UI/ml for effective prevention in humans against rabies virus contamination. Some studies argue that this neutralizing antibody titer is the minimal level required to protect cattle (Albas *et al.*, 2005).

On day 0, the results from titration of rabies neutralizing antibodies, sampled from the serum of cattle under the different experimental treatments, were not reactive, indicating that none of the animals had either prior contact with the rabies virus or vaccination against rabies. Thus, the variations found among rabies antibody titers were induced by rabies vaccination as well as by the probiotic treatment.

Significant interaction was found between the probiotic level and vaccination time ( $F_{(4,63)} = 2.58$ ;  $p = 0.046$ ). After rabies vaccination, all the cattle had rabies-antibody titers, irrespective of whether or not they had received probiotic added to the mineral mixture. However, animals from groups  $G_4$  and  $G_8$  had a significant increase ( $p < 0.05$ ) in the production of rabies antibodies 30 and 60 days after vaccination (**figure 1**), and also the frequency of immunized animals (antibody titers = 0.50 UI/ml) (**figure 2**) on both observation days. In this way, these supplementations caused a persistent immunostimulant effect. But their action mechanism is not completely known (Silva *et al.*, 2006). These results agree with previous studies in swine (Ávila *et al.*, 1998) and poultry (Kabir *et al.*, 2004)

which indicate that the immune response is improved when probiotic administration is associated with vaccination.

In group  $G_{33}$ , there was a significant increase ( $p < 0.05$ ) in rabies antibody titers on day 30, but on day 60 the titers decreased significantly (**figure 1**). Thus, the immu-



**Figure 2.** Frequency of rabies-immunized cattle. The cattle received daily feed supplementation with 0 ( $G_c$ ), 2 ( $G_2$ ), 3 ( $G_3$ ), 4 ( $G_4$ ) or 8 ( $G_8$ ) g probiotic (Proenzime<sup>®</sup>)/animal/day. Means followed by same upper-case letter indicate that no significant difference was found between the groups ( $p < 0.05$ ) within the same day. Means followed by the same lower-case letter indicate that no significant difference was found between the days ( $p < 0.05$ ) within the same group. (Frequência de bovinos imunizados contra a raiva. Os bovinos receberam suplementação alimentar diária com 0 ( $G_c$ ), 2 ( $G_2$ ), 3 ( $G_3$ ), 4 ( $G_4$ ) ou 8 ( $G_8$ ) g de probiótico (Proenzime<sup>®</sup>)/animal/dia. Médias seguidas por uma mesma letra maiúscula indicam que, dentro de um mesmo dia, diferença não significativa entre os grupos ( $p < 0,05$ ). Médias seguidas por uma mesma letra minúscula indicam que, dentro de um mesmo grupo, diferença não significativa entre os dias ( $p < 0,05$ ).

nostimulant action caused was not persistent.

In group G<sub>2</sub>, there was not a significant increase ( $p < 0.05$ ) in the titers of rabies antibodies (**figure 1**). Moreover, in this group, the frequency of immunized animals decreased significantly ( $p < 0.05$ ) 60 days after vaccination, similarly to group Gc (**figure 2**). The supplementation was not an immunostimulant.

In group control cattle had the protective rabies antibody titers in any of the two observation days, i.e., all the animals were exposed to the risk of rabies contamination after 30 and 60 days of vaccination (**figure 2**). In fact, Albas *et al.* (2005), and Reis *et al.*

(2008) report that cattle which are first-vaccinated against rabies require a booster vaccination to achieve protective titers of rabies anti-bodies.

The results found in the present study lead to the conclusion that the probiotic (Proenzime<sup>®</sup>) increases both the humoral immune response and the frequency of cattle immunized by rabies vaccine.

#### ACKNOWLEDGEMENTS

This research was supported by the EMBRAUPEC - Empresa Brasileira de Aumento de Produtividade Pecuária, Paranavaí, PR, Brazil.

#### REFERENCES

- Albas, A., P.E. Pardo, H. Bremer-Neto, N.M.F. Gallina, R.M. Mourão-Fuches e A. Sartori. 2005. Vacinação anti-rábica em bovinos: comparação de cinco esquemas vacinais. *Arq. Inst. Biol.*, 72: 153-159.
- Arenas, S.E., L.S.L.S. Reis, N.M. Frazatti-Gallina, R. Giuffrida e P.E. Pardo. 2007. Efeito do probiótico Proenzime<sup>®</sup> no ganho de peso em bovinos. *Arch. Zootec.*, 56: 75-78.
- Ávila, F.A., A.C. Paulillo, R.P. Schocken-Iturrino, F.A. Lucas, A. Orgaz and J.L. Quintana. 1998. Use of vaccine and probiotic in the control of swine diarrhea caused by enterotoxigenic *Escherichia coli*. *Arq. Bras. Med. Vet. Zootec.*, 50: 505-511.
- Consales, C.A., C.A. Pereira, E.C. Passos, M.L. Carreiri, N.M.F. Galina and A.O. Sant'Anna. 2006. Lack of correlation between rabies virus replication in the brain and antibody isotype profile in genetically modified mice. *J. Venom. Anim. Toxins. Incl. Trop. Dis.*, 12: 423-434.
- Hamilton-Miller, J.M.T. 2004. Probiotics and prebiotics in the elderly. *Postgrad. Med. J.*, 80: 447-451.
- Havenaar, R. and J.H.J. Huis In't Veld. 1992. Probiotics a general review. In: Wood B. (ed.). *The Lactic Acid Bacteria in Health and Disease*. Barking Elsevier. pp. 151-170.
- Kabir, S.M.L., M.M. Rahman, M.B. Rahman, M.M. Rahman and S.U. Ahmed. 2004. The dynamics of probiotics on growth performance and immune response in broilers. *Poult. Sci.*, 33: 61-364.
- Oliveira, N.A., M.C.R. Andrade, M.V. Silva, W.C. Moura and E.C. Contreras. 2000. Immune response in cattle vaccinated against rabies. *Mem. Inst. Oswaldo Cruz*, 95: 83-88.
- Reis, L.S.L.S., N.M. Frazatti-Gallina, R.L. Paoli, R. Giuffrida, A. Albas, E. Oba and P.E. Pardo. 2008. Efficiency of *Matricaria chamomilla* CH12 and number doses of rabies vaccine on the humoral immune response in cattle. *J. Vet. Sci.*, 9: 435-437.
- Silva, C.A., E.H. Hoshi, G.D.Pacheco e M.V. Briganó. 2006. Avaliação de probióticos (*Pediococcus acidilactici* e *Bacillus subtilis*) após o desmame e efeitos no desempenho dos leitões. *Semina: Ciências Agrárias*, 27: 133-140.
- Smith, J.S., P.A. Yager and G.M. Baer. 1996. A rapid fluorescent focus inhibition test (RFFIT) for determining rabies virus-neutralizing antibody. In: Mestin FX, Kaplan MM, Koprowski H (eds). *Laboratory techniques in rabies*. World Health Organization. Geneva. pp. 181-192.
- Zalan, E., C. Wilson and D.A. Pukitis. 1979. Microtest for the quantitation of rabies virus neutralizing antibodies. *J. Biol. Stand.*, 7: 213-220.
- Zar, J.H. 1999. *Biostatistical Analysis*. 4<sup>th</sup> ed. New Jersey. Prentice Hall. 663 p. + 123 ap.