

CASE REPORT

Hepatic hematoma after ERCP: two new case reports

María-del-Moral-Martínez, Ana Delgado-Maroto, María Eloísa Cervilla-Sáez-de-Tejada, Francisco Javier Casado-Caballero and Francisco Javier Salmerón-Escobar

Clinical Management Unit of Digestive Diseases. Hospital Universitario San Cecilio. Granada, Spain

ABSTRACT

ERCP is an endoscopic procedure with a complication risk ranging from 2.5 to 8%. The most frequent complications are pancreatitis, cholangitis, hemorrhage or perforation. Hepatic hematoma after ERCP is a potentially serious, rare complication. Not many cases are reported in the literature. We present here two new cases of hepatic hematoma following ERCP along with a review of the literature and possible therapeutic options.

Key words: Hepatic hematoma. ERCP. Complication.

INTRODUCTION

The endoscopic retrograde cholangiopancreatography ERCP is an endoscopic procedure with a complication risk ranging from 2.5 to 8%. The most frequent complications are pancreatitis, cholangitis, hemorrhage or perforation. Hepatic hematoma after ERCP is a potentially serious, rare complication. Not many cases are reported in the literature. We present here two new cases of hepatic hematoma following ERCP along with a review of the literature and possible therapeutic options.

CASE REPORT 1

A 37-year-old woman was admitted to hospital, as programmed, to perform an ERCP due to a diagnosis of a common-bile-duct stone the previous week via a magnetic resonance (MR) cholangiography. On admission, she was asymptomatic. The patient had no significant past medical and family history. During the examination, a marked cutaneous-mucous jaundice was observed. Laboratory tests showed: total bilirubin 7.91 mg/dl, direct bilirubin 7.13 mg/dl, aspartate aminotransferase (AST) 366 U/l, alanine aminotransferase (ALT) 407 U/l, GGT 812 U/l, and alkaline phosphatase (ALP) 332 U/l; the rest of the parameters showed no alteration.

An ERCP was performed, revealing a small papilla difficult to cannulate, and gaining access into the pancreatic duct on two occasions. By performing a precut, access was gained into the common bile duct, which was dilated, and an impacted calculus about 2 cm in diameter was found. A wide papillotomy was performed, the calculus was removed with the help of the Fogarty balloon and an outflow of retained bile and pus remains was observed.

Six hours after the ERCP was performed, the patient began to complain of intense diffuse abdominal pain associated with diffuse discomfort. She did not present with fever or hyperthermia. An urgent laboratory test showed amylase at 1,702 U/l and no other alterations. The patient was diagnosed with acute pancreatitis post-ERCP and she was prescribed serum therapy, *nil per os* (NPO), analgesic perfusion and vital signs monitoring. A few hours later, the patient had a presyncopal episode and her diffuse discomfort and intense abdominal pain persisted, observing a blood pressure (BP) of 85/45 mmHg, heart rate (HR) of 120 bpm and SatO₂ of 97% with no O₂ supply, so a new laboratory test was requested in order to evaluate possible complications.

A new laboratory test revealed a marked drop in hemoglobin from 12 to 7.6 g/dl, along with the already known hyperamylasemia; there was no rise in acute phase reactants and no other alterations. The patient presented neither tarry stools nor hematemesis, and no blood exteriorization was observed. In view of this findings, an abdominopelvic computed tomography (CT) scan was requested in order to preclude possible complications. The CT scan revealed two large subcapsular/intraparenchymal hypodense collections, one of them (measuring 70 x 107 x 120 mm) in the left hepatic lobe and the other one affecting the entire right hepatic lobe, compatible with hematomas. A small amount of subphrenic intraparenchymal air in contact with the right hepatic lobe collection was observed. Hepatic

Received: 28-02-2016
Accepted: 27-12-2016

Correspondence: María del Moral Martínez. UGC Digestive Diseases. Hospital Universitario San Cecilio. Calle Dr. Oloriz, 16. 18012 Granada, Spain
e-mail: mdlmoral87@gmail.com

Del-Moral-Martínez M, Delgado-Maroto A, Cervilla-Sáez-de-Tejada ME, Casado-Caballero FJ, Salmerón-Escobar FJ. Hepatic hematoma after ERCP: two new case reports. *Rev Esp Enferm Dig* 2017;109(6):470-473.

DOI: 10.17235/reed.2017.4237/2016

artery and portal vein permeability was noticed, with no contrast extravasation (Fig. 1).

A red blood cell transfusion and intravenous antibiotic therapy were initiated, and the patient was transferred to the Intensive Care Unit for stabilization and monitoring. An urgent arteriography revealed no contrast extravasation. Since the patient was successfully stabilized, afebrile and in good condition, with a stable hemoglobin level, she was managed by close observation.

Two weeks later she remained afebrile and there was no evidence of additional bleeding, so she was discharged from the hospital. The patient attended monthly follow-up visits in the clinic, which included laboratory tests and ultrasound monitoring. No clinical or analytical repercussion was observed and the imaging tests showed a progressive reduction in the size of the injuries. Six months after being discharged, a follow-up abdominal CT scan was performed, which revealed that the two hypodense collections in the subphrenic space remained over the right and the left hepatic lobes, though they had decreased significantly, measuring about 11 x 8.5 x 10.5 cm and 2.8 x 2.6 x 2.3 cm (anteroposterior x lateral x transverse diameters, respectively).

CASE REPORT 2

A 60-year-old woman with no significant past medical history was admitted from the Emergency Department with clinical symptoms suggestive of cholangitis. During her stay, an MR cholangiography was performed, which revealed a retained 7.7 mm calculus in the common bile duct, about 4.4 cm from the papilla, with intra- and extra-

hepatic bile duct dilatation. In view of this finding, we decided to perform an ERCP, but the papilla of Vater could not be cannulated. Two days later, as the jaundice persisted, a new ERCP was performed. A fresh clot covering the papilla was detected. After removing it, access was gained into the bile duct, observing the calculus described during the MR cholangiography, which was removed after performing a biliary sphincterotomy. Finally, diluted epinephrine was injected into the papilla, still with signs of recent bleeding.

Seven days after being discharged from the hospital, the patient presented to the Emergency Department with fever, abdominal pain and intense asthenia. Laboratory tests showed hemoglobin (Hb) 8.6 g/dl, hematocrit (Hct) 27.2%, white blood cells (WBC) 10,000 U/ml with 68% polymorphonuclear, platelets 627,000 U/ml, and C-reactive protein (CRP) 187.4 mg/l. An abdominal ultrasound was performed, during which a hepatic space occupying lesion (SOL) in segment III measuring about 10 x 6 cm with ultrasound signs of an abscess was observed. Percutaneous drainage was attempted, but it was not possible due to the lesion density. A fine-needle biopsy was performed and *Enterobacter cloacae* was isolated in the culture of the obtained material.

A follow-up abdominal CT scan after antibiotic therapy revealed the lesion described above, corresponding to a possible hepatic hematoma (Fig. 2). Since a high fever persisted, an 8-Fr drainage catheter was placed.

After drainage, the patient remained asymptomatic, afebrile and hemodynamically stable. The patient underwent weekly abdominal ultrasound examinations to monitor the hematoma evolution, observing its progressive reduction. Six weeks later, the drainage was removed with no complications.



Fig. 1. Abdominopelvic CT scan (coronal): two subcapsular/intraparenchymal hypodense collections, one of them (measuring 70 x 107 x 120 mm) in the left hepatic lobe, and the other one affecting the entire right hepatic lobe, compatible with hematomas.




Fig. 2. Abdominal CT scan with IV contrast (axial): hypodense collection in hepatic segment III measuring about 10 x 6 cm.

DISCUSSION

ERCP is an endoscopic procedure with a complication rate ranging from 2.5 to 8% and a mortality rate ranging from 0.5 to 1%. The most frequently described complications are acute pancreatitis (1-7%), acute cholangitis (1.4%), hemorrhage (1%) and duodenal perforation (1%) (1). Nevertheless, there are some exceptional complications, such as hepatic hematoma.

Hepatic hematoma after ERCP is a rare complication, the frequency of which is scarcely reported. We have

reviewed the cases reported in the literature, and when entering the terms “hepatic hematoma after ERCP” and “ERCP complications” in PubMed 25 cases reported to date were found (Table I).

Etiopathogenesis is not completely clear. Two hypothesis have been proposed. One of them states that the lesion would be caused by the guide wire usually used to cannulate the common bile duct, which would perforate the bile duct, thus damaging the hepatic parenchyma. As a result of this, blood vessels would break and air would enter the bile duct and the hematoma, which would explain the

Table I. Description of published case reports

<i>Author</i>	<i>Age</i>	<i>Sex</i>	<i>Indication for ERCP</i>	<i>Use of guidewire</i>	<i>Onset</i>	<i>Symptoms</i>	<i>Treatment</i>	<i>ATB</i>
Ortega-Deballon et al., 2000	81	Male	Common bile duct stone	NA	NA	Abdominal pain	Percutaneous drainage	Yes
Horn et al., 2004	88	Female	Pancreatic pseudocyst	Yes	48 h	Abdominal pain/anemia	Conservative	Yes
Chi et al, 2004	43	Female	Pancreatic cancer	Yes	NA	Abdominal pain	Embolization	Yes
Erturk et al., 2006	41	Male	Cholangio-carcinoma	Yes	48 h	Fever/abdominal pain	Conservative	Yes
Priego et al., 2007	30	Female	Obstructive jaundice	NA	NA	Abdominal pain	Surgery	Yes
Petit-Laurent et al., 2007	98	Male	Common bile duct stone	Yes	NA	NA	Percutaneous drainage	NA
Bhati et al., 2007	51	Female	Common bile duct stone	Yes	NA	Pain/hypotension	Percutaneous drainage	NA
McArthur et al., 2008	71	Male	Common bile duct stone	Yes	12 h	Pain/leukocytosis	Conservative	Yes
De la Serna et al., 2008	71	Female	Common bile duct stone	Yes	48 h	Pain/leukocytosis	Conservative	Yes
Cárdenas et al., 2008	54	Female	Bile leak	Yes	24 h	Pain/anemia	Conservative	Yes
Nari et al., 2009	15	Female	Pancreatitis	NA	NA	Fever/anemia	Conservative	Yes
Revuelto et al., 2010	41	Male	Common bile duct stone	NA	6 h	Pain	Conservative	Yes
Baudet et al., 2010	69	Female	Common bile duct stone	Yes	24 h	Pain/fever/anemia	Embolization/surgery	Yes
Pérez-Legaz et al., 2011	72	Female	Common bile duct stone	Yes	2 h	Pain/anemia	Surgery	Yes
Del Pozo et al., 2011	NA	NA	Common bile duct stone	Yes	5 days	Pain/anemia	Conservative	Yes
Orellana et al., 2012	96	Male	Periampullary tumor	Yes	4 h	Pain	Conservative	Yes
	49	Male	Biliary stent exchange		2 h	Pain/hypotension	Embolization	NA
	55	Female	Gallbladder cancer		NA	Pain	Conservative	NA
Oliviera et al., 2013	84	Male	Common bile duct stone	Yes	10 days	Pain/fever	Percutaneous drainage	Yes
Fei et al., 2013	56	Male	Common bile duct stone	Yes	2 h	Fever	Percutaneous drainage	Yes
Carrica et al., 2014	37	Female	Common bile duct stone	Yes	72 h	Pain/fever	Percutaneous drainage	Yes
Klimová et al., 2014	54	Male	Main pancreatic duct stone	Yes	6 h	Anemia/pain/hypotension	Embolization/surgery/percutaneous drainage	Yes
Zizzo et al., 2015	52	Female	Common bile duct stone	Yes	24 h	Pain	Embolization	Yes
González-López et al., 2015	30	Female	Benign strictures after surgery	Yes	72 h	Pain/hemodynamic shock	Surgery	Yes
Servide MJ et al., 2016	83	Male	Common bile duct stone	NA	15 days	Abdominal pain/anemia	Conservative	Yes

NA: Not available.

presence of subphrenic air in our first patient's CT scan. The other hypothesis states that the hepatic damage is secondary to the traction force exerted with the balloon on the bile duct when trying to remove a retained calculus. This force would cause the rupture of bile ductules and vessels, as well as the consequent bleeding (2).

The clinical manifestations are variable, including abdominal pain, anemia, fever and hemodynamic shock. According to the cases reported in the literature, abdominal pain was the first clinical manifestation in 87% of the cases, associated with anemia (29%) and fever (25%). The onset of symptoms can vary as well, ranging from hours to days. The longest period described in a case was fifteen days after ERCP (3).

The treatment must be personalized depending on the case. Those patients clinically and hemodynamically stable can be managed conservatively (3,4,5). Prophylactic antibiotic therapy is usually administered, given the high risk of infection. For those cases presenting with hemodynamical instability or associated complications, such as hematoma breakage, urgent surgical treatment should be considered (6,7,8). There are some case reports in which an arteriography with embolization of the bleeding vessel has successfully resolved the clinical situation (9,10).

REFERENCES

1. García-Cano J, González Martín JA, Morillas Ariño J, et al. Complications of endoscopic retrograde cholangiopancreatography. A study in a small ERCP unit. *Rev Esp Enferm Dig* 2004;96:163-73.
2. Baudet JS, Arguiñarena X, Redondo I, et al. Subcapsular hepatic hematoma: An uncommon complication of endoscopic retrograde cholangiopancreatography. *Gastroenterol Hepatol* 2011;34(2):79-82. DOI: 10.1016/j.gastrohep.2010.11.002
3. Servide MJ, Prieto M, Marquina T. Una rara complicación tardía tras CPRE: Hematoma hepático subcapsular. *Rev Esp Enferm Dig* 2016;108(4):234-5.
4. Orellana F, Irarrazaval J, Galindo J, et al. Subcapsular hepatic hematoma post ERCP: A rare or an underdiagnosed complication? *Endoscopy* 2012;44(Supl 2)UCTN:E108-9. DOI: 10.1055/s-0031-1291493
5. Oliveira-Ferreira A, Tato-Marinho R, Velosa J. Infected hepatic hematoma 10 days after ERCP. *Endoscopy* 2013;45(Supl 2)UCTN:E402-3.
6. González-López R, García-Cano E, Espinosa-González O, et al. Surgical treatment for liver haematoma following endoscopic retrograde cholangiopancreatography. An unusual case. *Cir Cir* 2015;83(6):506-9. DOI: 10.1016/j.circir.2015.05.028
7. Pérez-Legaz J, Santos J, Ruiz-Tovar J, et al. Subcapsular hepatic hematoma after ERCP (endoscopic retrograde cholangiopancreatography). *Rev Esp Enferm Dig* 2011;103(10):550-1. DOI: 10.4321/S1130-01082011001000011
8. Priego P, Rodríguez G, Mena A, et al. Subcapsular liver hematoma after ERCP. *Rev Esp Enferm Dig* 2007;99:53-4.
9. Klimová K, Padilla-Suárez C, González-Asanza C, et al. Subcapsular hepatic hematoma after ERCP: A case report and revision of literature. *Clin Med* 2014;03(03):161-6. DOI: 10.4236/crcm.2014.33039
10. Chi KD, Waxman I. Subcapsular hepatic hematoma after guide wire injury during endoscopic retrograde cholangiopancreatography: Management and review. *Endoscopy* 2004;36(11):1019-21. DOI: 10.1055/s-2004-825861