

## Revisión

# Chemical composition, antioxidant capacity and content of phenolic compounds in meals collected in hospitals in Bolivia and Sweden

L. Tejada<sup>1,2</sup>, M. Dębiec<sup>1,3</sup>, L. Nilsson<sup>2</sup>, J. M. Peñarrieta<sup>1</sup> and J. A. Alvarado<sup>1</sup>

<sup>1</sup>School of Chemistry. San Andrés University. La Paz, Bolivia. <sup>2</sup>Food Colloids Group. Dept. of Food Technology, Engineering and Nutrition. Faculty of Engineering LTH. Lund University. Lund, Sweden. <sup>3</sup>Pure and Applied Biochemistry. Faculty of Engineering LTH. Lund University. Lund, Sweden.

## Abstract

The objective of this study was to evaluate the proximal composition, as well as Total Antioxidant Capacity (TAC) and Total Phenols (TPH) in meals that represent a complex food matrix, from different hospitals in Bolivia and Sweden. Protein, fat, ash, dietary fiber and carbohydrate contents were measured in 29 samples: 20 from two Bolivian hospitals and 9 from the university hospital in Lund, Sweden. The antioxidant capacity was measured by three spectrophotometric methods: the ferric reducing antioxidant power (FRAP) method, the 2, 2'-azinobis-3-ethylbenzothiazoline-6-sulfonic acid (ABTS) method and Total Phenolic Compounds (TPH) using the Folin-Ciocalteu reagent.

The results show that fat, protein, carbohydrate and dietary fiber in Bolivian and Swedish hospital meals are following internationally established recommendations. Regarding the main courses, TPH contents in both countries were in the same range. However, TAC and dietary fiber content were higher in Swedish meals than in Bolivian meals and the TAC was far lower, in both cases, in comparison with the value obtained from individual food items reported from literature. The results show that antioxidant levels can be easily overestimated by considering only individual uncooked ingredients. An interesting consideration is, the fiber content in the meals, which can be an important source of antioxidants and non-extractable phenolic compounds.

(*Nutr Hosp.* 2012;27:1009-1016)

DOI:10.3305/nh.2012.27.4.5849

Key words: *Total antioxidant capacity. Phenolic compounds. Bolivian meal. Swedish meal.*

## COMPOSICIÓN QUÍMICA, ACTIVIDAD ANTIOXIDANTE Y COMPUESTOS FENÓLICOS EN ALMUERZOS COLECTADOS EN HOSPITALES DE BOLIVIA Y SUECIA

## Resumen

El objetivo de este estudio fue evaluar la composición proximal, así como la capacidad antioxidante total (CAT) y los fenoles totales (FT) en alimentos que representan una matriz compleja en diferentes hospitales de Bolivia y Suecia. Se midieron las proteínas, las grasas, la ceniza, la fibra dietética y el contenido en hidratos de carbono en 29 muestras: 20 de dos hospitales bolivianos y 9 del hospital universitario de Lund, Suecia. La capacidad antioxidante se midió mediante tres métodos espectrofotométricos: el método del poder antioxidante reductor férrico (PARF), el método del 2, 2'-azinobis-3-etilbenzotiazolina-6-ácido sulfónico (ABTS) y el de Compuestos fenólicos totales (FT) empleando el reactivo Folin-Ciocalteu.

Los resultados muestran que las comidas de los hospitales bolivianos y sueco siguen las recomendaciones internacionales con respecto al contenido de grasa, proteínas, hidratos de carbono y fibra dietética. En cuanto a las comidas principales, el contenido de FT estaba en el mismo rango en ambos países. Sin embargo, la CAT y el contenido de fibra dietética fue superior en las comidas suecas que en las bolivianas y la CAT estaba muy por debajo, en ambos casos, en comparación con el valor obtenido para los alimentos individuales reportado en la bibliografía. Estos resultados muestran que los niveles de antioxidantes pueden sobrestimarse fácilmente si sólo se considera los ingredientes no cocinados. Una consideración interesante es el contenido de fibra en las comidas, que puede ser una fuente importante de antioxidantes y compuestos fenólicos no extraíbles.

(*Nutr Hosp.* 2012;27:1009-1016)

DOI:10.3305/nh.2012.27.4.5849

Palabras clave: *Capacidad antioxidante total. Compuestos fenólicos. Comida boliviana. Comida sueca.*

**Correspondence:** Leslie Tejada.  
School of Chemistry.  
San Andrés University.  
P.O. Box 303 La Paz, Bolivia.  
E-mail: Leslie.Tejeda@food.lth.se

Recibido: 12-III-2012.

1.ª Revisión: 13-III-2012.

Aceptado: 27-III-2012.

## Abbreviations

ABTS: 2, 2'-azinobis-3-ethylbenzotiazoline-6-sulfonic acid.

FAO: Food and Agricultural Organization.

FRAP: Ferric Reduction Antioxidant Power.

HO: Hospital Obrero.

HPLC: High Performance Liquid Chromatography.

LUH: Lund University Hospital.

p.a.: pro analysis.

SD: Standard deviation.

SSU: Hospital of the Seguro Social Universitario.

TAC: Total Antioxidant Capacity.

TPH: Total phenolic compounds.

TPTZ: 2,4,6-tripyridyl-s-triazine.

Trolox: 6-hydroxy-2,5,7,8-tetramethyl-chroman-2-carboxylic acid, 97%.

WHO: World Health Organization.

## Introduction

In recent years the study of food quality through the measurement of active compounds has evolved to be an important area. There is great scientific interest in the nutritional importance of antioxidants and other compounds e.g. vitamins C and E, polyphenols and carotenoids. Both epidemiological and clinical studies have indicated that increased consumption of phenolic antioxidants present in cereals, fruits, and vegetables may be linked to reduced incidences of chronic and degenerative diseases.<sup>1-6</sup>

In contrast, literature data on the content and composition of polyphenols in food is limited and often insufficient to determine the total dietary intakes. Certain studies have provided data concerning the intake of some types of polyphenols such as flavonols, flavanones, catechins, phenolic acids and flavan-3-ols, but there is a lack of comprehensive data on total polyphenol intake.<sup>7,8</sup> Mostly the data found in the literature regarding antioxidant content in food are referred to uncooked food and plant material<sup>9-12</sup> while information regarding cooked meals is scarce. It is, however, important to investigate the antioxidant content in whole diets.

A drawback of studies on individual foods is that they may overestimate their relative contributions to the antioxidant capacity within a whole diet. Individual foods known to have a high antioxidant capacity may contribute very little to the antioxidant capacity of whole diets.<sup>7</sup> Furthermore, when foods are consumed together in a diet, the total antioxidant capacity may be influenced via synergistic, additive, or antagonistic interactions among the components, which may in turn alter their physiological impacts.<sup>13-16</sup>

Hospital meals would be a good indicator of macro-nutrient content and the importance of antioxidants in a healthy diet. They are also prepared under somewhat standardized conditions. Oxidative stress that occurs following injury results in significant depletion of

many endogenous antioxidants (vitamin C, E, selenium). Increasing evidence suggests antioxidant supplementation reduces infectious complications and organ dysfunction following injury and hemorrhagic shock.<sup>2,3,5,6,17</sup>

Furthermore, higher antioxidant status may improve survival after stroke in patients at nutritional risk. Energy- and protein supplementation might further reduce the mortality risk which can be considered important for diets in hospitals.<sup>18</sup>

In the present study, which is part of a research programme on the antioxidant capacity of Bolivian foods<sup>12,19,20</sup> we investigated the chemical composition, TAC and TPH content in meals collected in hospitals in Bolivia and Sweden. Our objective was to compare these values in order to have a presentation of what the patients are eating as food in the hospitals in terms of macronutrients and antioxidant levels. Furthermore, the aim was to illustrate the importance of determining antioxidant content in whole prepared meals as compared to individual food stuffs.

## Materials and methods

### Chemicals

Folin-Ciocalteu reagent, gallic acid, sodium carbonate, acetone (p.a.) were purchased from Merck (Darmstadt, Germany), ABTS [2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid)], potassium persulphate, Trolox (6-hydroxy-2,5,7,8-tetramethyl-chroman-2-carboxylic acid, 97%), TPTZ (2,4,6-tripyridyl-s-triazine) were obtained from Sigma-Aldrich (St. Louis, MO, USA), ferric chloride from ICN Biomedicals (Costa Mesa, CA, USA), glacial acetic acid (p.a.) and sodium acetate from BDH Chemicals (Poole, UK) and methanol HPLC grade from Laboratory Supplies (Poole, UK).

Sulfuric acid (95-97%), reagent grade ISO, was obtained from Scharlan (Barcelona, Spain). Sodium hydroxide (p.a.) was purchased from Biopack (Buenos Aires, Argentina).

### Collection of meals

The description of the ingredients for the whole meals collected at the different hospitals is reported in table I. Lunch meals were collected from three hospitals. The hospitals were Lund University Hospital (LUH, Lund, Sweden), Hospital of the Seguro Social Universitario in La Paz (SSU) and Hospital Obrero in La Paz (HO). The hospital meals varied regarding the type of diet served to the patients, which include standard diets (LUH, HO) as well as soft diet, diabetic diet, low sodium diet and staff diet (SSU).

For the collection of fresh lunch meals from SSU a packing cooler for each meal (soup and main course)

**Table I**  
*Description of meals served by Bolivian and Swedish hospitals*

<i>Code</i>	<i>Description of meals</i>	<i>Initial weight (g)</i>
<i>Main course</i>		
<i>Seguro Social Universitario (SSU)</i>		
SSU_1	Pejerrey (fish), rice and boiled vegetables <sup>1)</sup>	294.5
SSU_3	Fried chicken, boiled vegetables and manioc <sup>2)</sup>	285.5
SSU_5	Steak, piece of fresh cheese, boiled vegetable and potatoes <sup>3)</sup>	302.5
SSU_7	Hamburger , boiled vegetables and manioc <sup>4)</sup>	332.0
SSU_9	Charquekan ( fried dry meat , boiled egg, piece of fresh cheese, potato and corn) <sup>3)</sup>	360.0
<i>Soups</i>		
SSU_2	Semolina <sup>1)</sup>	156.0
SSU_4	Semolina <sup>2)</sup>	333.5
SSU_6	Pumpkin <sup>3)</sup>	194.0
SSU_8	Oat <sup>4)</sup>	433.0
SSU_10	Vegetable cream <sup>3)</sup>	433.0
<i>Main course</i>		
<i>Hospital Obrero (HO)</i>		
HO_1	Oven cooked chicken, boiled vegetables (potato, carrot, green bean) mayonnaise sauce and tomato <sup>3)</sup>	386.0
HO_3	Breaded beef fillet, fried potatoes, rice and salad (boiled red beet, tomato and onion) <sup>1)</sup>	440.5
HO_5	Boiled chicken with peanut sauce, pasta and potato <sup>b)</sup>	582.5
HO_7	Stewed meat with lentils and rice <sup>1)</sup>	504.0
HO_9	Paella type rice (vegetables, minced meat and some spice) and fried egg <sup>1)</sup>	363.5
<i>Soups</i>		
HO_2	Quinoa <sup>2)</sup>	362.0
HO_4	Semolina <sup>2)</sup>	355.0
HO_6	Rice <sup>2)</sup>	435.5
HO_8	Wheat <sup>2)</sup>	444.5
HO_10	Chairo (Potato, chuño, corn, wheat and vegetables) <sup>2)</sup>	528.0
<i>Main course</i>		
<i>Lund University Hospital (LUH)</i>		
LUH_1	Vegetable cream, meatballs, bread and semolina cake with fruit sauce (lingonberry)	371.93
LUH_2	Beef with apple sauce, boiled potatoes and green peas	351.94
LUH_3	Fried pork with onion sauce, boiled potatoes, cauliflower and carrot	407.83
LUH_4	Bolognese pasta, cauliflower and carrot	350.27
LUH_5	Meatball, gravy, boiled potatoes and American vegetable mix	369.23
LUH_6	Mashed potatoes ball, fried bacon, lingonberry and American vegetable mix	285.33
LUH_7	Fishballs with mashed turnips, dill sauce, boiled potatoes and peas	474.69
LUH_8	Beef stew and boiled beetroot	501.55
LUH_9	Marengo chicken, boiled potatoes and broccoli	393.09

The meals of the Seguro Social Universitario were different, as Soft diet<sup>1)</sup>; Diabetic diet<sup>2)</sup>; Normal diet<sup>3)</sup> and Low salt diet<sup>4)</sup>. The meals (Main course<sup>a)</sup> and Soup<sup>b)</sup>) of Hospital Obrero were normal diets, as the meals from Lund University Hospital.

was used. The meals were kept at -20 °C before the start of the analysis. The collection was made every lunch time during 5 days. A main course and soup were picked up, and on each day a different kind of meal was collected, namely: soft diet, diabetic diet, low sodium diet, standard diet and staff diet. The collection of fresh lunch meals from HO followed the same procedure as for the SSU but all meals were standard diets.

Fresh lunch meals were collected from LUH on five different days (two meals per day during four days and one meal the fifth day) and kept at -20 °C. These meals were standard diets for patients.

#### *Sample preparation*

The collected meals were thawed and homogenized with 200-400 mL of sodium acetate buffer 0.1 mol/L (pH 5.0) for 5 min (the amount of added buffer varied depending on the viscosity of the homogenate). Two aliquots (50 g) of each homogenate were lyophilized. The lyophilized samples were extracted with methanol: water (9:1, by volume) in a liquid: sample proportion of 10:1 by vortexing and then sonicating the sample in an ice-water bath (0 °C, 15 min). The mixture was centrifuged at 20,000 G for 30 min at 4 °C, and the aspi-

rated supernatant was stored at  $-80\text{ }^{\circ}\text{C}$ . The extraction was performed in duplicates during three different days.

#### *Chemical composition*

Protein content (N 6.25) was determined by Kjeldahl digestion technique followed by volumetric analysis of the result of ammonia. Fat content was determined by exhaustively extracting samples in a Soxhlet apparatus with petroleum ether.<sup>21</sup> The dietary fiber content was determined after preceding separation of apolar compounds after which the sample was subjected to acid and alkaline digestion and the dietary fiber content was determined gravimetrically.<sup>22</sup> Ash was determined based on the content of inorganic matter after the incineration.<sup>23</sup> Moisture content was determined by Bolivian norm.<sup>24</sup> Total carbohydrates were calculated by difference.

#### *Measurement of TAC*

##### The ABTS method

To oxidize the colorless ABTS to the blue-green ABTS<sup>+</sup> radical cation, 5 mL of ABTS solution (7 mmol/L) was mixed with 88  $\mu\text{L}$  of potassium persulfate (140 mmol/L) and stored at room temperature in the dark overnight. On the day of analysis the ABTS<sup>+</sup> radical cation solution was diluted with acetate buffer to an absorbance of 0.70 ( $\pm 0.02$ ) at 734 nm. A Trolox standard stock solution, 5 mmol/L in ethanol, was diluted with acetate buffer to concentrations of 20-200  $\mu\text{mol/L}$ .<sup>11,25</sup> Different standards or samples (100  $\mu\text{L}$ ) were added to 1 mL of ABTS<sup>+</sup> solution, mixed for 30s, after which the absorbance reading was started after another 30 s and maintained during 6 min at 734 nm and  $25\text{ }^{\circ}\text{C}$ . The concentration was plotted against percent inhibition which was used for the calculation. The results were expressed as Trolox equivalents.

##### The FRAP method

The FRAP method was performed as described by Benzie et al. (1996).<sup>26</sup> A solution of TPTZ (10 mmol/L) was made in HCl (40 mmol/L). The FRAP reagent solution was prepared on the day of analysis by mixing 25 mL of 0.1 M acetate buffer (pH 3.6), 2.5 mL of TPTZ, and 2.5 mL of ferric chloride (20 mmol/L). A Trolox standard stock solution (5 mmol/L), prepared in ethanol, was diluted with acetate buffer to a concentration range of 100-1,000  $\mu\text{mol/L}$ . Each standard and sample (30  $\mu\text{L}$ ) were mixed with 900  $\mu\text{L}$  of FRAP solution and 90  $\mu\text{L}$  of water. A blank sample was prepared by mixing 900  $\mu\text{L}$  of FRAP solution with 120  $\mu\text{L}$  of water. The mixtures were measured after 10 min in a

spectrophotometer at 593 nm. The results were expressed as Trolox equivalents.

#### Measurement of total phenolic compounds

Total Phenolic Compounds (TPH) was determined by the Folin-Ciocalteu reagent which oxidizes the phenolic compounds to phenolates at alkaline pH in a saturated solution of sodium carbonate, resulting in a blue molybdenum-tungsten complex.<sup>27</sup> The Folin-Ciocalteu reagent was diluted with water (1:10 by volume) prior to analysis. A gallic acid stock solution was prepared in 80% aqueous acetone (1:1 by volume), and the gallic acid standard solution was diluted with water to concentrations of 235-1,180  $\mu\text{mol/L}$ . From each standard solution and sample 50  $\mu\text{L}$  was mixed with 2.5 mL of Folin-Ciocalteu reagent and 2.0 mL of sodium carbonate solution. The samples were mixed and incubated at  $45\text{ }^{\circ}\text{C}$  for 30 min. The absorbance was read at 765 nm after cooling the sample to room temperature. The results were expressed as gallic acid equivalents (GAE).

#### *Statistical analysis*

The data is reported as mean and standard deviation SD of two extractions each measured in triplicates for ABTS, FRAP and TPH.

## Results

The results of the proximal analysis for the whole meals are reported in table II and III.

Comparing the results with the recommendation by FAO/WHO,<sup>28</sup> the meals prepared at HO, showed higher protein content representing 17% of the total sample while soups showed lower fat content (4%). The main courses in SSU were prepared for different types of special diets and the protein content was higher, approximately 30%, (the highest in comparison with the other hospitals) as well the carbohydrate content in soups (84%).

On the other hand the main dishes prepared by LUH were standard diets showed somewhat higher protein content (19%).

Comparing the standard diets from HO and LUH the macronutrient content was similar.

Regarding the dietary fiber content in the foods, the Swedish national food administration recommends<sup>29</sup> a consumption of approximately 30 g of dietary fiber per day, which could be around 6 g per meal.

Comparing the main dishes from the three hospitals, the meals prepared in LUH were richer in dietary fiber (2.3 g), followed by SSU (1.45 g). The dietary fiber content in soups, from SSU (0.9 g) had higher dietary fiber content than from HO (0.6 g). The dietary fiber

**Table II**  
*The average energy and macronutrient contents of meals served by Bolivian and Swedish hospitals*

	Moisture (g)	Ash (g)	Fat (g)	Dietary fiber (g)	Protein (g)	Carbohydrate (g)	Energy (kJ)
<i>HO</i>							
Main course	81.0 (0.7)	0.9 (0.2)	3.9 (1.7)	0.003 (0.006)	3.2 (1.1)	12.1 (2.4)	390 (42)
Soups	88.0 (2.4)	0.9 (0.1)	0.4 (0.2)	0.002 (0.001)	1.5 (0.4)	9.1 (2.3)	190 (40)
<i>SSU</i>							
Main course	90.0 (2.5)	0.9 (0.2)	0.7 (0.6)	0.004 (0.006)	2.5 (0.5)	5.0 (2.9)	190 (40)
Soups	94.0 (1.1)	0.8 (0.2)	0.3 (0.1)	0.002 (0.002)	0.5 (0.3)	4.3 (1.4)	96 (31)
<i>LUH</i>							
Main course	87.6 (1.8)	0.8 (0.1)	2.0 (0.9)	0.006 (0.07)	2.0 (0.8)	6.4 (1.6)	230 (40)

Values are expressed by 100 g of fresh weight.

**Table III**  
*Comparison of the macronutrient content in a portion regarding WHO and the Swedish National Food Administration recommendations*

	Fat (%)	Protein (%)	Carbohydrate (%)	Fiber (g)
<i>HO</i>				
Main course	20	17	63	0.98
Soups	4	14	83	0.59
<i>SSU</i>				
Main Course	9	30	61	1.45
Soups	6	10	84	0.89
<i>LUH</i>				
Main course	19	19	62	2.30
Recommended by WHO	15-30	10-15	55-75	
Recommended by the Swedish National Food Administration	30	15	55	6

The consumption of dietary fiber is 30 g per day according to the Swedish National Food Administration, but on the chart is considered one of the five meals per day.

content in main courses and soups did not fulfill the quantity of dietary fiber recommended in any investigated case.

Analysis of meals from the three hospitals showed a similar range of total antioxidant capacity and total phenolic compounds (table IV). However, Bolivian lunches apart from main courses, include soups and for this reason the TAC value content is expressed in separated items. The results obtained by the ABTS and FRAP methods are expressed as median (range) table IV.

In the Bolivian meals the TAC values obtained by the ABTS and FRAP were 0.16 (0.11-0.39) and 0.22 (0.10-0.38) Trolox eq./g fw respectively for SSU samples, and 0.23 (0.15-0.33) and 0.22 (0.12-0.47) for HO.

The LUH menu showed TAC values somewhat higher than in Bolivian hospitals. For instance, 0.36 (0.12-0.43) and 0.41 (0.14-0.62) Trolox eq./g fw were obtained respectively by the ABTS and FRAP method.

TPH values measured in Bolivian soups showed lower amount of total phenols than main courses, 0.4

(0.22-0.68) and 0.39 (0.24-0.49) Trolox eq./g fw respectively. The TPH found in main Bolivian and Swedish courses are in the same range 1.11 (0.38-1.34), 0.98 (0.62-1.03) and 0.92 (0.38-1.28) Trolox eq./g fw respectively in SSU, HO and LUH samples.

## Discussion

In general, the proximal analysis showed similar patterns for the hospitals. However HO showed higher amount of fat and carbohydrates in comparison with the other parameters which is also reflected when energy content is calculated.

In general, fat, carbohydrate and protein contents of the three hospitals were within the nutrition recommendation by FAO/WHO.<sup>28</sup>

The higher protein content was found in one of the Bolivian hospital (SSU) which was almost the double in comparison with the others. The higher dietary fiber content was found in Swedish meals. However, both

**Table IV**  
*The total antioxidant capacity (TAC) and total phenols content (TPH) in the investigated meals.*  
*The number in brackets is the standard deviation*

<i>Sample code</i>	<i>ABTS (TAC)</i> <i>[<math>\mu</math>mol Trolox/g fw]</i>	<i>FRAP (TAC)</i> <i>[<math>\mu</math>mol Trolox/g fw]</i>	<i>TPH</i> <i>[<math>\mu</math>mol GAE/g fw]</i>
<i>Seguro Social Universitario (SSU)</i>			
<i>Main courses:</i>			
SSU_1	0.11 (0.01)	0.10 (0.01)	0.38 (0.05)
SSU_3	0.16 (0.03)	0.18 (0.02)	0.94 (0.14)
SSU_5	0.15 (0.01)	0.22 (0.02)	1.20 (0.18)
SSU_7	0.39 (0.05)	0.38 (0.07)	1.34 (0.21)
SSU_9	0.27 (0.02)	0.26 (0.03)	1.11 (0.07)
Median	0.16	0.22	1.11
Range	(0.11-0.39)	(0.10-0.38)	(0.38-1.34)
<i>Soups:</i>			
SSU_2	0.06 (0.01)	0.08 (0.01)	0.44 (0.05)
SSU_4	0.03 (0.01)	0.05 (0.005)	0.26 (0.03)
SSU_6	0.13 (0.02)	0.15 (0.02)	0.68 (0.08)
SSU_8	0.09 (0.01)	0.06 (0.01)	0.22 (0.02)
SSU_10	0.10 (0.02)	0.10 (0.01)	0.40 (0.06)
Median	0.09	0.08	0.4
Range	(0.03-0.13)	(0.05-0.15)	(0.22-0.68)
<i>Obrero Hospital (HO)</i>			
<i>Main courses:</i>			
HO_1	0.19 (0.01)	0.22 (0.03)	0.97 (0.17)
HO_3	0.23 (0.05)	0.47 (0.05)	0.98 (0.15)
HO_5	0.26 (0.03)	0.23 (0.03)	1.03 (0.19)
HO_7	0.33 (0.04)	0.20 (0.02)	0.62 (0.12)
HO_9	0.15 (0.02)	0.12 (0.01)	1.00 (0.17)
Median	0.23	0.22	0.98
Range	(0.15-0.33)	(0.12-0.47)	(0.62-1.03)
<i>Soups:</i>			
HO_2	0.12 (0.01)	0.13 (0.02)	0.39 (0.06)
HO_4	0.06 (0.01)	0.07 (0.01)	0.24 (0.05)
HO_6	0.09 (0.01)	0.11 (0.01)	0.49 (0.10)
HO_8	0.07 (0.01)	0.09 (0.01)	0.33 (0.06)
HO_10	0.08 (0.01)	0.10 (0.01)	0.41 (0.07)
Median	0.08	0.1	0.39
Range	(0.06-0.12)	(0.07-0.13)	(0.24-0.49)
<i>Lund University Hospital (LUH)</i>			
<i>Main courses:</i>			
LUH_1	0.12 (0.01)	0.14 (0.02)	0.38 (0.03)
LUH_2	0.36 (0.05)	0.43 (0.07)	0.94 (0.15)
LUH_3	0.37 (0.02)	0.51 (0.06)	0.94 (0.06)
LUH_4	0.38 (0.03)	0.40 (0.06)	0.92 (0.1)
LUH_5	0.17 (0.01)	0.24 (0.03)	0.76 (0.04)
LUH_6	0.41 (0.06)	0.62 (0.07)	1.26 (0.2)
LUH_7	0.25 (0.04)	0.28 (0.05)	0.89 (0.18)
LUH_8	0.28 (0.02)	0.35 (0.04)	0.91 (0.04)
LUH_9	0.43 (0.03)	0.52 (0.03)	1.28 (0.06)
Median	0.36	0.41	0.92
Range	(0.12-0.43)	(0.14-0.62)	(0.38-1.28)

Bolivian and Swedish hospital were below the Swedish National Food Administration, European and US recommendation for dietary fiber.<sup>29-31</sup>

In a recent study, reported from South Africa, the antioxidants and the nutritional levels were measured in meals for older people in a day care centre.<sup>32</sup> The data from the present study shows similar protein content. However, the amounts of carbohydrates and fat were higher than in the present study.

The antioxidant levels were somewhat higher in the Swedish hospital meals. The reason could be due to the higher amount of dietary fiber, to which antioxidants are known to sometimes be associated.<sup>33</sup> Furthermore, how the food is combined and processed could influence the antioxidant capacity.<sup>14-16,34</sup>

In comparison with the data obtained from the South African study our values are in general lower. This may be explained as above considering food processing conditions etc.

The antioxidants and phenolic compounds measured in the whole meals were lower than those values obtained in individual uncooked foods.<sup>10-12</sup> This fact is also reflected in the South African study. The reason could be due to the fact that the combination of items could give a whole matrix that inhibits the antioxidant capacity, and we also can expect some loss in antioxidant capacity due to cooking. We can conclude that antioxidant levels measured in individual food items can be overestimated in comparison to whole meals.<sup>33,35,36</sup>

The present study is among the first that reflect the antioxidant levels and phenolic content in hospital meals. The results give an interesting indication of how antioxidant levels could be evaluated in hospital and other meals. Furthermore, the results could serve as a starting-point for the quality of meals with regards to antioxidants.

In summary, the antioxidant level in whole diets has a great importance, often overestimated by considering only individual uncooked ingredients. An interesting consideration is the dietary fiber content in the meals which can be an important source of antioxidants and non-extractable phenolic compounds.

## Acknowledgements

The studies were supported by the Swedish International Development Agency (SIDA/SAREC) in a collaborative project between Universidad Mayor de San Andrés (Bolivia) and Lund University (Sweden). Helpful collaboration is acknowledged from Hospital Obrero and Seguro Social Universitario, La Paz, Bolivia as well as Lund University Hospital, Lund, Sweden.

## References

1. Shahidi F. Natural Antioxidants. *Review* 1997; 9.
2. Lampe WJ. Health effects of vegetables and fruits: assessing mechanisms of action in human experimental studies. *The American Journal of Clinical Nutrition* 1999; 70: 47S-90S.

3. Cox BD, Whichelow M, Prevost AT. Seasonal consumption of salad vegetables and fresh fruit in relation to the development of cardiovascular disease and cancer. *Public Health Nutrition* 2000; 3: 19-29.
4. Stanner SA HJ, Kelly CNM, Buttriss J. A review of the epidemiological evidence for the antioxidant hypothesis. *Public Health Nutrition* 2003; 7: 407-22.
5. Hung H-Ch JK, Jiang R, Hu FB, Hunter D, Smith-Warner SA, Colditz GA, Rosner B, Spiegelman D, Willett WC. Fruit and Vegetable Intake and Risk of Major Chronic Disease. *Journal of the National Cancer Institute* 2004; 96: 1577-84.
6. Quiñones M, Miguel M, Aleixandre A. Los polifenoles, compuestos de origen natural con efectos saludables sobre el sistema cardiovascular. *Nutrición Hospitalaria* 2012; 27: 76-89.
7. Saura-Calixto F, Goñi I. Antioxidant capacity of the Spanish Mediterranean diet. *Food Chemistry* 2006; 94: 442-7.
8. Saura-Calixto F, Goñi I. Intake and bioaccessibility of total polyphenols in a whole diet. *Food Chemistry* 2007; 101: 495-501.
9. Halvorsen BL, Holte K, Myhrstad MCW, et al. A systematic Screening of Total Antioxidants in dietary Plants. *J Nutr* 2002; 132: 461-71.
10. Halvorsen BL, Carlsen ML, Phillips KM et al. Content of redox-active compounds (ie. antioxidants) in foods consumed in the United States. *American Journal of Clinical Nutrition* 2006; 84: 95-135.
11. Nilsson J, Pillai D, Önning G, Persson C, Nilsson Å, Åkesson B. Comparison of the 2,2,2-azino-bis(3-thylbenzotiazoline-6-sulfonic acid (ABTS) and ferric reducing antioxidant power (FRAP) methods to assess the total antioxidant capacity in extracts of fruit and vegetables. *Mol Nutr Food* 2005; 49: 239-46.
12. Peñarrieta JM, Alvarado JA, Åkesson B, Bergenståhl B. Total antioxidant in Andean food species from Bolivia. *Revista Boliviana de Química* 2005; 22: 89-93.
13. Parker TL, inventor Synergistic interactions of Phenolic Compounds Found in Food. United States. 2011.
14. Wang S, Meckling KA, Marcone MF, Kakuda Y, Tsao R. Synergistic, additive, and antagonistic effects of food mixtures on total antioxidant capacities. *J Agric Food Chem* 2011; 59: 960-8.
15. Freeman BL, Eggett DL, Parker TL. Synergistic and antagonistic interactions of phenolic compounds found in navel oranges. *Food Sci* 2010; 75: C570-6.
16. Iribhogbe OI, Emordi JE, Idonije BO, Aigbiremolen A, Nwoke EO, Akpamu U. Synergistic Effects of Antioxidant Vitamins on Lipid Profile in Pregnancy. *Journal of Biological Sciences* 2011; 3: 104-9.
17. Collier BR, Giladi A, Dossett LA, Dyer L, Fleming SB, Cotton BA. Impact of High-Dose Antioxidants on Outcomes in Acutely Injured Patients. *Journal of Parenteral and Enteral Nutrition* 2008; 32: 384-8.
18. Ha L, Sakhi AK, Bohn SK et al. Antioxidants status after an acute stroke and the association with survival in elderly at nutritional risk. *The European e-Journal of Clinical Nutrition and Metabolism* 2011; 6: 135-41.
19. Peñarrieta JM, Alvarado J, Åkesson B, Bergenståhl B. Total antioxidant capacity and content of flavonoids and other phenolic compounds in canihua (Chenopodium pallidicaule): An Andean pseudocereal. *Molecular Nutrition and Food Research* 2008; 52: 708-17.
20. Peñarrieta JM, Alvarado JA, Bergenståhl B, Åkesson B. Total antioxidant capacity and content of phenolic compounds in wild strawberries (*Fragaria vesca*) collected in Bolivia. *International Journal of Fruit Science* 2009; 9: 344-59.
21. Gutiérrez V, Bonifacio A, Requena G et al. Cereales-Quinoa en grano-Determinación del contenido de materia grasa. *Norma Boliviana NB 665* 1996: 13-5.
22. Cortés J, Donoso O, Gonzáles V. Cereales. Método de ensayo para determinar la fibra cruda. *Norma Boliviana NB 104-75* 1975: 18-20.
23. Lopez JJ, Achá J, Scott R et al. Cereales - Métodos para determinación de cenizas. *Norma Boliviana NB 075* 1997: 1-3.

24. Lopez JJ, Achá J, Scott R et al. Cereales - Métodos para determinar el contenido de humedad. *Norma Boliviana NB074* 1997: 1-3.
25. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidants activity applying an improved ABTS radical cation decolorization assay. *Free Radicals in Biology and Medicine* 1999; 26: 1231-7.
26. Benzie IFF, Strain JJ. The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power" the FRAP assay. *Analytical Biochemistry* 1996; 239: 70-6.
27. Singleton VL, Rossi JA. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagent. *Am J Enol Vitic* 1965; 16: 144-58.
28. FAO/WHO. Launch expert report on diet nutrition and prevention of chronic diseases. Report of Food and Agriculture Organization (FAO) and the World Health Organization (WHO) expert consultation. In: Roma 2003.
29. Enghardt Barbieri H, Lindvall C. Swedish Nutrition recommendations Objectified (SNO). National Food Administration. In: National Food Administration 2003: 16.
30. EFSA Panel on dietetic products nutrition and allergies. Scientific opinion on dietary reference values for carbohydrates and dietary fibre. *EFSA Journal* 2010; 8 (3): 1-77.
31. Dietary guidelines for Americans. Washington. US Department of Agriculture 2005 DU. Dietary guidelines for Americans. In: Washington: US Department of Agriculture, DC:USDA; 2005.
32. Medoua GN, Egal AA, Oldewage-Theron WH. Nutritional value and antioxidant capacity of lunch meals consumed by elderly people of Sharpeville, South Africa. *Food Chemistry* 2009; 115: 260-4.
33. Saura-Calixto F. Dietary Fiber as a Carrier of Dietary Antioxidants: An Essential Physiological Function. *J Agric Food Chem* 2011; 59: 43-9.
34. Ragaei S, Guzar I, Dhull N, Seetharaman K. Effects of fiber addition on antioxidant capacity and nutritional quality of wheat bread. *Food Science and Technology* 2011; 44: 2147-53.
35. Gökmen V, Serpen A, Fogliano V. Direct measurement of the total antioxidant capacity of foods: the "Quencer" approach. *Trends in Food Science & Technology* 2009; 20: 278-88.
36. Serrano J, Goni I, Saura-Calixto F. Food antioxidant capacity determined by chemical methods may underestimate the physiological antioxidant capacity. *Food Research International* 2007; 40: 15-21.