



Nutrición Hospitalaria



Trabajo Original

Paciente crítico

Nitrogenous content in parenteral nutrition: a four-year experience in a general hospital. Critically-ill patient specificity

Contenido nitrogenado de la nutrición parenteral: un estudio de cuatro años en un hospital general. Especificidad en paciente crítico

Juan Luis Villalobos-Gómez^{1,2,3}, Carlos Lara-Ramos¹, Yolanda Domínguez-Rivas¹, Antonio Vallejo-Báez^{3,4}, Francisco Cota-Delgado⁴, Efrén Márquez-Fernández¹, José Manuel García-Almeida^{2,3,5}, José Antonio López-Medina^{3,5}, Rosalía Rioja-Vázquez^{2,3,5}, Albert Santacreu-Regí⁶, Francisca Rius-Díaz⁷ and Alfredo Mínguez-Mañanes^{3,8}

¹Hospital Pharmacy Department. Nutrition Section. Hospital Universitario Virgen de la Victoria. Málaga, Spain. ²Nutritional Support Team. Hospital Universitario Virgen de la Victoria. Málaga, Spain. ³Nutrition Committee. Hospital Universitario Virgen de la Victoria. Málaga, Spain. ⁴Intensive Care Department. Hospital Universitario Virgen de la Victoria. Málaga, Spain. ⁵Endocrinology and Nutrition Department. Hospital Universitario Virgen de la Victoria. Málaga, Spain. ⁶Informatics Development Department. B. Braun Medical. Barcelona, Spain. ⁷Biostatistics Department. Universidad de Málaga. Málaga, Spain. ⁸Anesthesiology and Recovery Department. Hospital Universitario Virgen de la Victoria. Málaga, Spain

Abstract

Introduction: There have been several studies focusing on caloric intake during the last years, while protein content relevance has been underestimated. Some recent evidence has shown that protein deficiency has also an impact on patient outcomes. We have studied the nitrogen (N) content in parenteral nutrition (PN) bags administered to adult patients in a Spanish tertiary level hospital for four years.

Material and methods: Patients who received parenteral nutrition in the general ward and Intensive Care Unit (ICU) were recorded. Caloric and protein content were registered and adjusted to weight and length of stay. Data were compared among three groups of patients: those in the general ward, those in the ICU and those requiring renal replacement therapy (RRT). The one-factor analysis of variance (ANOVA) test was used after checking data normality and homoscedasticity

Results: There was an increase in the mean g N/stay year after year ($p < 0.01$) from 14 to 15.5 g, with a decrease in non-protein caloric content ($p < 0.001$) from 111.6 to 101.8 kcal/g N. The range was established from 4.1 to 32.6 g. PN diets with ≥ 18 g N% ranged from 12.8% (2010) to 19.6% (2013). There were significant differences among the groups when comparing the variable g N/stay ($p < 0.0001$): 13.5 general ward vs 15.9 ICU patients vs 17.6 ICU with RRT, also when referring to adjusted weight.

Conclusions: According to most recent recommendations nitrogen has been provided in higher amounts than previously, especially in critical care patients with RRT.

Key words:

Nitrogen. Protein. Parenteral nutrition. Critical care. Renal replacement therapies. Individualized diets.

Resumen

Introducción: algunos estudios recientes sugieren que se ha dado gran importancia al aporte calórico en la nutrición parenteral (NP) del paciente adulto, infraestimando su contenido proteico. Sin embargo, se ha demostrado su relación con los resultados clínicos. Con este objetivo se ha estudiado el contenido en nitrógeno (N) de las NP administradas en un hospital terciario a lo largo de cuatro años.

Material y métodos: se recogieron datos de la NP de pacientes hospitalizados en planta, así como en la Unidad de Cuidados Intensivos (UCI). El peso del paciente, su índice de masa corporal (IMC), el contenido en nitrógeno (total y por peso), el aporte calórico no proteico y la duración de la NP fueron algunas de las variables estudiadas. Se compararon en 2013 los aportes en la planta general, en UCI y en aquellos que recibieron algún tipo de terapia renal sustitutiva (TRS). Se utilizó el análisis de varianza (ANOVA) de un factor, previa comprobación de la normalidad y homocedasticidad.

Resultados: se ha observado un aumento progresivo en aporte nitrogenado medio diario cada año ($p < 0,01$) de 14 a 15,05 g, con descenso del contenido calórico no proteico ($p < 0,001$) de 111,6 a 101,8 kcal /g N. El rango de N en bolsa fue de 4,1 a 32,6 g. Aumentó el porcentaje de bolsas con ≥ 18 g N (12,8 en 2010 vs. 19,6 en 2013). También hubo diferencias entre grupos de pacientes en g N/estancia ($p < 0,0001$): 13,5 plantas de hospitalización vs. 15,9 UCI vs. 17,6 UCI con TRS, igualmente si referidos a peso ajustado.

Conclusiones: En consonancia con las recomendaciones más recientes, el contenido en nitrógenos ha aumentado con los años, en especial en la NP del paciente crítico, siendo aún mayor en los sometidos a TRS.

Palabras clave:

Nitrógeno. Proteína. Nutrición parenteral. Cuidados intensivos. Técnicas de depuración extrarrenal. Dietas individualizadas.

Received: 28/10/2016
Accepted: 23/02/2017

Villalobos-Gómez JL, Lara-Ramos C, Domínguez-Rivas Y, Vallejo-Báez A, Cota-Delgado F, Márquez-Fernández E, García-Almeida JM, López-Medina JA, Rioja-Vázquez R, Santacreu-Regí A, Rius-Díaz F, Mínguez-Mañanes A. Nitrogenous content in parenteral nutrition: a four-year experience in a general hospital. Critically-ill patient specificity. Nutr Hosp 2017;34:548-554

DOI: <http://dx.doi.org/10.20960/nh.699>

Correspondence:

Juan Luis Villalobos Gómez. Hospital Pharmacy Department. Nutrition Section. Hospital Universitario Virgen de la Victoria. Campus de Teatinos, s/n. 29010 Málaga, Spain
e-mail: juanluisvillalobos@gmail.com

INTRODUCTION

There have been several recent studies focusing on caloric intake in parenteral nutrition (PN) while protein content relevance has been underestimated. Protein deficiency has also an impact on patient outcomes (1,2).

There is a clear need to identify safe amounts of nutrients for each acute disease stage in order to avoid under or overfeeding (3). In critically ill patients, undernourishment has been correlated with a higher number of infections, respiratory and immunology function impairment, as well as an increase in hospital length of stay and mortality (4). The results of a European study in 2012 including 102 patients showed a higher mortality rate on those patients who received a lower protein load, less than 1.2-1.5 g/kg/day (5). Other recent trials have shown that most critically ill patients do not reach nutritional requirements, being a low protein intake the most remarkable one (5-9), although there are some exceptions (10).

Based on our experience, we wanted to design a study to quantify the nitrogenous content in parenteral nutrition (PN) during 2010-2013 periods, comparing patients in the general ward and in the Intensive Care Unit. At the same time, we recorded the number of standardized/tailored solutions as well as insulin and glutamine addition.

MATERIAL AND METHODS

A retrospective descriptive study was performed. The pharmacist recorded daily, clinical, anthropometric and analytical information using the Nutridata® program. PN composition was also recorded. This program offers the possibility to explore data through SPSS®. Some content parameters, considered as targets, were obtained to estimate differences among different groups of patients.

Some of the analyzed variables were current weight, present weight, ideal weight (IW) and the % of weight loss/time. Body mass index (BMI) was calculated after exporting weight and height information to Excel, as well as the adjusted weight ($AW = IW \pm 25\%$ of the excess or defect in weight). Current weight data are the result of the analysis done through the Process InforNut® (11), when a nutritional risk alarm arises in the filter FILNUT (12,13), during the hospital stay. If there was no opportunity to record the patient's weight, it was obtained through digital clinical records (Diraya specialized care - DAE - oncology pharmacy - Farmis Oncofarm®, reports from preanesthesia, hemodynamics, etc.). In any case, it is mandatory to fulfill patient's weight and height into the justification sheets for PN (14). Any missing information would be obtained through the visit done by a member of the nutritional team. When a new weight is recorded into Nutridata® from a patient on long-term PN, the mean weight is used as the calculator for global weight/case. When it is not feasible to get patient's height, it can be estimated from the ulna bone length. Variables such as g N/kg or the non-protein kcal/g N are calculated by the program on a daily basis. The percentage and number of PN diets

and their distribution as per the different target variables were calculated using the Nutridata® database per case and day. The information regarding g N/kg is referred to the current weight in the four-year period. Enteral nutrition intake was not considered.

Three groups of patients were established: general ward, ICU, and ICU requiring renal replacement therapy (RRT). Patients on RRT are those who received PN at any time during the RRT therapy, even if it was for a short period of time.

In 2011, the Nutritional Support Team proposed a new PN protocol. It included 27 different diets ranging from standard, fat-free, cholestasis, renal or liver failure and sepsis mixtures with different nutrient content. As a novelty, it also included diets with 18 and 20 g N with a low non-protein kcal/g N ratio (highly stressed patients) (Smofkaviben® and Olime® triple chamber bags); glutamine dipeptide (Dipeptiven®) was added in certain cases, mainly in critically ill patients. A personalized diet was defined as the one with different macronutrients and/or volume composition when compared with those in the protocol, but not different in terms of electrolytes, insulin addition or glutamine composition.

The one-way analysis of variance (ANOVA) test was used after checking data for normality and homoscedasticity.

RESULTS

Table I shows results in the three groups. The most severe and the longer the stay, the longer the length of PN use, and the higher the nitrogen load. The number of tailored formulations was higher in ICU patients requiring RRT. Table II shows yearly evolution in the study period. The number of PN bags remains similar, with a trend to higher nitrogen content and lower nitrogen/non-protein calorie ratio.

Figure 1 shows the evolution of the nitrogen intake as per weight, in the four-year period and other indicators fixed as targets. Figure 2 shows average values/stay (day) and 95% CI for nitrogen (g), non-protein kcal/g N, adjusted weight (AW) and g N/Kg AW in the three groups of patients. Patients in the ICU requiring RRT received a significantly higher nitrogen load as well as a lower N/non-protein kcal ratio.

DISCUSSION

Standard PN solutions are widely used in most hospitals. There is also the possibility of designing standardized formulations based on standard patient profiles, including different stress situations, sepsis, cholestasis, and renal or hepatic impairment. Individualized formulations could be of interest in those patients with a complex clinical situation (up to 25-40% of the cases) (15). The ready-to-use bags (triple chamber bags) have dramatically reduced the need for manual compounding, by significantly reducing manipulation and liquid transfer (16). Martínez Romero et al. (17) demonstrated that 75% of their metabolically stable patients had their needs addressed by only three standardized formulations they had designed according to case-mix. Schoenenberger

Table I. Comparison of nitrogen content in patients according to their location

2013 Total patients 365 ⁽¹⁾	Hospitalization wards	ICUs	ICUs with RRT ⁽²⁾
Cases	251	168	53
Administered bags	2,973	2,178	1,015
PN days/case	11.84	12.97	19.15
BMI/case	25.19 ± 5.28	27.06 ± 6.23	28.58 ± 6.6
Current weight/case (kg)	69.46 ± 15.82	75.89 ± 17.74	81.81 ± 15.33
Current weight/stay	68.68 ± 16.19	79.60 ± 18.20	85.13 ± 17.19
Ideal weight/stay	58.83 ± 7.87	61.67 ± 7.76	61.96 ± 8.64
Adjusted weight/stay	61.37 ± 8.28	66.32 ± 8.25	67.99 ± 8.13
<i>g N/ stay (mean ± σ)</i>	<i>13.56 ± 3.13</i>	<i>15.87 ± 4.19</i>	<i>17.60 ± 4.28</i>
% ≤ 18	94.8	72.4	52.7
% 18.1-22	4.4	20.4	33.8
% > 22	0.8	7.2	13.5
<i>Non-prot kcal/g N/stay (mean ± σ)</i>	<i>112.2 ± 20.1</i>	<i>97.3 ± 22.6</i>	<i>89.68 ± 18.4</i>
% ≤ 80	2.9	14.9	27.5
% 80.1-95	14.5	44.5	44.7
% > 95	82.6	40.6	27.8
<i>g N/kg*/stay (mean ± σ)</i>			
*/adjusted weight	<i>0.223 ± 0.046</i>	<i>0.240 ± 0.059</i>	<i>0.260 ± 0.060</i>
Protein g equivalent	1.39	1.5	1.63
*/current weight	<i>0.204 ± 0.052</i>	<i>0.205 ± 0.056</i>	<i>0.211 ± 0.056</i>
*/ideal weight	<i>0.233 ± 0.055</i>	<i>0.260 ± 0.068</i>	<i>0.288 ± 0.070</i>
% ≥ 0.26-current-	15.6	20.4	22.8
% personalized diets	17.1	25	41.3
% protocol (% triple chamber bags)	82.9 (38)	75 (36.7)	58.7 (25.7)
PN bags with insulin (%)	8.6	22.5	31.3
Glutamine g/bag ⁽³⁾	0.8	4.83	7.43

¹Fifty-four patients stayed both in ICUs and wards; 731 bags were excluded (five homecare patients). ²Critically ill patients requiring RRT at any time. It includes all bags with or without RRT. ³Equivalent to 6.1; 36.6 and 56.3 ml of Dipeptiven®/bag.

et al. showed that a standardized protocol which included 20 diets helped to prepare 6,300 formulations; up to 30% of the cases were prepared from different macronutrients, although only 8% of the formulations were really individualized (18).

As our own program allowed collecting data on prescription, elaboration and dispensation, we were able to assess the protein load in our PN solutions in all three scenarios: general ward, ICU patients, and those in the ICU requiring RRT. There was an increase in the g N/stay year after year ($p < 0.01$) except from 2012 and 2013 ($p = 0.181$). The range was established from 4.1 to 32.6. PN (%) diets with ≥ 18 g N have increased every year, ranging from 12.8% (2010) to 19.6% (2013). Data from 2014 show the same trend (21.3% ≥ 18 g N). There were no yearly significant differences for mean weight or BMI per case. Nevertheless, the mean weight/stay on PN in 2013 was significantly higher ($p < 0.001$) than in the three previous years.

In Weijs paper, including 886 patients, an optimal caloric and protein intake (between 1.2 and 1.5 g/kg/day) was associated with a decrease in mortality, while only reaching the caloric targets

was not sufficient to establish that association (19). In agreement with this, in a pilot study including 2,772 ICU patients a higher caloric and protein intake was associated with better clinical outcomes, especially if BMI was < 25 or ≥ 35 on admission (20). Other studies confirm these findings (21,22), suggesting that classically recommended protein intake (ESPEN) may not be sufficient.

Regarding the use of glutamine, most published studies in critically ill patients have shown beneficial effects (23), leading to diverse scientific societies to recommend glutamine use (24,25). Heyland described that very high doses administered through enteral and parenteral route (> 0.5 g/kg/day) and during the acute phase in patients with multiple organ dysfunction and shock increased mortality (26).

According to recent recommendations, we have increased the g N/stay in our PN from 14 to 15.05, with a decrease in the non-protein caloric content, from 112 to 102. This nitrogen increase has happened despite the increase on PN average duration (two days) and partly due to the increase in the number of complementary PN, but also as a result of an improvement on

Table II. Nitrogen content evolution between 2010 a 2013

Year	2010	2011	2012	2013
<i>PN cases⁽¹⁾</i>	411	419	409	365
Mean age/case	61	61.5	62.7	59.7
Males %	62	60.6	62.1	61.4
<i>Stays (bags)</i>	5,738	6,142	6,141	5,882
Days/case	14.2	14.9	15.2	16.6
<i>BMI</i>	25.82 ± 5.38	25.77 ± 5.97	25.91 ± 5.66	25.86 ± 5.49
Current weight/case	71.28 ± 14.9	71.24 ± 16.44	71.88 ± 16.82	71.98 ± 16.37
Current weight/stay	72.44 ± 17.51	71.82 ± 18.58	71.63 ± 20.11	74.25 ± 19.56
<i>g N/stay (mean ± σ)</i>	14.00 ± 3.28	14.62 ± 4.21	14.89 ± 4.08	15.05 ± 4.24
% ≥ 16 g N	29.65	37.87	43.09	43.9
% ≥ 18 g N	12.8	17.2	17.4	19.6 (2014: 21.3)
% ≥ 22 g N	1.73	5.37	5.06	7.6
<i>g N/kg-current-/stay (mean ± σ)</i>	0.201 ± 0.057	0.211 ± 0.066	0.218 ± 0.070	0.210 ± 0.059
Protein g equivalent	1.256	1.319	1.363	1.313
% ≥ 0.26 g N/kg	15.9	21.7	25	21.3
% ≥ 0.3 g N/kg	4.7	6.9	12.2	10.1
<i>Non-prot kcal/g N/stay (mean ± σ)</i>	111.6 ± 22.2	107.6 ± 25.1	106.1 ± 20.3	101.8 ± 25.4
% ≤ 95 non-prot kcal/g N	24.8	36.8	37.2	41.4
% ≤ 110 non-prot kcal/g N	36.8	48.5	48.6	56.4
% >110 non-prot kcal/g N	63.2	51.5	51.4	42.9
% Personalized diets ⁽²⁾	7	15.5	20.7	26.5
% Protocol (% triple chambre bags)	93 (46.5)	86.5 (44.7)	79.3 (33.7)	73.5 (31.5)
% bags with insulin	11.1	11.1	16.8	15

¹Hospitalization wards, ICUs and homecare patients included. ²Personalized: those different from the protocol in terms of macronutrients and/or volume, not those with added electrolytes and/or insulin or glutamine dipeptide.

the way PN is gradually initiated in patients at risk of refeeding syndrome. There was a yearly significant increase in g N/kg/stay from 0.20 to 0.22 until 2012 ($p < 0.001$). In 2013 there was a significant decrease when compared with 2012. The value reached was similar to 2011 value (0.21) and higher than 2010 value. This could be explained by the higher current weight/stay in 2013. Its equivalent in g prot/kg/d varied between 1.26 and 1.38, in accordance with ESPEN guidelines for PN (27).


The reduction in non-protein kcal/g of N turns out to be significant ($p < 0.001$ year by year, excepting 2011-2012, $p < 0.05$). An important reduction appeared in this ratio, lowered from 111.6 (2010) to 101.8 (2013). The need for personalized diets was more necessary (26.5% in 2013) when the nitrogenous content increased.

We also found a significant difference among groups when comparing the variable g N/stay (13.5 vs 15.9 vs 17.6, $p < 0.0001$). The same significance was reached for the difference in non-protein kcal/g N/stay (112 vs 97 vs 90). The increase in g

N/kg/stay in ICU did not reach statistical significance when compared with hospitalization wards; however, those on RRT did ($p < 0.01$). These differences were statistically significant among the three groups when the variable g N/kg considered the IW or AW/stay ($p < 0.0001$). Patients on the general wards received 0.22 by average, while the ICU ones received 0.24. Those on RRT received 0.26 g N/kg AW, raising up to 0.29 g N when referred to the IW.

Critically ill patients, and those requiring RRT, had higher weight than those in hospital wards, received PN for a longer period, and also received higher nitrogenous content, in absolute terms and with regards to weight kg, and less non-protein calories per gram. Finally, glutamine dipeptide, as a PN component, was mainly used in these patients.

If a clinical pharmacist is fully integrated in the interdisciplinary nutrition support team, as it was in our case, in close collaboration with the Intensive Care Unit as a reference consultant, it is more feasible to adjust nutrient supplies to patient needs and, therefore, to favor quality of care in patients requiring PN.

**Figure 1.**

Four-year evolution 2010-2013 (mean/stay with PN, CI 95%).




Figure 2.

Comparison during 2013: Wards/ICUs/ICUs with RRT (mean/stay with PN, CI 95%).

REFERENCES

- Burgos R. La importancia del aporte proteico en la nutrición parenteral. En simposio «Balance optimizado en nutrición parenteral»; 29^a Congreso SENPE. Nutr Hosp 2014;30(3):7-10.
- Irlas JA. Coste-eficacia del aporte proteico. En simposio «Balance optimizado en Nutrición Parenteral»; 29^o Congreso SENPE. Nutr Hosp 2014;30(3):11-4.
- Singer P, Hiesmayr M, Biolo G, Felbinger TW, Berger MM, Goeters C, et al. Pragmatic approach to nutrition in the ICU: Expert opinion regarding which calorie protein target. Clin Nutr 2014;33(2):246-51.
- Dvir D, Cohen J, Singer P. Computerized energy balance and complications in critically ill patients: An observational study. Clin Nutr 2006;25(1):37-44.
- Casaer MP, Mesotten D, Hermans G, Wouters PJ, Schetz M, Meyfroidt G, et al. Early versus late parenteral nutrition in critically ill adults. N Engl J Med 2011 11;365(6):506-17.

6. Singer P, Anbar R, Cohen J, Shapiro H, Shalita-Chesner M, Lev S, et al. The tight calorie control study (TICACOS): A prospective, randomized, controlled pilot study of nutritional support in critically ill patients. *Intensive Care Med* 2011;37(4):601-9.
7. Heidegger CP, Berger MM, Graf S, Zingg W, Darmon P, Costanza MC, et al. Optimization of energy provision with supplemental parenteral nutrition in critically ill patients: A randomized controlled clinical trial. *Lancet* 2013;381(9864):385-93.
8. Doig GS, Simpson F, Sweetman EA, Finfer SR, Cooper DJ, Heighes PT, et al. Early parenteral nutrition in critically ill patients with short-term relative contraindications to early enteral nutrition: A randomized controlled trial. *Jama* 2013;309(20):2130-8.
9. Ferrie S, Allman-Farinelli M, Daley M, Smith K. Protein requirements in the critically ill: A randomized controlled trial using parenteral nutrition. *JPEN* 2015;20(10):1-11.
10. Oliveira G, Tapia MJ, Ocon J, Cabrejas-Gómez C, Ballesteros-Pomar MD, Vidal-Casariago A, et al. Parenteral nutrition-associated hyperglycemia in non-critically ill inpatients increases the risk of in-hospital mortality (multi-center study). *Diabetes Care* 2013;36(5):1061-6.
11. Villalobos Gómez JL, González Pérez C, García-Almeida JM, Martínez Reina A, Del Rio Mata J, Márquez Fernández E, et al. InforNut® Process improves accessibility to diagnosis and nutritional support for the malnourished hospitalized patient. Impact on management indicators. Two-year assessment. *Nutr Hosp* 2014;29(6):1210-23.
12. Villalobos Gamez JL, Garcia-Almeida JM, Guzman de Damas JM, Rioja Vazquez R, Osorio Fernandez D, Rodríguez-García LM, et al. Proceso INFORNUT: validación de la fase de filtro -FILNUT- y comparación con otros metodos de detección precoz de desnutrición hospitalaria. *Nutr Hosp* 2006;21(4):491-504.
13. Villalobos Gómez JL, Guzmán de Damas JM, García-Almeida JM, Galindo MM, Rioja Vázquez R, Enguix Armada A, et al. FILNUT-escala: justificación y utilidad en el cribaje de riesgo por desnutrición dentro del proceso INFORNUT. *Farm Hosp* 2010;34(5):231-6.
14. Villalobos JL, Tutau F, Mora R, Mazure R, Mínguez A, Rodríguez F, et al. Prescripción y costes de nutrición artificial antes y después de un nuevo protocolo de utilización. *Nutr Hosp* 2000;15(2):71-8.
15. Llop Talaveron JM, Berlana Martín D, Badia Tahull MB, Fort Casamartina E, Vinent Genestar JL, Tubau Mola M, et al. Preparados estándar de nutrición parenteral en situaciones clínicas complejas. *Nutr Hosp* 2004;19(4):229-35.
16. Llop Talaveron JM, Machi Ribes JJ, Gracia García B, Badia Tahull MB, Tubau Molas M, Jodar Masanes R. Nutrición parenteral modular: un nuevo concepto. *Nutr Hosp* 2007;22(4):402-9.
17. Martínez Romero G, Pérez Ruixo JJ, Jiménez Torres NV. Nutrición parenteral e identificación de subpoblaciones con necesidades nutricionales similares. *Nutr Hosp* 2002;17(2):80-92.
18. Schoenenberger JA, Rodríguez A. Protocolización de la nutrición artificial por vía parenteral. Bases metodológicas y organizativas para el diseño y revisión del proceso. *Nutr Hosp* 2010;25(1):26-33.
19. Weijs PJ, Stapel SN, De Groot SD, Driessen RH, De Jong E, Girbes AR, et al. Optimal protein and energy nutrition decrease mortality in mechanically ventilated, critically ill patients: A prospective observational cohort study. *JPEN* 2012;36:60-8.
20. Alberda C, Gramlich L, Jones N, Jeejeebhoy K, Day AG, Dhaliwal R, et al. The relationship between nutritional intake and clinical outcomes in critically ill patients: Results of an international multicenter observational study. *Intensive Care Med* 2009;35(10):1728-37.
21. Germano Borges de Oliveira Nascimento Freitas R, Negro Nogueira RJ, Hessel G. Las necesidades proteicas de pacientes en estado crítico que reciben nutrición parenteral. *Nutr Hosp* 2015;32(n01):250-5.
22. Dickerson RN, Medling TL, Smith AC, Maish GO 3rd, Croce MA, Minard G, et al. Hypocaloric, high-protein nutrition therapy in older vs younger critically ill patients with obesity. *JPEN J Parenter Enteral Nutr* 2013;37(3):342-51.
23. Burgos Peláez R, Escudero Álvarez E, García Almeida JM, García de Lorenzo A, García Luna PP, Gil Hernández A, et al. Farmaconutrición en el paciente grave. *Nutr Hosp* 2015;32:478-86.
24. Vanek VW, Matarese LE, Robinson M, Sacks GS, Young LS, Kochevar M, et al. A.S.P.E.N. position paper: Parenteral nutrition glutamine supplementation. *Nutr Clin Pract* 2011;26(4):479-94.
25. Mesejo A, Vaquerizo Alonso C, Acosta Escribano J, Ortiz Leyba C, Montejo González JC, Spanish Society of Intensive Care M, et al. Recomendaciones para el soporte nutricional y metabólico especializado del paciente crítico. Actualización. Consenso SEMICYUC-SENPE: introducción y metodología. *Med Intensiva* 2011;35(Suppl 1):1-81.
26. Heyland D, Muscedere J, Wischmeyer PE, Cook D, Jones G, Albert M, et al. A randomized trial of glutamine and antioxidants in critically ill patients. *N Engl J Med* 2013;368(16):1489-97.
27. Singer P, Berger MM, Van den Berghe G, Biolo G, Calder P, Forbes A, et al. ESPEN Guidelines on Parenteral Nutrition: Intensive care. *Clin Nutr* 2009; 28(4):387-400