



Trabajo Original

Pediatría

Physical activity values in two- to seven-year-old children measured by accelerometer over five consecutive 24-hour days

Valores de actividad física en niños de dos a siete años, medidos mediante actimetría durante cinco días consecutivos las 24 horas diarias

Ana Gutiérrez-Hervás¹, Ernesto Cortés-Castell², Mercedes Juste-Ruiz², Antonio Palazón-Bru³, Vicente Francisco Gil-Guillén³ and María Mercedes Rizo-Baeza¹

¹Department of Nursing. University of Alicante. San Vicente del Raspeig. Alicante, Spain. Departments of ²Pharmacology, Pediatrics and Organic Chemistry and ³Clinical Medicine. Miguel Hernández University. San Juan de Alicante, Alicante. Spain

Abstract

Introduction: interpretation of accelerometer-derived physical activity in preschool children is confounded by differences in cut-off points.

Aim: the purpose of this study was to analyze physical activity in 2-to-7-year-old children to establish reference values for daily activity.

Methods: observational study in children aged 2-7 years, without chronic diseases and whose parents provided informed consent. The main variable was physical activity, measured continuously over 120 hours (three workdays and two weekend days) by accelerometer. Secondary variables were weight status (body mass index [BMI] Z-score) and gender. The relationship between the main variable and secondary variables was determined through the t-test, ANOVA and the Pearson correlation coefficient. A multivariate model was used to obtain the standard deviation (SD) of all possible combinations of values, constructing percentiles of normality ($x \pm SD$ and $x \pm 2 \cdot SD$).

Results: one hundred and thirty-six children (35% of municipality children) were included in the study (54.4% of them were girls). Their weight status distribution was: 25 underweight (18.4%), 54 normal weight (39.7%), 12 risk of overweight (8.8%), 22 overweight (16.2%) and 23 obese (16.8%). The median age was 5.7 years and the mean physical activity was 592 counts/minute. The boys undertook more physical activity ($p = 0.031$) and the underweight and normal-weight children undertook more physical activity than the overweight and obese children ($p = 0.012$). There were no significant differences according to age. The multivariate analysis showed significant differences ($p < 0.001$) according to gender and weight status. In boys, physical activity decreased as weight status increased. In contrast, the girls in the extreme BMI groups obtained higher levels of physical activity.

Conclusion: overweight and obese preschool children had lower levels of physical activity than normal weight children. Physical activity levels were higher in boys.

Key words:

Preschool child.
Physical activity.
Pediatric obesity.
Normal distribution.

Resumen

Introducción: la interpretación de la actividad física medida mediante actimetría en preescolares es confusa debido a los diferentes puntos de corte.

Objetivo: el objetivo de este estudio fue analizar la actividad física en niños de dos a siete años para establecer valores de actividad física diaria.

Método: estudio observacional en niños de dos a siete años, sin enfermedades crónicas y cuyos padres hubieran firmado el consentimiento informado. La variable principal fue la actividad física, medida durante 120 horas ininterrumpidas (tres días laborables y dos días festivos) mediante actímetros. Las variables secundarias fueron el estado de peso (puntaje z de índice de masa corporal [IMC]) y el sexo. La relación entre la variable principal y las variables secundarias fue determinada mediante el test-t, ANOVA y el coeficiente de correlación de Pearson. Se utilizó un modelo multivariable para obtener estándares de desviación en todas las posibles combinaciones de valores, construyendo percentiles de normalidad ($x \pm DE$ y $x \pm 2 \cdot DE$).

Resultados: participaron en el estudio 136 niños (35% de los niños del municipio); el 54,4% fueron chicas. La distribución de estado de peso fue: 25 niños con bajo peso (18,4%), 54 normopeso (39,7%), 12 en riesgo de sobrepeso (8,8%), 22 con sobrepeso (16,2%) y 23 con obesidad (16,8%). La media de edad fue 5,7 años y de actividad física, 592 cuentas/minuto. Los chicos realizaron mayor actividad física ($p = 0,031$) y los niños con bajo peso y normopeso realizaron mayor actividad física que los niños con sobrepeso y obesidad ($p = 0,012$). No hubo diferencias significativas respecto a la edad. El análisis multivariable mostró diferencias significativas ($p < 0,001$) respecto al sexo y el estado de peso. En los niños, la actividad física decreció cuando aumentó el peso. Por el contrario, las niñas en grupos de IMC extremos obtuvieron mayores niveles de actividad física.

Conclusión: los preescolares en situación de sobrepeso y obesidad presentaron niveles menores de actividad física que los niños normopeso. Los niveles de actividad física fueron mayores en los chicos.

Palabras clave:

Preescolar.
Actividad física.
Obesidad pediátrica.
Distribución normal.

Received: 02/07/2017 • Accepted: 25/10/2017

Gutiérrez-Hervás A, Cortés-Castell E, Juste-Ruiz M, Palazón-Bru A, Gil-Guillén VF, Rizo-Baeza MM. Physical activity values in two- to seven-year-old children measured by accelerometer over five consecutive 24-hour days. *Nutr Hosp* 2018;35:527-532

DOI: <http://dx.doi.org/10.20960/nh.1403>

Correspondence:

Ernesto Cortés Castell. Department of Pharmacology, Pediatrics and Organic Chemistry, Miguel Hernández University. Ctra. de Valencia-Alicante, s/n. 03550 San Juan de Alicante, Alicante. Spain
e-mail: ernestocort@gmail.com

INTRODUCTION

The global prevalence of overweight and obesity in pre-school-age children has increased dramatically since 1990, from approximately 4% in 1990 to 7% in 2010 (1). Furthermore, obesity was more prevalent in children aged 4-5 years (18.3%) and overweight was more prevalent in children aged 8-9 years (25.5%) (2). This great increase in the incidence of overweight and obesity at early ages results in a high prevalence of metabolic syndrome (3). Consequently, obesity is related to decreasing quality of life (4) and to increased healthcare costs due to associated comorbidities, equivalent to 7% of the gross domestic product in developed countries (5).

During childhood and adolescence, environmental factors are the main causes of obesity, due to the energy imbalance caused by a high-energy intake (6) and/or a low level of physical activity (7). Thus, it is crucial to promote preventive interventions in early childhood to impact on lifestyle and prevent the development of overweight and obesity (7).

Over the past decade, a large body of empirical studies has described the correlates and determinants of physical activity in preschoolers (8-11). Though several tools exist to estimate physical activity approximately, these tools are not so effective at providing a more accurate determination, whether daily or over a certain period (12). For example, physical activity questionnaires obtain a subjective estimation of physical activity. In this field, accelerometry has been shown to be the most reliable method to record the quantity and level of physical activity that each subject performs in each period (13). Accordingly, accelerometers are considered to be the gold standard for free-living physical activity assessment (14).

A meta-analysis of several studies reporting accelerometry-derived daily physical activity levels of preschool-age children concluded that is necessary to develop physical activity guidelines for this range of age due to the confusion between several different cut-off points applied in the literature (15). Consequently, an equation based on these cut points in preschoolers was created (16).

Furthermore, physical activity levels have been assessed in different ways. Some authors define these levels following cut points using ROC curves over 3-7 minutes performing different physical activities in three different sessions (17). Others have measured physical activity levels in various ways: for three minutes using interclass correlation coefficients to modify Children's Activity Rating Scales, obtaining different cut points for three-, four-, and five-year-old children (18); for 20 minutes using the standard deviation (SD) of the counts and VO_2 , random-coefficient models and multivariate models for gender, race, age, height, quadratic terms for age and height and interactions between these variables and accelerometry counts, determining moderate-to-vigorous physical activity cut points at 420 counts/15 seconds and vigorous physical activity cut points at 842 counts/15 seconds (19); and for an average of 100 minutes, constructing physical activity cut points by ROC analysis, specifically to determine sedentary behaviors at $< 1,100$ counts/minute (20). A more recent study determined physical activity cut points for preschoolers during

a physical exercise session over four consecutive days (during activity time) using the Children's Activity Rating Scales and ROC curve and the ANOVA and Chi-squared test to determine differences between physical activity cut points, obtaining light physical activity at 373 counts/15 seconds, moderate physical activity at 585 counts/15 seconds and vigorous physical activity at 881 counts/15 seconds (21).

Nevertheless, these physical activity levels have recently been revised. The authors concluded there is a need to unify these cut points to determine the true physical activity of these children through methodological studies to advance the understanding of physical activity in this age group (22). Previous studies have focused on analyzing the amount of time that children spend each day in moderate-to-vigorous physical activity, not their total daily physical activity (23). Thus, physical activity such as playing games or in park areas that does not involve moderate-to-vigorous physical activity is not included in these activity measurements though it could be important in the quantification of total physical activity at the end of the day.

Determining reference values for this population is difficult because the variables that could influence physical activity do not use the same time period to determine cut points, or the same hours per day and the same days per week to analyze the physical activity of the children, even though some authors indicate the need to obtain data for ten hours per day to be considered as a valid day (23). The choice of accelerometer cut points can result in large discrepancies (21). Also, determining true physical activity in children requires recording over more days, with the best option being the analysis of consecutive days (8) and throughout the entire day. For this reason, we examined physical activity in two- to seven-year-old children, 24 hours per day over five consecutive days (three workdays and two weekend days) with the aim of establishing daily reference values without the influence of variables. The hypothesis of this study was that by measuring physical activity during this period accurate physical activity levels of preschool children would be obtained.

MATERIALS AND METHODS

STUDY POPULATION

Rafal is a small town of 3,091 inhabitants in Alicante, Spain. It is situated in a rural area in which most of the population lives off the cultivation of fruits and vegetables and the production of wine. Consequently, access to food representative of the Mediterranean diet, such as fruit and vegetables, is not an issue. The socioeconomic status of the families was medium-low, and 22.5% were children of immigrant parents, mostly of Moroccan origin.

STUDY DESIGN AND PARTICIPANTS

We undertook a cross-sectional observational study with quantitative analysis between September 2014 and June 2015, inviting

the entire population of children aged between two and seven years (through their parents), either from the pediatric office at the health center, the school or the two kindergartens in the municipality. The parents of all the children read and signed the consent form. Children were excluded if they had any disease, such as diabetes or Down syndrome, which could result in lack of control with the accelerometer.

A nutritionist took the children to the multipurpose classroom in small groups of four. The children who were measured in the pediatric examination room were accompanied by their parents. Weight and height were recorded with the children in underwear and without shoes. Next, accelerometers were placed on the abdominal area of the participants, and the nutritionist explained that the accelerometer could only be removed to aquatic activities (swimming or bathing). The parents helped during this time to ensure the accelerometers remained in place without interruptions and to record the time the child went to bed and woke up as well as any interruption during this period.

VARIABLES AND MEASUREMENTS

The main outcome variable in this study was 24-hour-per-day physical activity. Physical activity was measured for 120 uninterrupted hours (three workdays and two weekend days) by accelerometer (counts/minute), validated to determine physical activity in this age range (12,24,25). The data procedure was recorded every 15 epochs to maximize opportunities to more accurately capture the sporadic nature of young children's physical activity (26). A valid day may include at least ten hours per day (23), and a minimum of four consecutive days (three weekdays and one weekend day) were required to include a child (27). Half an hour of consecutive zeros was established as inactivity. To determine non-wear time as aquatic activity or some periods of time that children refuse to wear the accelerometer, these periods of time were compared with parent's reports, as well as sleep time, at night and day (27), which was excluded to calculate the mean of counts/minute of the sample.

Secondary variables were weight status (determined by the body mass index [BMI] Z-score), gender, and age. BMI (kg/m^2) was calculated from the weight and height parameters obtained, and the BMI Z-score was calculated in relation to age and gender using the Seinaptraker program (28), based on the Orbegozo Foundation standards (29). Children under five years were classified by BMI Z-score into five subgroups: underweight Z-score < -1 , normal-weight Z-score ($[-1]-1$), risk of overweight Z-score > 1 to ≤ 2 , overweight Z-score > 2 to ≤ 3 and obesity Z-score > 3 . On the other hand, children over five years were classified by BMI Z-score into four subgroups: underweight Z-score < -1 , normal-weight Z-score ($[-1]-1$), overweight Z-score > 1 to ≤ 2 and obese Z-score > 2 , according to World Health Organization (WHO) recommendations (30).

Auxological parameters were measured by two trained nutritionists. Tools used were: Seca® weighing-scales (761 Class III, accuracy 0.5 kg), a Harpenden stadiometer (Holtain Ltd.,

Crymych, Dyfed, UK), which accurately determines height within 0.1 cm, and 13 accelerometers (Actigraph™ GT1M and ActiLife program).

SAMPLE SIZE

Sample size was calculated to estimate the mean of our main variable, counts per minute (cpm). Since the analysis was carried out in volunteers, sample size was calculated *a posteriori*. As the sample was selected from the total study population ($n = 391$) with an expected SD of 170 cpm (31), an accuracy in the determination of the population mean of 23.07 cpm was obtained.

STATISTICAL ANALYSIS

The descriptive analysis was performed through absolute and relative frequencies for qualitative variables, while the mean and SD were used for quantitative variables. The relationship between the main variable and secondary variables was calculated using the t-test, ANOVA and the Pearson correlation coefficient. A linear regression model was constructed with the dependent variable of physical activity (in cpm) and the independent variables of weight status, age and gender, considering possible interactions between them. Non-significant interactions were eliminated from the model. The goodness-of-fit of the model was obtained through ANOVA. The model was used to obtain the SD from all the possible combinations of values (for example, children with childhood obesity). With each of the SD, normal percentiles were computed ($\bar{x} \pm \text{SD}$ and $\bar{x} \pm 2\text{-SD}$). For each relevant parameter its associated confidence interval (CI) was calculated. The type I error was set at 5%. IBM SPSS Statistics 24 software was used.

In the ANOVA and normal distribution test and the determination of percentiles of normality for daily physical activity, the BMI groups of risk of overweight and overweight were unified due to the sample size of risk of overweight children.

ETHICAL ISSUES

The study protocol was approved by the Ethics Committee of the University of Alicante prior to initiation (18 March, 2014). The researchers informed the parents about the study, the ethical principles and confidentiality, and written consent was provided by the parents of the participants.

RESULTS

We offered the entire population in this age range (391 children) the opportunity to participate, but 252 of them refused (through their parents) and two children met the exclusion criteria. Thus, the final sample included 136 children (35% of the children in this age range in the municipality).

Table I shows the weight status distribution of the sample of children studied: 25 underweight children (18.4%), 54 with normal weight (39.7%), 12 with risk of overweight (8.8%), 22 overweight (16.2%) and 23 obese (16.8%). Consequently, almost half the sample had excess weight and one in four preschool children was obese. The median age of the participants was 5.7 years (IQ 2.5) and the range of age was 2.4 to 7.3 years. Table I also shows their physical activity as recorded by accelerometer. In the entire sample, the mean was 592 cpm, with a SD of 125 cpm. Concerning gender, weight status group and age, the boys undertook more physical activity, underweight and normal-weight children undertook more physical activity in comparison with the overweight and obese groups, and there were no significant differences by age.

In the multivariate analysis (Table II), the integration of gender and weight status showed significant differences. Specifically, boys had a lower weight status than girls. From the normal distribution analysis, the mean plus two levels of SD for gender and weight status were obtained (Table III). When the number of cpm measured in a child exceeded two SD, above or below, the physical activity level was considered as abnormal. Figure 1 illustrates this distribution. In boys, physical activity decreased as weight status increased. In contrast, the girls in the extreme weight status groups (underweight and obese) undertook greater amounts of physical activity.

Table I. Descriptive and inferential analysis of physical activity in children aged two to seven years

Variable	Total n = 136 n (%) / x ± SD	Cpm x ± SD/r	p-value
Cpm	N/A	592 ± 125	N/A
<i>Gender:</i>			
Boys	62 (45.6)	618 ± 135	0.031
Girls	74 (54.4)	571 ± 113	
<i>Weight status:</i>			0.012
Underweight	25 (18.4)	604 ± 128	
Normal weight	54 (39.7)	627 ± 118	
Risk of overweight*	12 (8.8)	555 ± 154	
Overweight	22 (16.2)	563 ± 130	
Obese	23 (16.9)	538 ± 114	
Age (years)	5.5 ± 1.5	-0.064	0.458

Cpm: counts per minute; n (%): absolute frequency (relative frequency); N/A: not applicable; r: Pearson correlation coefficient; x ± SD: mean ± standard deviation.
*This category of weight status includes only children under five years.

DISCUSSION

A significant association was found between a higher level of physical activity and a healthier weight status and gender. The

Table II. Multivariate analysis of physical activity in children aged two to seven years

Variable	B (95% CI)	p-value
Intercept	567.3 (494.8, 639.7)	< 0.001
Weight status*	1.26 (-25.2, 27.7)	0.925
Boys	191.2 (82.5, 299.8)	< 0.001
Boys · Weight status*	-52.4 (-90.4, -14.4)	0.007

B: regression coefficient; CI, confidence interval. *: defined as: 1, underweight; 2, normal weight; 3 and 4, risk of overweight and overweight; and 5, obese.
Goodness-of-fit of the model (ANOVA test): $F = 6.35$, $p < 0.001$.

multivariate regression model produced curves of normal physical activity values for both genders and considering weight status.

The relationship between physical activity and weight status has been shown in school-age youth (32) and preschool children (9). However, it is difficult to compare physical activity between studies due to the considerable variation of prevalence estimates used to quantify this physical activity (22). A review concluded that a valid day may include at least ten hours per day, eliminating inactive hours in which children were asleep and aquatic activities (23).

In our case, we measured 120 consecutive hours distributed over workdays and weekend days, eliminating an average of ten hours of total inactivity (sleep). The mean physical activity in our sample was approximately 1,000 cpm. This value situates our sample in the light activity category, according to recent studies (19,33,34), and confirms that young children are not achieving an adequate amount of physical activity (27). However, we believe that uninterrupted physical activity measurements afford children and their families more freedom of activity, trying to measure a more accurate physical activity. When only a short period in which children are doing physical exercise is measured, these children could be forced to do more physical activity than they normally would during a day, resulting in a biased measurement. Furthermore, most authors have focused their studies on quantifying the number of minutes that preschool children spend in moderate-to-vigorous physical activity, following public health recommendations (15,16,19,21,26,35). However, a recent review concludes that the decision to select from among various cut points significantly influences the optimal levels of moderate-to-vigorous physical activity for obesity prevention in children (36). Moreover, none of the studies reviewed considered the weight status of preschool children to determine their physical activity levels or to provide physical activity level recommendations to prevent childhood obesity.

We measured total physical activity, which provides more accurate values for physical activity levels in preschool-age children during their normal daily life. Reference values are given by gender and weight status, which provide clinicians with standardized physical activity level values with which to compare their patients. Since we considered total daily physical activity, these values could be a reference, adaptable to the special needs of each group of children depending on their characteristics. According to sex,

Table III. Normal distribution of physical activity (in counts per minute) in children aged two to seven years, considering weight status and gender

Weight status	Boys					Girls				
	x-2·SD	x-SD	x	x + SD	x + 2·SD	x-2·SD	x-SD	x	x + SD	x + 2·SD
Underweight	650	679	707	736	764	519	544	569	593	618
Normal weight	620	638	656	675	693	539	554	570	585	601
Risk of overweight and overweight	574	590	605	620	636	541	556	571	586	601
Obese	508	531	554	577	600	525	548	572	596	620

X: mean; SD: standard deviation.




Figure 1.

Percentiles for normality for daily physical activity (counts per minute) according to the weight of the child. Thick lines denote mean \pm 2 · SD, and continuous lines denote mean \pm SD. *The overweight group includes children in risk of overweight and overweight, depending of their range of age.

boys were more active than girls (27) and overweight and obesity groups presented a lower amount of physical activity, especially in the boys. For this reason, activities requiring more effort in these children should be promoted during both school (37) and family time (38). This is of great importance given the epidemiologic problem in countries such as Spain, where the prevalence and incidence of obesity have constituted an alarming situation in the population in this age range during the last decades (39).

The most significant contribution of the study is the reference values obtained according to gender and weight status, which provide clinicians with the standardized daily physical activity levels of preschool-age children. Since we considered the total daily physical activity, these values could be used as a reference and adapted to the special necessities of each child depending on their weight status. In future studies, the use of accelerometers throughout the entire day is recommended to quantify physical activity and to choose an appropriate physical exercise intervention to improve weight status in preschool-age children (7,27,40).

The strength of this study is the evidence shown about the association between the level of daily physical activity in Spanish

preschool children, weight status and gender. By monitoring the children 24 hours per day and measuring workdays and weekend days we could obtain a more accurate determination of total physical activity in preschool children. This is of great importance given the need to study the population in this age range and the need for reliable data to quantify the true level of physical activity.

The main limitation of this study is that it was carried out in a small population. Therefore, it is necessary to replicate the study in other communities before the results can be extrapolated to the general population in this age range.

ACKNOWLEDGMENTS

The authors thank the Rafal town hall for their cooperation and Maria Repice and Ian Johnstone for their review of the English version of this paper.

ETHICAL STANDARDS

All human and animal studies have been approved by the appropriate Ethics Committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki and its later amendments.

In addition, all people gave their informed consent prior to their inclusion in the study.

REFERENCES

1. Wang Y, Lim H. The global childhood obesity epidemic and the association between socio-economic status and childhood obesity (Review). *Int Rev Psychiatry* 2012;24:176-88.
2. Valdés-Pizarro J, Royo-Bordonada MA. Prevalence of childhood obesity in Spain: National Health Survey 2006-2007. *Nutr Hosp* 2012;27:154-60.
3. Gutiérrez-Hervás AI, Rizo-Baeza MM, Martínez-Amorós N, Cortés-Castell E. Systolic pressure, abdominal obesity and body fat, metabolic syndrome predictors in Spanish preschoolers. *Nutr Hosp* 2015;31(5):2109-14. DOI: 10.3305/nh.2015.31.5.8685
4. Michels N, Susi K, Marqués-Vidal PM, Nydegger A, Puder JJ. Psychosocial quality-of-life, lifestyle and adiposity: a longitudinal study in pre-schoolers (Ballabeina study). *Int J Behav Med* 2016;23:383-92.
5. Frew E. Economic evaluation of childhood obesity interventions: reflections and suggestions. *Pharmacoeconomics* 2016;34:733-40.

6. Hasnain SR, Singer MR, Bradlee ML, Moore LL. Beverage intake in early childhood and change in body fat from preschool to adolescence. *Child Obes* 2014;10:42-9.
7. Moreno LA, Bel-Serrat S, Santaliestra-Pasias AM, Rodríguez G. Obesity prevention in children. Review. *World Rev Nutr Diet* 2016;106:119-26.
8. Butte NF, Wong WW, Wilson TA, Adolph AL, Puyau MR, Zakeri IF. Revision of dietary reference intakes for energy in preschool-age children. *Am J Clin Nutr* 2014;100:161-7.
9. Carson V, Clark D, Orgen N, Harber V, Kuzik N. Short-term influence of revised provincial accreditation standards on physical activity, sedentary behavior, and weight status in Alberta, Canada child care centers. *Early Child Educ J* 2015;43:459-65.
10. Creamer M, Decker E, De Bourdeaudhuij I, Verloigne M, Manios Y, Cardon G. The translation of preschoolers' physical activity guidelines into a daily step count target. *J Sports Sci* 2015;33:1051-7.
11. Hinkley T, Crawford D, Salmon J, Okely AD, Hesketh K. Preschool children and physical activity: a review of correlates. *Am J Prev Med* 2008;34:435-41.
12. Aguilar-Cordero MJ, Sánchez-López AM, Guisado-Barrilao R, Rodríguez-Blanco R, Noack-Segovia J, Pozo-Cano MD. Descripción del acelerómetro como método para valorar la actividad física en los diferentes periodos de la vida. *Nutr Hosp* 2014;29:1250-61.
13. Aguilar-Cordero MJ, González-Jiménez E, García-García CJ, García-López PA, Álvarez-Ferre J, Padilla-López CA, et al. Obesidad de una población de escolares de Granada: evaluación de la eficacia de una intervención educativa. *Nutr Hosp* 2011;26:636-41.
14. Pate RR, O'Neill JR, Mitchell J. Measurement of physical activity in preschool children. *Med Sci Sports Exerc* 2010;42:508-12.
15. Bornstein DB, Beets MW, Byun W, McIver K. Accelerometer-derived physical activity levels of preschoolers: a meta-analysis. *J Sci Med Sport* 2011;14:504-11.
16. Bornstein DB, Beets MW, Byun W, Welk G, Bottai M, Dowda M, et al. Equating accelerometer estimates of moderate-to-vigorous physical activity: in search of the Rosetta Stone. *J Sci Med Sport* 2011;14:404-10.
17. Evenson KR, Catellier DJ, Gill K, Ondrak KS, McMurray RG. Calibration of two objective measures of physical activity for children. *J Sports Sci* 2008;26:1557-65.
18. Sirard JR, Trost SG, Pfeiffer KA, Dowda M, Pate RR. Calibration and evaluation of an objective measure of physical activity in preschool children. *J Phys Act Public Health* 2005;2:345-57.
19. Pate RR, Almeida MJ, McIver KL, Pfeiffer KA, Dowda M. Validation and calibration of an accelerometer in preschool children. *Obesity* 2006;14:2000-6.
20. Reilly JJ, Coyle J, Kelly L, Burke G, Grant S, Paton JY. An objective method for measurement of sedentary behavior in 3- to 4-year olds. *Obes Res* 2003;11:1155-18.
21. Van Cauwenberghe E, Labarque V, Trost SG, Bourdeaudhuij I, Cardon G. Calibration and comparison of accelerometer cut points in preschool children. *Int J Pediatr Obes* 2005;6:e582-9.
22. Hnatiuk A, Salmon J, Hinkley T, Okely AD, Trost S. A review of preschool children's physical activity and sedentary time using objective measures. *Am J Prev Med* 2014;47:487-97.
23. Tudor-Locke C, Cahmi SM, Troiano RP. A catalog of rules, variables, and definitions applied to accelerometer data in the National Health and Nutrition Examination Survey, 2003-2006. *Prev Chronic Dis* 2012;9:110332.
24. Arvidsson D, Fitch M, Hudes ML, Tudor-Locke C, Fleming SE. Accelerometer response to physical activity intensity in normal-weight versus overweight African American children. *J Phys Act Health* 2001;8:682-92.
25. Pulsford R, Cortina-Borja M, Rich C, Kinnafick FE, Dezaux C, Griffiths LJ. Actigraph accelerometer-defined boundaries for sedentary behaviour and physical activity intensities in 7 year old children. *PLoS One* 2006;6:e21822.
26. Cliff D, Okely A, Smith L, Mckeen K. Relationships between fundamental movement skills and objectively measured physical activity in pre-school children. *Pediatr Exer Sci* 2009;21:436-9.
27. Hinkley T, Salmon J, Okely AD, Crawford D, Hesketh K. Preschoolers' physical activity, screen time, and compliance with recommendations. *Med Sci Sports Exerc* 2012;44(3):458-65.
28. Sociedad Española de Investigación en Nutrición y Alimentación en Pediatría-SEINAP. Nestlé Nutrition Institute and InterCath Medical-One. Aplicación Informática para gestión de pacientes y cálculos auxológicos y nutricionales en pediatría. Barcelona, Spain: Nestlé Nutrition Institute; 2007.
29. Sobradillo B, Aguirre A, Aresti U, Bilbao A, Fernández-Ramos C, Lizárraga A, et al. Curvas y tablas de crecimiento (Estudio longitudinal y transversal). Bilbao, Spain: Fundación Faustino Orbeagoza Eizaguirre; 1988.
30. World Health Organization. Training course on child growth assessment. WHO Child Growth Standards. Module C - Interpreting Growth Indicators. 2008. Available from: http://www.who.int/childgrowth/training/c_interpretando.pdf
31. Chaput JP, Lambert M, Methieu ME, Tremblay MS, Loughlin JO, Tremblay A. Physical activity vs. sedentary time: independent association with adiposity in children. *International Association for the Study of Obesity. Pediatr Obes* 2012;7:251-8.
32. Strong WB, Malina RM, Blimkie CJ, Daniels SR, Dishman RK, Gutin B, et al. Evidence based physical activity for school-age youth. *J Pediatr* 2005;146:732-7.
33. España-Romero V, Mitchell JA, O'Neill JR, Pate RR. Objectively measured sedentary time, PA and markers of body fat in preschool children. *Pediatr Exer Sci* 2013;25:154-63.
34. Puyau MR, Adolph AL, Vohra FA, Butte NF. Validation and calibration of physical activity monitors in children. *Obes Res* 2002;10:150-7.
35. Costa S, Barber SE, Cameron N, Clemes S. Calibration and validation of the ActiGraph GT3X+ in 2-3 year old. *J Sci Med Sport* 2014;17:617-22.
36. Gába A, Dygrýn J, Mitás J, Jakubec L, Frömel K. Effect of accelerometer cut-off points on the recommended level of physical activity for obesity prevention in children. *PLoS One* 2016;11:e0164282.
37. Vanderloo LM, Tucker P. Weekly trends in preschoolers' physical activity and sedentary time in childcare. *Int J Environ Res Public Health* 2015;12:2454-64.
38. Ruiz R, Gesell SB, Buchowski MS, Lambert W, Barkin SL. The relationship between Hispanic parents and their preschool-aged children's physical activity. *Pediatrics* 2011;127:888-95.
39. Ruiz-Pérez L, Zapico M, Zubiaur A, Sánchez-Paya J, Flores J. Increase in the prevalence of overweight and obesity in the pediatric population of the province of Alicante (Spain) in the last 10 years. *Endocrinol Nutr* 2008;5:389-95.
40. Ogata BN, Hayes D. Position of the Academy of Nutrition and Dietetics: nutrition guidance for healthy children ages 2 to 11 years. *J Acad Nutr Diet* 2014;114:1257-76.