



Nutrición Hospitalaria



Trabajo Original

Valoración nutricional

Hydration habits before, during and after training and competition days among amateur basketball players

Hábitos de hidratación antes, durante y tras el ejercicio y la competición en jugadores de baloncesto amateurs

Maria del Mar Bibiloni¹, Eulàlia Vidal-García², Marta Carrasco², Alicia Julibert¹, Antoni Pons¹ and Josep A. Tur¹

¹Research Group on Community Nutrition and Oxidative Stress. University of the Balearic Islands and CIBEROBN (Physiopathology of Obesity and Nutrition). Palma de Mallorca, Spain. ²Blanquerna's School of Health Sciences. Ramon Llull University. Barcelona, Spain

Abstract

Background: fluid intake before, during and after exercise is the most important way to replace water lost during exercise and avoid dehydration.

Aim: to assess fluid intake habits before, during and after exercise in amateur basketball players on both training and competition days.

Methods: one hundred and eighty-three amateur basketball players (87 women and 96 men, 19-29 years old) were interviewed. Data was obtained from a drinking habits questionnaire.

Results: overall, 20.8%, 5.5% and 2.7% of subjects did not drink before, during and after exercise on training days, respectively; 17.5% of subjects did not drink before exercise on competition days. Water was the preferred beverage before, during and after exercise on both training and competition days, with fruit juice being the second most consumed beverage before exercise. All subjects also drank fizzy drinks and other beverages during exercise on competition days. According to the recommendations, good hydration habits were found in 54.6%, 74.2% and 76.5% of subjects before, during and after training days, respectively.

Conclusions: most amateur basketball players drink before, during and after exercise on both training and competition days, but not all of them complied with the hydration recommendations.

Key words:

Adults. Liquid intake. Survey. Hydration recommendations.

Resumen

Introducción: la ingesta de líquidos antes, durante y tras el ejercicio es la forma más importante de recuperar el agua perdida durante el ejercicio y de evitar la deshidratación.

Objetivo: analizar los hábitos de ingesta de líquidos, durante y tras el ejercicio en jugadores de baloncesto amateurs tanto durante el entrenamiento como durante la competición.

Métodos: fueron entrevistados 183 jugadores de baloncesto amateurs (87 mujeres y 96 hombres, de 19-29 años de edad). Los datos se obtuvieron por aplicación de un cuestionario de hábitos de bebida.

Resultados: globalmente, el 20,8%, 5,5% y 2,7% de los sujetos no bebieron antes, durante y tras el entrenamiento, respectivamente; el 17,5% de los sujetos no bebieron antes de la competición. El agua fue la bebida preferida antes, durante y tras el ejercicio, tanto durante el entrenamiento como durante la competición, en tanto que los zumos de frutas fueron la segunda bebida consumida antes del ejercicio. Todos los sujetos tomaron bebidas gaseosas y de otro tipo durante la competición. De acuerdo con las recomendaciones, se encontraron buenos hábitos de hidratación en el 54,6%, 74,2% y 76,5% de los sujetos antes, durante y tras el entrenamiento, respectivamente.

Conclusiones: la mayoría de los jugadores de baloncesto amateurs beben antes, durante y tras el ejercicio, tanto durante el entrenamiento como durante la competición, pero no todos ellos cumplen con las recomendaciones de hidratación.

Palabras clave:

Adultos. Ingesta líquida. Entrevista. Recomendaciones de hidratación.

Received: 25/07/2017 • Accepted: 05/01/2018

Authors' contributions: MMB, EV and JAT conceived, designed, devised and supervised the study. EV and MC collected and supervised the samples. MMB, AJ, MC, EV and JAT analyzed the data and wrote the manuscript. AP and JAT obtained funds. All authors read and approved the final manuscript.

Bibiloni MM, Vidal-García E, Carrasco M, Julibert A, Pons A, Tur JA. Hydration habits before, during and after training and competition days among amateur basketball players. *Nutr Hosp* 2018;35:612-619
DOI: <http://dx.doi.org/10.20960/nh.1462>

Correspondence:

Josep A. Tur. Research Group on Community Nutrition and Oxidative Stress. Universitat de les Illes Balears. Campus Guillem Colom Bldg. E-07122 Palma de Mallorca, Spain
e-mail: pep.tur@uib.es

INTRODUCTION

A balanced diet and appropriate hydration are fundamental to sport performance, but specific needs will depend on several factors, such as individual physiological conditions, intensive physical activity, time of the season and training or competition period (4,5). An average daily water intake of 2.2 to 2.6 ml/d in men and 1.9 to 2.4 ml/d in women meets the needs of most adult people (1). However, strenuous physical exercise and heat stress can greatly increase daily water needs, and individual variability between athletes can be substantial (6).

Dehydration can negatively affect different physiological systems including the nervous system (7,8), cardiovascular, thermoregulation and endocrine systems or metabolism (3), which may have negative consequences on health (9), affect athletic performance (10-12) and increase the risk of exertional heat injury (13) in both anaerobic and aerobic sports (14). Physical and mental performance during physical exercise and sport practice is impaired in the under-hydrated individual (15) and to avoid this, as well as the negative effects on physical performance, the athlete must drink enough fluids before (16-18), during (19-21) and after practicing physical exercise, maintaining proper hydration throughout the day (3). Furthermore, not only the amount of beverage is important, but also the type of beverage to be drunk.

Currently, many people practice some kind of recreational intensive activity sport, such as basketball, for more than one hour a day (4), but there are few studies evaluating whether drinking habits are in line with the recommendations made by national and international sports institutions. Basketball is an aerobic-anaerobic sport, characterized by high-intensity intermittent exercise (22) and mainly played indoors, so dehydration may also be affected by high temperature and humidity. Health and disease prevention do not only depend on physical activity practice (23), but also on eating and hydration habits (20). Therefore, the aim of this work was to assess the drinking habits of amateur basketball players before, during and after exercise on both training and competition days.

METHODS

STUDY DESIGN

The study was a population-based cross-sectional study carried out in the city of Barcelona between October 2011 and February 2012.

SELECTION OF PARTICIPANTS, RECRUITMENT AND APPROVAL

Barcelona city's basketball clubs ($n = 50$) were contacted by e-mail and phone and all players registered as senior category (1st, 2nd and 3rd) in the Catalan Federation of Basketball (CFB) and aged between 19 to 29 years were invited to participate in the study. One hundred and eighty-three players (96 men and 87 women), from 45 clubs, performing intensive physical activity 2-3 times per week, accounting for 60 to 120 min/d, agreed to participate.

ETHICS

This study was conducted according to the guidelines laid down in the Declaration of Helsinki, and the URL's Committee of Ethics in Research (CER-URL, Barcelona, Spain, ref. 2010_05) approved all procedures involving human subjects. Written informed consent was obtained from all subjects.

ANTHROPOMETRIC MEASUREMENTS

Anthropometric characteristics are part of the biological variables related to sport performance. Anthropometric characterization of subjects was included in table I. All anthropometric variables were measured according to the protocol recommended by the

Table I. Anthropometric characteristics of the amateur basketball players

	Women (n = 87)	Men (n = 96)	p [‡]
Missing (n)	1	4	
Age (years)*	22.4 (20.1-24.6)	21.7 (20.4-25.3)	0.883
Body mass (kg)*	61.7 (56.7-67.1)	78.7 (71.4-83.9)	< 0.001
Height (m)*	169.0 (164.8-173.0)	184.5 (178.0-190.0)	< 0.001
BMI (kg/m ²)*	21.9 (20.2-23.4)	23.0 (21.6-24.6)	< 0.001
WHR*	0.42 (0.41-0.45)	0.44 (0.42-0.47)	< 0.001
<i>Somatotype components</i>			
Endomorphic [†]	4.33 ± 1.07	3.06 ± 0.90	< 0.001
Mesomorphic [†]	2.84 ± 1.54	4.02 ± 1.16	< 0.001
Ectomorphic [†]	2.68 ± 1.22	2.92 ± 1.09	0.182

BMI: body mass index; WHR: waist-to-height ratio. Values are the *median (1st quartile-3rd quartile) and [†]mean ± standard deviation. [‡]Statistical significant differences between genders were assessed by means of unpaired Student t test or Mann-Whitney U test.

International Society for the Advancement of Kinanthropometry (ISAK) (24). All the anthropometric measurements were objectively obtained by trained personnel. Height was determined using a mobile anthropometer (Seca 217®, Hamburg, Germany), to the nearest millimeter, with the subject's head in the Frankfurt plane. Body mass was determined to the nearest 100 g using a digital scale (Seca 874®, Hamburg, Germany). The subjects were weighed in bare feet and light underwear. Height and weight measures were used to calculate the body mass index (BMI, kg/m²). Waist circumference (WC) was also measured. Height and WC measures were used to calculate the waist-to-height ratio (WHtR).

The three somatotype components (endomorphism, mesomorphism and ectomorphism) were calculated according to the Carter and Heath anthropometric somatotyping method (25). The ten variables used to calculate the anthropometric somatotype were height, mass, four skinfolds (triceps, subscapular, supraspinale, medial calf), biepicondylar breadth of humerus and femur, and arm (flexed and tensed) and calf circumferences. Skinfold thicknesses were measured using a skinfold caliper (Holtain®, Crosswell, UK), to the nearest 0.2 mm. Circumferences were measured using a non-stretch measuring tape (Lufkin Executive® w606pm, Lufkin, USA) to the nearest 1 mm. Bone widths were measured with a small sliding bone caliper (Holtain® Ltd., Crosswell, UK) to the nearest 1 mm. The subjects were asked to stand erect in a relaxed position with both feet together on a flat surface. The median was used for statistical analysis if the measurements had to be taken three times, while the mean was used if the first two measurements were within the acceptable range (26).

ASSESSMENT OF BEVERAGE CONSUMPTION

A drinking habits questionnaire based on a previously published questionnaire (27) was self-administered by computer. This questionnaire included the following questions about beverage intake before, during and after exercise on both training and competition days:

1. Type of beverage, which was categorized into five groups: fruit juice (i.e., natural fruit juice and fruit juice sweetened with sugar), water (i.e., tap, bottled and spring water), energy drinks (i.e., cola and *guarana* drinks), soda (i.e., carbonated soft drinks) and others (i.e., carrot juice, alcohol-free beer, chocolate, vanilla and strawberry milkshakes and diet milkshakes, soya milk, rice milk, oat milk, fermented milk drinks with sugar, fermented milk drinks, kefir and sweetened iced tea).
2. The amount of fluid drunk, which was categorized into three groups: < 250, 250-500, and ≥ 500 ml.
3. The beverage time, which was categorized into four groups before and after the exercise: < 10, 10-20, 20-30, and > 30 min; and into three groups during the exercise: 1-3, 4-6 and > 6 times. Fluid intake was only considered if it occurred within four hours before starting or after finishing physical exercise.

According to the drinking habits recommendations published by the American College of Sport Medicine (ACSM) (28,29) and the

Spanish Federation of Sports Medicine (FEMEDE) (4), the following criteria were established to assess healthy hydration habits:

1. Before exercise: less than one glass (< 250 ml) between 10-30 minutes beforehand, or more (≥ 250 ml), 30 minutes before exercise.
2. During exercise: less than one glass (< 250 ml) more than six times; between one to two glasses (250 to 500 ml) 4-6 times; or more than two glasses (≥ 500 ml) regardless of the frequency.
3. After exercise: more than one glass (≥ 250 ml) within 30 minutes after exercise ended.

STATISTICS

Analyses were performed with the Statistical Package for the Social Sciences version 24.0 (IBM SPSS Statistics for Windows, Armonk, NY: IBM Corp USA). All tests were stratified by gender. Categorical variables were presented as frequencies and/or proportions. Significant differences in prevalence were calculated by means of χ^2 or Fisher's exact test. Normality of data was assessed using Kolmogorov-Smirnov test. Continuous variables were expressed as follows: mean (standard deviation [SD]) for normally distributed data (somatotype components: endomorphy, mesomorphy, ectomorphy), and median (1st-3rd quartile) for non-normally distributed data (age, weight, height, BMI, and WHtR). For normally distributed data, the unpaired Student's t-test was used to compare the mean of two independent groups. For non-normally distributed data, the Mann-Whitney U test was used to compare the median of two independent groups. The level of significance was established for p values at < 0.05.

RESULTS

Table II shows the type of beverage consumed before, during and after exercise on both training and competition days. Overall, 20.8%, 5.5% and 2.7% of the subjects reported that they did not drink before, during and after exercise on training days, respectively. More women than men reported that they did not drink before exercise (27.6% vs 14.6%, $p < 0.05$) while more men than women did not drink during exercise (9.4% vs 1.2%, $p < 0.05$). No statistically significant difference in the prevalence of non-drinkers was found between men and women after exercise (2.1% and 3.4%, respectively). In competition days, 17.5% of subjects reported that they did not drink before exercise, with a higher prevalence among women than in men (25.3% vs 10.4%, $p < 0.01$). Almost none of them reported that they did not drink during or after exercise on competition days.

Water was the preferred beverage before, during, and after exercise on both training and competition days. In fact, 64.5% of the subjects reported that they drank water before exercise on training days, followed by fruit juice (21.3%), soda (10.9%), energy drinks (6.0%) and other beverages (4.9%). Overall, a

Table II. Type of beverages consumed before, during and after the exercise in training and competition days among amateur basketball players

	n	Training day			Competition day		
		Before	During	After	Before	During	After
<i>Nothing</i>							
All	183	38 (20.8)	10 (5.5)	5 (2.7)	32 (17.5)	0 (0.0)	2 (1.1)
Women	87	24 (27.6)	1 (1.2)	3 (3.4)	22 (25.3)	0 (0.0)	1 (1.1)
Men	96	14 (14.6)	9 (9.4)	2 (2.1)	10 (10.4)	0 (0.0)	1 (1.0)
p		0.030	0.020	0.670	0.008	1.000	1.000
<i>Fruit juice</i>							
All	183	39 (21.3)	1 (0.5)	11 (6.0)	25 (13.7)	3 (1.6)	14 (7.7)
Women	87	19 (21.8)	1 (1.1)	4 (4.6)	10 (11.5)	1 (1.1)	6 (6.9)
Men	96	20 (20.8)	0 (0.0)	7 (7.3)	15 (15.6)	2 (2.1)	8 (8.3)
p		0.868	0.475	0.444	0.416	1.000	0.715
<i>Water</i>							
All	183	118 (64.5)	172 (94.0)	149 (81.4)	126 (68.9)	180 (98.4)	142 (77.6)
Women	87	51 (58.6)	85 (97.7)	75 (86.2)	54 (62.1)	86 (98.9)	68 (78.2)
Men	96	67 (69.8)	87 (90.6)	74 (77.1)	72 (75.0)	94 (97.9)	74 (77.1)
p		0.115	0.044	0.113	0.059	1.000	0.861
<i>Energetic drinks</i>							
All	183	11 (6.0)	2 (1.1)	21 (11.5)	15 (8.2)	8 (4.4)	25 (13.7)
Women	87	3 (3.4)	1 (1.1)	5 (5.7)	3 (3.4)	3 (3.4)	4 (4.6)
Men	96	8 (8.3)	1 (1.0)	16 (16.7)	12 (12.5)	5 (5.2)	21 (21.9)
p		0.165	1.000	0.021	0.026	0.723	0.001
<i>Soda</i>							
All	183	20 (10.9)	0 (0.0)	33 (18.0)	10 (5.5)	183 (100.0)	45 (24.6)
Women	87	10 (11.5)	0 (0.0)	11 (12.6)	6 (6.9)	87 (100.0)	25 (28.7)
Men	96	10 (10.4)	0 (0.0)	22 (22.9)	4 (4.2)	96 (100.0)	20 (20.8)
p		0.816	1.000	0.071	0.522	1.000	0.215
<i>Others</i>							
All	183	9 (4.9)	0 (0.0)	8 (4.4)	13 (7.1)	183 (100.0)	3 (1.6)
Women	87	6 (6.9)	0 (0.0)	2 (2.3)	10 (11.5)	87 (100.0)	2 (2.3)
Men	96	3 (3.1)	0 (0.0)	6 (6.3)	3 (3.1)	96 (100.0)	1 (1.0)
p		0.313	1.000	0.283	0.028	1.000	0.605
<i>≥ 2 type of beverages</i>							
All	183	48 (26.2)	2 (1.1)	39 (21.3)	35 (19.1)	8 (4.4)	45 (24.6)
Women	87	24 (27.6)	1 (1.1)	12 (13.8)	17 (19.5)	3 (3.4)	18 (20.7)
Men	96	24 (25.0)	1 (1.0)	27 (28.1)	18 (18.8)	5 (5.2)	27 (28.1)
p		0.691	1.000	0.018	0.892	0.723	0.243

Values are: n (%). Statistical significant differences between genders were assessed by χ^2 or Fisher's exact test.

quarter of the subjects reported more than one type of beverage before exercise on training days. On competition days, 68.9% of the subjects also reported that they drank water before exercise,

followed by fruit juice (13.7%), energy drinks (8.2%), other beverages (7.1%) and soda (5.5%). However, men were more likely to show a preference for energy drinks than women. Almost all of

the subjects reported that they only drank water during exercise on training days. However, all the subjects reported that they also drank soda and other beverages during exercise on competition days. Despite the fact that the preferred beverage after exercise on both training and competition days included water (81.4% and 77.6%, respectively), some subjects reported that they also drank soda (18.0% and 24.6%), energy drinks (11.5% and 13.7%), fruit juice (6.0% and 7.7%) and/or other beverages (4.4% and 1.6%). After exercise, men were also more likely to prefer energy drinks than women were.

The overall amount of beverage intake before, during and after exercise on both training and competition days is shown in figure 1. Overall, 1.4%, 41.4% and 57.2% of the subjects reported that they drank < 250, 250-500, and ≥ 500 ml, respectively, before exercise on training days, with no statistically significant differences between genders ($p = 0.265$). On competition days, 62.3% of subjects reported that they drank ≥ 500 ml before exercise. On the other hand, statistically significant differences in the distribution of the sample according to the amount of beverage intake during exercise were obtained on both training ($p = 0.006$) and competition ($p = 0.042$) days, with men more likely to drink ≥ 500 ml




Figure 1. Amount of beverage consumed before, during and after exercise in training (A) and competition (B) days among amateur basketball players.

than women were. After exercise, around 70% of the subjects reported that they drank ≥ 500 ml.

Figure 2 shows the time of fluid intake for those subjects who reported drinking some beverage before (A), during (B) and after (C) exercise on both training and competition days. Most of the subjects reported that they drank < 10 or > 30 minutes before exercise on both training and competition days. During exercise, most subjects (89.0%) reported that they drank 1-3 times on training days, while only 52.1% of men and 28.7% of women did




Figure 2. Time of the fluid intake among amateur basketball players who reported to drink several beverages before (A), during (B) and/or after (C) the exercise in training and competition days.

so 1-3 times on competition days. Furthermore, 21.9% of men and 25.3% of women reported that they drank > 6 times during exercise on competition days. Overall, 80.3% and 83.4% of the subjects reported that they drank < 30 minutes after exercise on training and competition days, respectively.

Table III shows the prevalence of good hydration habits before, during and after exercise on training days according to the recommendations released by the ACSM and FEMEDE. Good hydration habits were found in 54.6%, 74.2% and 76.5% of the subjects, respectively.

DISCUSSION

The first main finding of this study was that 20.8%, 5.5% and 2.7% of amateur basketball players reported that they did not drink before, during and after exercise on training days, respectively. Furthermore, 17.5% of the subjects reported that they did not drink before exercise on competition days, with a higher prevalence among women than in men. To our knowledge, there is little previous data regarding the drinking habits of amateur sportspeople. Previously, Alarcon et al. (27) assessed the drinking habits of 35 basketball players (14-32 years) from the Polaris World basketball Club (Murcia, Spain) on a competition day. In this study, 94.3% of the participants drank some fluid before exercise against the 82.5% from this study.

In agreement with the research by Alarcon et al. (27), water was the preferred beverage before, during, and after exercise on both training and competition days. Moreover, fruit juice was the second most-consumed beverage on both training and competition days

prior to exercise. However, in contrast to the study by Alarcon et al. (27), all the subjects in our work reported that they also drank soda and other beverages during exercise on competition days. Finally, beverage intake in the Alarcon et al. study (27) included water (40%), fruit juice (22.8%), soda (20%) and energy drinks (17.1%) after exercise. Likewise, in this study participants also reported that they drank water, soda, energy drinks, fruit juice and other beverages after exercise on both training and competition days. Finally, 31.8% of the subjects in our study reported that they drank some type of beverage 30 minutes before exercise on competition days, against the 8.6% seen in the Alarcon et al. study (27). Furthermore, 16.6% of the subjects in this study reported that they drank some type of beverage 30 minutes after exercise on competition days, against the 22.8% observed in the study by Alarcon et al. (27).

The National Athletic Trainers Association (NATA) (13) proposed that, to ensure proper pre-exercise hydration, athletes should consume approximately 500-600 ml of water or sport drinks 2-3 hours before exercise. If hydration levels are low, athletes should drink 200-300 ml of water or sport drinks 10-20 minutes before exercise. In this study, most subjects reported drinking at least 250 ml before exercise; however, only 26 subjects reported that they drank it 10-20 minutes before exercise. The NATA (13) recommended that, during exercise, fluid replacement generally requires 200-300 ml of water or sports drinks every 10-20 minutes and proposed that post-exercise hydration should aim, within two hours afterwards, to correct fluid losses accumulated during the practice or event and that, ideally, rehydration should contain water to restore hydration levels.

A review by Rowland (17) pointed out that the amount of liquid intake could be calculated as a fluid intake of

Table III. Hydration habits according to the recommendations released by the American College of Sports Medicine (ACSM) and the Spanish Federation of Sports Medicine (FEMEDE) among amateur basketball players in a training day

	Training day		
	All (n = 183)	Women (n = 87)	Men (n = 96)
<i>Before the exercise</i>			
≥ 250 ml; 10-30 min before the exercise	54 (29.5)	26 (29.9)	28 (29.2)
250-500 ml; > 30 min before the exercise	19 (10.4)	9 (10.3)	10 (10.4)
≥ 500 ml; > 30 min before the exercise	27 (14.8)	12 (13.8)	15 (15.6)
Total	100 (54.6)	47 (54.0)	53 (55.2)
<i>During the exercise</i>			
< 250 ml; > 6 times	1 (0.5)	0 (0.0)	1 (1.0)
250-500 ml; 4-6 times	1 (0.5)	1 (1.1)	0 (0.0)
≥ 500 ml; independently of times	134 (73.2)	58 (66.7)	76 (79.2)
Total	136 (74.2)	59 (67.8)	77 (80.2)
<i>After the exercise</i>			
≥ 250 ml within 30 min	140 (76.5)	64 (73.6)	76 (79.2)

Values are: n (%).

13 ml/kg body weight/h. According to the ACSM (28,29) and FEMEDE (4), prior to exercise individuals should drink beverages at least four hours before exercise and around 5-7 ml/kg body weight (300-600 ml) or 3-5 ml/kg body weight approximately two hours before exercise. In the present study, about 60% of the subjects reported that they drank at least 500 ml before exercise. However, post-exercise fluid intake is equally important to avoid commencing subsequent bouts of exercise in a dehydrated state (4,17,26). In fact, the ACSM (29) and FEMEDE (4) also pointed out that, 30 minutes after exercise starts, it is necessary to compensate fluid loss, and after one hour it is essential to drink 400-500 ml/h or 150-200 ml/20 min. Moreover, post-exercise hydration should aim to restore fluid losses accumulated during the practice or event within two hours. In this study, around 70% of the subjects reported that they drank at least 500 ml after exercise, but around 80% of basketball players rehydrated by drinking water and 25% drank also soda. Ideal rehydration should contain water to restore hydration levels. Finally, good hydration habits were found in 54.6%, 74.2% and 76.5% amateur basketball players before, during and after exercise on training days, respectively.

STRENGTHS AND LIMITATIONS

The main strength of this study is that there are many studies regarding the relationship between hydration and health and performance in the field of elite sport, but few studies have been conducted on adults in the field of recreational sports. However, this study has also several limitations. Firstly, good hydration habits should consider the intake of liquid throughout the day to learn about the level of hydration before starting exercise in order to determine the amount of liquid that each subject should drink after completing the exercise. This should be 150% (4) of the amount of weight lost during exercise. Secondly, the evaluation of good hydration was only carried out on a training day, because not all subjects played for more than one hour on a competition day.

CONCLUSIONS

Good hydration habits were found in 54.6%, 74.2% and 76.5% of subjects before, during and after the training day, respectively. It is difficult to instil hydration habits into all sportsmen, and even more so in non-professional categories and in the lower divisions. Because good hydration substantially contributes to the improvement of physical performance, and drinking water *ad lib* is not enough, hydration patterns should be an important part of training (30). These results should be taken into account to establish recommendations by sports organizations in relation to the amount and quality of liquid to drink before, during and after exercise, taking into account the type of sport, duration and climate.

FUNDING SOURCES

The study was supported by the Instituto de Salud Carlos III (Projects 14/00636 and 17/01827, Red Predimed-RETIC RD06/0045/1004, and CIBEROBN CB12/03/30038), Grant of support to research groups no. 35/2011 (Balearic Islands Gov.), EU COST Action CA16112, and EU FEDER funds. The authors also wish to thank the Catalan Basketball's Federation to help contact with the Barcelona Basketball Clubs, and Claudia Martinez, Elena Carrillo and Carla Not for assistance in the development of this survey.

REFERENCES

1. EFSA. Scientific Opinion on Dietary Reference Values for Water. 2010. Available from: <http://www.efsa.europa.eu/en/scdocs/scdoc/1459.htm>. Accessed on 12/13/2012.
2. Guyton AC. Tratado de fisiología médica. Madrid: Interamericana; 1984.
3. Iglesias-Rosado C, Villarino-Marin AL, Martínez JA, Cabrerizo L, Gargallo M, Lorenzo H, et al. Importance of water in the hydration of the Spanish population: FESNAD 2010 document. *Nutr Hosp* 2011;26(1):27-36.
4. Palacios N, Franco L, Manonelles P, Manuz B, Villegas J. Consensus on drinks for the sportsman. Composition and guidelines of replacement of liquids. Document of consensus of the Spanish Federation of Sports Medicine. *AMD* 2008;25(126):245-58.
5. Casa DJ, Clarkson PM, Roberts WO. American College of Sports Medicine roundtable on hydration and physical activity: consensus statements. *Curr Sports Med Rep* 2005;4(3):115-27.
6. Sawka MN, Cheuvront SN, Carter R. Human water needs. *Nutr Rev* 2005;63(6-2):S30-9.
7. Grandjean AC, Grandjean NR. Dehydration and cognitive performance. *J Am Coll Nutr* 2007;26(5 Suppl):549S-554S.
8. Maughan RJ, Shirreffs SM, Leiper JB. Errors in the estimation of hydration status from changes in body mass. *J Sports Sci* 2007;25(7):797-804.
9. Hillman AR, Vince RV, Taylor L, McNaughton L, Mitchell N, Siegler J. Exercise-induced dehydration with and without environmental heat stress results in increased oxidative stress. *Appl Physiol Nutr Metab* 2011;36(5):698-706.
10. Montain SJ, Smith SA, Mattot RP, Zientara GP, Jolesz FA, Sawka MN. Hypohydration effects on skeletal muscle performance and metabolism: a 31P-MRS study. *J Appl Physiol* 1998;84(6):1889-94.
11. Sawka MN, Cheuvront SN, Kenefick RW. High skin temperature and hypohydration impair aerobic performance. *Exp Physiol* 2012;97(3):327-32.
12. Below PR, Mora-Rodríguez R, González-Alonso J, Coyle EF. Fluid and carbohydrate ingestion independently improve performance during 1 h of intense exercise. *Med Sci Sports Exerc* 1995;27(2):200-10.
13. Casa DJ, Armstrong LE, Hillman SK, Montain SJ, Reiff RV, Rich BS, et al. National athletic trainers' association position statement: fluid replacement for athletes. *J Athl Train* 2000;35(2):212-24.
14. Ubiratan F. El efecto de la deshidratación en el rendimiento anaeróbico. *Rev Cienc Ej Salud* 2006;4(1):13-21.
15. Baker LB, Dougherty KA, Chow M, Kenney WL. Progressive dehydration causes a progressive decline in basketball skill performance. *Med Sci Sports Exerc* 2007;39(7):1114-23.
16. McMurray RG, Williams DK, Battaglini CL. The timing of fluid intake during an Olympic distance triathlon. *Int J Sport Nutr Exerc Metab* 2006;16(6):611-9.
17. Rowland T. Fluid replacement requirements for child athletes. *Sports Med* 2011;41(4):279-88.
18. Goulet ED. Dehydration and endurance performance in competitive athletes. *Nutr Rev* 2012;70(Suppl 2):S132-6.
19. Backhouse SH, Biddle SJ, Williams C. The influence of water ingestion during prolonged exercise on affect. *Appetite* 2007;48(2):193-8.
20. Miller VS, Bates GP. Hydration, hydration, hydration. *Ann Occup Hyg* 2010;54(2):134-6.
21. Williams M. Nutrition for health, fitness and sport. Barcelona: McGraw-Hill Companies; 2002.

22. World Health Organization. Estrategia mundial sobre régimen alimentario, actividad física y salud. Accessed on 12/12/2012. Available from: http://www.who.int/dietphysicalactivity/physical_activity_intensity/es/index.html
23. Warburton DE, Nicol CW, Bredin SS. Health benefits of physical activity: the evidence. *Can Med Assoc J* 2006;174(6):801-9.
24. Norton K, Whittingham N, Carter L, Kerr D, Gore C, Marfell-Jones M. Measurement techniques in anthropometry. In: Norton K, Olds T, eds. *Anthropometrica*. Sydney: UNSW Press; 1996. pp. 25-75.
25. Carter JEL, Heath BH. *Somatotyping: development and applications*. Cambridge: Cambridge University Press; 1990.
26. Ross W, Marfell-Jones M. Kinanthropometry. In: McDougall J, Wenger H, Green H, eds. *Physiological testing of the elite athlete*. Ithaca, NY: Movement Publications; 1982. pp. 223-308.
27. Alarcón López F, Ureña Ortin N, Piñar López M. Hábitos sobre hidratación durante la competición en baloncesto. *Efdeportes.com* 2009;137:1.
28. American Dietetic Association, Dietitians of Canada, American College of Sports Medicine, Rodríguez NR, Di Marco NM, Langley S. American College of Sports Medicine position stand. Nutrition and athletic performance. *Med Sci Sports Exerc* 2009;41(3):709-31
29. American College of Sports of Medicine. Exercise and fluid replacement. *Med Sci Sports Exerc* 2007;39(377):390.
30. Rush P, Gatti E. La hidratación en el básquetbol de primera de la FRBCF. *ISDe Sports Magazine* 2011;3(9):1-4.