



Trabajo Original

Nutrición en el anciano

Cost, microbiological, and nutritional properties of pureed food production in nursing homes. The ABADIA Study

Propiedades nutricionales, microbiológicas y costes de producción de las dietas de textura modificada en las residencias de ancianos. El estudio ABADIA

María D. Ballesteros-Pomar¹, Javier Pérez-Martín², María J. Mendiola³, José M García-García⁴, Soledad Parrado-Cuesta⁵, Ángel M. Caracuel⁶, Begoña Garcés⁷, Marta Comellas⁸, Héctor D. de Paz⁸, and Núria Barcons⁹

¹Unidad de Nutrición Clínica y Dietética. Sección de Endocrinología y Nutrición. Complejo Asistencial Universitario de León. León, Spain. ²Residencia de Atención a Personas Dependientes Vigo II. Vigo, Pontevedra. Spain. ³Residencia de Personas Mayores Fundación Caja de Burgos. Burgos, Spain. ⁴Residencia Jesán. Salamanca, Spain. ⁵Subdirección de Gestión y Servicios Generales. Complejo Asistencial Universitario de León. León, Spain. ⁶Unidad de Gestión Clínica de Endocrinología y Nutrición. Hospital Universitario Regional de Málaga. Málaga, Spain. ⁷Dietética y Nutrición. Grupo AUSOLAN. Mondragón, Guipúzcoa. Spain. ⁸Outcomes'10 S.L. Castellón de la Plana, Spain. ⁹Departamento Médico. Nestlé Health Science. Barcelona, Spain

Abstract

Introduction: although nutritional differences between different types of texture-modified diet (TMD) have been evaluated, the resources and costs associated with their preparation have been less studied.

Objective: to describe the nutritional, microbiological properties and costs of: 1) in-home produced pureed food (hTMD); 2) concentrated nutrient-dense commercial food products, hand-blended (cTMD); 3) food prepared using the MixxPro[®] automatic food mixer (cTMD-Mix).

Methods: an observational, prospective study carried out in three geriatric nursing-homes. Patients ≥ 65 years, receiving TMD, with a stable clinical condition, estimated survival/expected interment > 1 month, and sufficient cognitive capacity were included. The following data were recorded: 1) patient socio-demographic and clinical variables; 2) TMD compliance and symptoms related to dysphagia during the meal; 3) patient appetite; and 4) kitchen information and resources used to prepare a TMD.

Results: sixty-two residents were included (65.0 % women, 88.3 years (SD: 9.3); 43.5 % malnourished, 79.0 % with good appetite). The proportion of food eaten/median kcal served/portion/mean kcal consumed were: hTMD: 95.5 % (SD: 10.7)/92.4 kcal (IQR: 75.6-128.1)/88.2 kcal (IQR: 72.2-122.3); cTMD: 89.2 % (SD: 15.9)/323.4 kcal (IQR: 284.2-454.1)/288.5 kcal (IQR: 253.5-325.1); and cTMD-Mix: 80.3 % (SD: 21.4)/358.0 kcal (IQR: 344.0-372.1)/287.5 kcal (IQR: 276.5-298.8). No microorganisms were detected. The average time spent in preparing each portion and its costs were: hTMD: 11.2 min (SD: 3.89)/€2.33 (SD: 0.63); cTMD: 1.7 min (SD: 0.28)/€2.01 (SD: 0.39); and cTMD-Mix: 1.6 min (SD: 0.00)/€2.00 (SD: 0.33).

Conclusions: in patients with dysphagia and/or chewing difficulties, concentrated nutrient-dense food products, particularly those produced using the MixxPro[®] automatic food mixer, ensure a high caloric intake and allow quick and safe food preparation.

Keywords:

Nursing home.
Dysphagia.
Swallowing problems.
Malnutrition. Texture-modified diets.
Pureed.

Received: 02/12/2020 • Accepted: 17/02/2021

Funding: this work was supported by Nestlé Health Science.

Acknowledgments: the authors would like to thank all participants and professionals (nurses, physicians, kitchen employees, and medical managers) at the three geriatric nursing homes (Residencia Jesán, Residencia de Mayores Fundación Caja de Burgos, and Centro Residencial de Atención a Personas Dependientes Vigo I) involved in the study.

Conflicts of interest: this research was supported by Nestlé Health Science. NB is an employee of Nestlé Health Science. MC and HDP work for an independent research entity that received Nestlé Health Science funding to coordinate and conduct the study, and write up the manuscript. MDBP has received fees as a speaker at teaching and training activities for Nestlé Healthcare, Fresenius Kabi, Nutricia, Vegeat Healthcare, Persan Farma, Lilly, Novo Nordisk, and Adventia Farma. AMC has nothing to disclose. All authors declare their independence from the sponsor regarding the analysis of results and compilation of the conclusions.

Ballesteros-Pomar MD, Pérez-Martín J, Mendiola MJ, García-García JM, Parrado-Cuesta S, Caracuel AM, Garcés B, Comellas M, de Paz HD, Barcons N. Cost, microbiological, and nutritional properties of pureed food production in nursing homes. The ABADIA Study. *Nutr Hosp* 2021;38(3):470-477

DOI: <http://dx.doi.org/10.20960/nh.03465>

Correspondence:

María D. Ballesteros-Pomar. Unidad de Nutrición Clínica y Dietética. Sección de Endocrinología y Nutrición. Complejo Asistencial Universitario de León. Altos de Nava s/n. 24008 León, Spain
e-mail: mballesteros@telefonica.net

Resumen

Introducción: aunque existe evidencia acerca de las diferencias nutricionales entre los distintos tipos de dieta de textura modificada (DTM), los recursos y los costos asociados a su preparación se han estudiado menos.

Objetivo: describir las propiedades nutricionales, las microbiológicas y los costes de: 1) una dieta triturada de manera artesanal (hDTM); 2) una dieta preparada con alimentación básica adaptada (ABA) (cDTM); y 3) una ABA preparada con el mezclador automático de alimentos MixxPro® (cDTM-Mix).

Métodos: estudio observacional prospectivo realizado en tres residencias. Se incluyeron pacientes \geq 65 años que recibían DTM, con estado clínico estable, con supervivencia/internamiento estimado $>$ 1 mes y capacidad cognitiva suficiente. Se registraron: 1) las variables sociodemográficas y clínicas del paciente; 2) el cumplimiento y los síntomas relacionados con la disfagia durante la comida; 3) el apetito del paciente, y 4) la información de la cocina y los recursos utilizados para preparar la DTM.

Resultados: se incluyeron 62 residentes (65,0 % mujeres, 88,3 años (SD: 9,3), 43,5 % desnutridos, 79,0 % con buen apetito). La proporción de alimentos consumidos/mediana de kcal servidas/porción/mediana de kcal media consumidas fueron: hDTM 95,5 % (SD: 10,7)/92,4 kcal (IQR: 75,6-128,1)/88,2 kcal (IQR: 72,2-122,3); cDTM: 89,2 % (SD: 15,9)/323,4 kcal (IQR: 284,2-454,1)/288,5 kcal (IQR: 253,5-325,1), y cDTM-Mix: 80,3 % (SD: 21,4)/358,0 kcal (IQR: 344,0-372,1)/287,5 kcal (IQR: 276,5-298,8). No se detectaron microorganismos. El tiempo medio empleado en la preparación y el coste por porción fueron: hDTM: 11,2 min (SD: 3,89)/2,33 € (SD: 0,63); cDTM: 1,7 min (SD: 0,28)/2,01 € (SD: 0,39), y cDTM-Mix: 1,6 min (SD: 0,00)/2,00 € (SD: 0,33).

Conclusiones: en los pacientes con disfagia y/o dificultades para masticar, los productos de ABA comerciales, en particular los que se producen con el mezclador automático de alimentos MixxPro®, aseguran una elevada ingesta calórica y permiten una preparación rápida y segura.

Palabras clave:

Residencias. Disfagia. Problemas de deglución. Dietas de textura modificada. Purés.

INTRODUCTION

Swallowing difficulties are common in the elderly, especially among institutionalized people. Half of institutionalized individuals are estimated to suffer from oropharyngeal dysphagia (OD) (1,2). Malnutrition is one of the main consequences associated with OD. Although prevalence rates vary depending on the method used, over half of the institutionalized elderly are malnourished or at risk of suffering malnourishment (3,4). Besides, approximately 80 % of pneumonia cases in elderly people with swallowing disorders correspond to cases of aspiration pneumonia, one of the leading causes of death in this cohort (5).

Between 26 % and 67 % of the meals served in geriatric nursing homes are texture-modified diets (TMD) (6,7). The in-home production of TMD (hTMD) requires blending the food and then diluting it with water or broth, thus reducing its nutritional intake (8-11), and increasing variability between prepared meals (11). Up to 83 % of patients receiving hTMD do not meet the necessary nutritional requirements, leading to nutritional deficiency (12,13). Recently, the European Society for Clinical Nutrition and Metabolism (ESPEN) recommends providing enriched TMD to support adequate dietary intake in elderly with malnutrition or at risk of malnutrition and with signs of OD (14). Concentrated nutrient-dense commercial food products (cTMD) for adults with OD provide a suitable alternative to hTMD. Caloric and protein intake is significantly increased by cTMD, which improves the nutritional status of the patient (12,15,16).

It should be borne in mind that the high amount of handling required to prepare hTMD meals increases the risk of food poisoning (17-19), and is also labor-intensive, requiring significant investment in resources and time spent by the kitchen staff.

There are currently automatic food mixers that can automatically prepare cTMD (cTMD-Mix), with the potential to significantly increase time and resource savings, and to minimize microbiological risks.

Although there are studies that describe the nutritional differences between hTMD and cTMD, as far as we know this is the

first study to evaluate the resources and costs associated with TMD preparation. This observational study aims to describe the nutritional, microbiological properties, and the cost of three types of TMD: hTMD, cTMD, and cTMD-Mix.

METHODOLOGY

STUDY DESIGN

A proof-of-concept study with a descriptive, observational design was carried out in three geriatric nursing homes. The nursing homes were selected based on: 1) availability of own kitchen (not an external catering service); 2) cook with experience in the preparation of TMD \geq 6 months; 3) experience of the specialist prescribing the diet \geq two years.

In each of the nursing homes TMD was prepared using a method according to its routine practice: 1) in-home TMD (hTMD); 2) concentrated nutrient-dense commercial food products (cTMD); 3) cTMD-Mix prepared using the MixxPro® automatic food mixer to produce blended meals (cTMD-Mix).

The study was led by a committee of four experts (an endocrinologist, a catering manager, a bromatologist, and a food technologist) who participated in preparing the Case Report Form (CRF) and in the correct interpretation of study results.

PATIENTS

The study included patients \geq 65 years of age, who were receiving TMD at the start of the study, with a stable clinical condition and an estimated survival $>$ 1 month, with an expected internment \geq 30 days, with sufficient cognitive capacity to complete the questionnaires (as assessed by the specialist prescribing the diet) either alone or with the help of a caregiver, and who were not participating in any other clinical trial. The study excluded those patients who required a change in diet (oral, enteral, or parenteral) for more than two days.

The participants were identified by the healthcare professional who prescribed the TMD following a consecutive, non-random recruitment. Patients were followed up for 15 days.

The following tests were conducted for each participant: nutritional screening using the Mini Nutritional Assessment Short Form (MNA[®]-SF) (at baseline), risk of dysphagia assessment using the Eating Assessment Tool (EAT-10) (at baseline), and appetite determination using the Short Nutrition Assessment Questionnaire (SNAQ) (before the main course). Data of symptoms associated with swallowing during feeding (choking, fractional swallowing, throat clearing, coughing during and after ingestion) were also collected at each meal.

PREPARATION OF TEXTURE-MODIFIED DIETS (TMD)

For all three TMD types, the preparation methods of the meal's main course were evaluated for five different varieties: chicken, veal, fish, egg, or lentils.

The hTMD was prepared using traditional processing methods and further blending. The cTMD was based on Meritene[®] (Nestlé Health Science) commercial products in the instant puree range (chicken with rice and carrots, veal with vegetables, fish with rice, puréed eggs Provenzal style, and lentils with vegetables). This range of dehydrated products was reconstituted with water, either manually (cTMD) or mechanically using a MixxPro[®] automatic food mixer (cTMD-Mix).

STUDY VARIABLES

Four independent CRFs were designed to collect the study variables: 1) to be filled in by the specialist prescribing the diet, recording the patient's main socio-demographic and clinical variables; 2) to be filled in by nurses or healthcare assistants, collecting data on compliance and symptoms related to dysphagia occurring during the meal; 3) to be filled in by the patient, including data on appetite; and 4) to be filled in by the catering manager, including information on the kitchen and the resources used to prepare each diet variety (Table I).

The staff responsible for data collection received appropriate training in person. Additionally, if the staff required it, online training was available during the study.

An external laboratory conducted the microbiological and nutritional (estimation of kcal per portion) analysis.

STATISTICAL ANALYSIS

Data were analyzed with the STATA statistical package, version 14. The absolute and relative frequencies of qualitative variables were calculated. The mean, standard deviation (SD), and interquartile ranges were calculated for the quantitative variables.

The compliance of each patient was calculated based on: 1) the amount of TMD served during the 15 days: mean amount served

(according to the five cooking records) x 15 days; and 2) the average compliance of the patient during the 15 days of follow-up data recorded in the nursing questionnaire.

The average calorie count per portion was obtained based on the average number of calories per portion, and the average amount served per portion. The mean number of calories consumed per portion was calculated from the mean calories served per portion, and the mean compliance.

The costs related to food preparation were calculated on the following basis: 1) the cost of each of the ingredients used (€/kg, €/volume, or €/unit) according to the reference prices on the wholesale markets (20-23) or to the prices of origin; 2) the cost of human resources required for each of the processes — pre-processing, processing, blending (if applicable), homogenization (if applicable), and cleaning) — based on the time spent on each process, number of employees and wages (according to professional category (24,25) and updated to €/2019 in line with the CPI (26)); and 3) the number of portions prepared.

The time spent cooking, baking, frying, grilling, and blending/homogenizing was considered to calculate energy costs. For each process, the cost/hour was applied based on electricity or gas energy consumption, estimating the average cost of the corresponding energy supply at the time of the data analysis (27,28). Given the variability extant in the energy used by different types of blenders, a conservative 200 W value was applied. For instance, for the MixxPro[®] automatic food mixer, a maximum consumption of 2000 W was assumed, and a working time of 1 min per portion. Gas consumption was calculated by means of a direct extrapolation of the minutes recorded and energy consumption. The same energy tariffs were applied to all three centers, and similar appliances were assumed to have the same energy consumption.

The amount served per portion in each center was different, so the standardized cost per 100 g was calculated as follows: $\text{cost}/100 \text{ g} = 100 \text{ g} \times \text{cost per portion} [\text{€/portion} / \text{amount served per portion (g)}]$. Based on this result, the average cost per 100 g was obtained for each menu.

The average cost per 100 kcal portion was obtained from the average calories served per portion, and the cost per portion.

ETHICAL CONCERNS

The study was approved by the Clinical Research Ethics Committee at the *Complejo Asistencial Universitario de León* (E. OBS1676), and was conducted according to the principles of the Helsinki Declaration and good clinical practice (GCP). Patients were informed about the study and signed an informed consent. The data were duly anonymized.

RESULTS

PATIENT CHARACTERISTICS

Of the total number of patients starting the study (n = 64), two were excluded: one did not adjust to the diet, and one died.

Table I. Variables collected during the study; source and time of collection

Variable	Source	(No.) time of collection
Questionnaire completed by the prescriber		
Patient age	Medical record	(1) Baseline
Sex	Medical record	(1) Baseline
Body mass index (BMI)	Medical record	(1) Baseline
Diagnosis for which TMD is prescribed	Medical record	(1) Baseline
Chronic comorbidities	Medical record	(1) Baseline
Nutritional supplementation requirements	Medical record	(1) Baseline
Swallowing/gastrointestinal complications	Medical record	(1) Baseline
<i>Risk of dysphagia</i> [Risk of dysphagia ≥ 3]	EAT-10	(1) Baseline
<i>Malnutrition</i> [Malnutrition, < 7; risk of malnutrition, 8-11; good nutritional status, ≥ 12]	MNA® -SF	(1) Baseline
Questionnaire completed by nurses		
<i>Symptoms related to dysphagia</i> during eating (main course) [coughing during and after ingestion, throat-clearing, fractional swallowing and choking]	Reported by nursing staff	(15) Days 1 to 15
<i>Compliance</i> with the main course of the meal [0 %, 25 %, 50 %, 75 % or 100 %].	Reported by nursing staff	(15) Days 1 to 15
Questionnaire completed by the patient		
<i>Appetite</i> [good appetite, ≥ 14 points]	SNAQ	(3) Days 1, 7 and 15
Questionnaire completed by the cook		
<i>Kitchen features</i> [number of employees, professional category, number of daily regular meals and number of daily blended meals]	Catering staff	(1) Baseline
<i>Data for preparation of each meal variety</i> [number of meals prepared and ration weight, raw materials used (weight or volume or units of all ingredients required), human resources required (number of employees involved in the process*, professional category and time required). Energy consumption (time spent using kettles/pots, ovens, grills, fryers, pans, mixers, etc.)]	Catering staff	(5) During the preparation of each variety (chicken, veal, fish, egg, or lentils)
<i>Temperature</i> of the prepared meal [temperature after homogenisation and before serving]	Catering staff	(15) Three independent records per diet variety
<i>Microbiological analysis</i> [presence of coagulase-positive <i>Staphylococcus</i> , <i>Salmonella</i> spp, <i>Escherichia coli</i> β-glucuronidase+, and <i>Listeria monocytogenes</i>] and <i>energy content</i> (kcal/100 g)]	Accredited external laboratory (UNE-EN ISO/IEC 17025:2005)	(15) Three independent records per diet variety

*The time required for cleaning was estimated based on the time required to clean each utensil (by hand or machine) and the number of utensils used during the preparation of each diet variety.

Finally, 62 patients, 65 % women, with a mean age of 88.3 (SD: 9.3) years completed the follow-up and were included in the analysis (hTMD, n = 20; cTMD, n = 20; and cTMD-Mix, n = 22). Only 3.2 % of participants were well nourished (MNA®-SF ≥ 12). Baseline patient characteristics are shown in table II.

A total of 79.0 % of patients had a good appetite (SNAQ ≥ 14 points) during follow-up [hTMD: 100.0 %; cTMD: 75.0 %; cTMD-Mix: 63.6 %].

Overall, compliance with the diet was good. On average, the proportion of food eaten was above 80 %: hTMD: 95.5 % (SD: 10.7); cTMD: 89.2 % (SD: 15.9); cTMD-Mix: 80.3 % (SD: 21.4). Regarding compliance and the mean amount served per portion, patients receiving hTMD, cTMD, and cTMD-Mix ingested a daily mean of 197.7 g (SD: 21.6), 284.6 g (SD: 49.6), and 281.9 g (SD: 73.4), respectively. Similarly, considering the calories served, the calories consumed per portion were 88.2 (IQR: 72.2-122.3)

Table II. Baseline characteristics of study patients

	Total (n = 62)	hTMD (n = 20)	cTMD (n = 20)	cTMD-Mix (n = 22)
Age, years [mean (SD)]	88.3 (9.3)	89.9 (6.5)	94.2 (6)	81.6 (9.9)
Women [% (n)]	65.0 (39)	80.0 (16)	47.4 (9)	66.7 (14)
BMI, kg/m ² [mean (SD)]	23.9 (4.3)	22.7 (2.9)	22.8 (3.4)	25.9 (5.5)
<i>MNA</i> [®] , [% (n)]				
Malnourished	43.5 (27)	60.0 (12)	50.0 (10)	22.7 (5)
At risk of malnutrition	53.2 (33)	40.0 (8)	50.0 (10)	68.2 (15)
Normal nutritional status	3.2 (2)	0.0 (0)	0.0 (0)	9.1 (2)
Risk of dysphagia (EAT-10 \geq 3), [% (n)]	91.9 (57)	100.0 (20)	75.0 (15)	100.0 (22)
With nutritional supplements, [% (n)]	30.6 (19)	20.0 (4)	40.0 (8)	31.8 (7)
<i>Number of comorbidities</i> , [% (n)]				
0	1.6 (1)	0.0 (0)	0.0 (0)	4.5 (1)
1	32.3 (20)	85.0 (17)	10.0 (2)	4.5 (1)
\geq 2	66.1 (41)	15.0 (3)	90.0 (18)	90.9 (20)
<i>Previous swallowing and intestinal complications</i> , [% (n)]				
Aspiration pneumonia	33.9 (21)	80.0 (16)	25.0 (5)	0.0 (0)
Constipation	19.4 (12)	0.0 (0)	0.0 (0)	54.5 (12)
Nausea or vomiting	9.7 (6)	0.0 (0)	30.0 (6)	0.0 (0)
Diarrhea	3.2 (2)	0.0 (0)	10.0 (2)	0.0 (0)
Abdominal pain	1.6 (1)	0.0 (0)	0.0 (0)	4.5 (1)

hTMD: in-home TMD; cTMD: commercial TMD, hand blended; cTMD-Mix: commercial TMD, automatically blended; SD: standard deviation; BMI: body mass index; MNA: mini nutritional assessment; EAT-10: eating assessment tool.


for hTMD; 288.5 (IQR: 253.5-325.1) for cTMD; and 287.5 (IQR: 276.2-298.8) for cTMD-Mix.

The mean number of symptoms associated with swallowing during feeding (choking, fractional swallowing, throat clearing, coughing during and after ingestion) was 68 (hTMD), 46 (cTMD), and 10 (cTMD-Mix) for every 100 meals served, respectively.

CHARACTERISTICS OF PROCESSED MEALS

The average number of TMD served per portion, as well as the energy intake, differed in the three nursing homes: 207.0 g (SD: 9.0) and 92.4 kcal (IQR: 75.6-128.1) for hTMD; 319.2 g (SD: 35.4) and 323.4 kcal (IQR: 284.2-375.3) for cTMD; and 351.0 g (SD: 0.0) and 358.0 kcal (IQR: 344.0-372.1) for cTMD-Mix (Fig. 1). Greater variability was observed for those meals requiring higher levels of food handling for their processing: 56.1 % (hTMD), 26.1 % (cTMD), and 7.8 % (cTMD-Mix).

The mean temperature of meals after blending was 80.6 °C (SD: 1.6) for hTMD, 88.2 °C (SD: 1.02) for cTMD, and 85.0 °C (SD: 0.0) for cTMD-Mix; while at the time of serving to patients, it was 54.3 °C (SD: 1.22) for hTMD, 58.7 °C (SD: 0.31) for hTMD, and 75.0 °C (SD: 0.0) for cTMD-Mix.

**Figure 1.**

Kilocalories served per ration: interquartile ranges (hTMD: in-home TMD; cTMD: commercial TMD, hand blended; cTMD-Mix: commercial TMD, automatically blended).

No microorganisms were detected in any of the analyzed samples.

USE OF RESOURCES AND COSTS ASSOCIATED WITH FOOD PREPARATION

The number and type of kitchen employees involved in food preparation varied between nursing homes (Table III).

In each kitchen, 69 % (hTMD), 81.8 % (cTMD), and 53.8 % (cTMD-Mix) of the total number of prepared meals (including non-blended types) were TMD. The mean number of portions prepared during the study period was 28.8 (SD: 2.9) for hTMD, 26.0 (SD: 1.4) for cTMD, and 70.0 (SD: 0.0) for cTMD-Mix. The average

time spent in preparing each portion was 11.2 min (SD: 3.89) for hTMD, 1.7 min (SD: 0.28) for cTMD, and 1.6 min (SD: 0.00) for cTMD-Mix (Table III).

The total cost per portion (human resources, ingredients, and energy consumption) amounted to €2.33 (SD: 0.63), €2.01 (SD: 0.39), and €2.00 (SD: 0.33) for hTMD, cTMD, and cTMD-Mix, respectively. In the nursing home employing hTMD the main cost component was associated with human resources, whereas in the centers using cTMD and cTMD-Mix the main cost was related to raw materials (Fig. 2).

Table III. Use of resources and time spent in meal preparation

Type of diet	No. of portions prepared	Process	Resource	Total time (SD) min	Time/portion (SD) min
hTMD	28.8	Pre-processing	1 cook	77.0 (45.3)	2.7 (1.7)
		Processing	1 cook	112.0 (24.0)	3.9 (0.9)
		Blending/homogenization	1 cook	12.8 (1.9)	0.4 (0.1)
		Serving up	1-2 cooks	8.8 (2.8)	0.3 (0.1)
		Cleaning	1 cook	110.8 (24.3)	4.0 (1.0)
		Total	-	321.4 (96.1)	11.2 (3.9)
cTMD	26.0	Pre-processing	1 assistant	4.4 (5.4)	0.2 (0.2)
		Processing	1 assistant	15.0 (0.7)	0.6 (0.0)
		Blending/homogenization	1 assistant	5.2 (0.4)	0.2 (0.0)
		Serving up	2 healthcare assistants	10.4 (0.9)	0.4 (0.0)
		Cleaning	1 catering assistant	7.00 (0.0)	0.3 (0.0)
		Total	-	42.0 (6.2)	1.7 (0.3)
cTMD-Mix	70.0	Pre-processing	1 cook	15.0 (0.0)	0.2 (0.0)
		Processing	2 cooks 1 assistant	90.0 (0.0)	1.3 (0.0)
		Blending/homogenizing	-	-	-
		Serving up	-	-	-
		Cleaning	1 catering assistant	10.0 (0.0)	0.1 (0.0)
		Total	-	115 (0.0)	1.6 (0.0)

hTMD: in-home TMD; cTMD: commercial TMD, hand blended; cTMD-Mix: commercial TMD, automatically blended; SD: standard deviation; Min: minutes.




Figure 2.

Breakdown of cost per ration (hTMD: in-home TMD; cTMD: commercial TMD, hand blended; cTMD-Mix: commercial TMD, automatically blended).

The average cost per 100 g was €1.12 (SD: 0.29) for hTMD, €0.65 (SD: 0.18) for cTMD, and €0.57 (SD: 0.09) for cTMD-Mix. Finally, the average cost per 100 kcal was €2.80 (SD: 1.32) for hTMD, €0.62 (SD: 0.18) for cTMD, and €0.58 (SD: 0.13) for cTMD-Mix.

DISCUSSION

TMD is recommended for patients who have difficulty swallowing. To facilitate its intake, the texture of the original food is modified by blending, and very often water or broth is added, which can lead to nutritional deficiency in the prepared meal and greater variability between portions (15). In line with previous studies (11), our work has demonstrated a lower energy intake and a higher variability associated with hTMD meals.

The need to blend food to adapt it to a patient's swallowing requirements can also modify its organoleptic qualities and be associated with suboptimal intake (8,29). In our study, the percentage of food eaten, as compared to food served, was high (> 80 %), and it was higher in patients receiving hTMD than in those receiving cTMD or cTMD-Mix. However, this difference may be accounted for because the portions served in the nursing homes serving hTMD (approximately 200 g) were smaller than those served in the centers offering commercial diets (approximately 350 g). In fact, the total amount of ingested food per portion was about 30 % higher in the homes serving commercial meals. This fact, together with the lower caloric intake of hTMD, could diminish the feeling of fullness, and explain the greater appetite observed in patients with hTMD. Similarly to our study, Rubio et al. compared the intake of an hTMD with a hyperproteic and hypercaloric powdered preparation (cTMD), observing that the total amount ingested by the patient was lower in the cTMD group, with a higher energy and protein intake (30). Other studies have described that in both institutionalized elderly patients and elderly outpatients with swallowing difficulties, a higher energy and protein intake was achieved with cTMD as compared with hTMD (31,32). Similarly, our results show that, although compliance was higher in the nursing home with hTMD, the final kilocalories ingested were much higher in the patients receiving cTMD or cTMD-Mix than in the patients receiving hTMD.

Another reason for prescribing TMD is to reduce the risk of choking and prevent aspiration pneumonia. Up to 70 % of institutionalized people may suffer from choking during meals (33). Evidence of symptoms related to swallowing (such as choking) associated with hTMD vs. cTMD is minimal; moreover, results on the effectiveness of TMD to prevent aspiration pneumonia are inconclusive (34,35). In the present study, the number of swallowing-related symptoms observed was higher in patients receiving hTMD, decreasing with the diets requiring lower levels of manipulation during preparation, as was the case of cTMD-Mix. These results would be consistent with the fact that patients with hTMD reported more aspiration pneumonia (before study initiation) than patients receiving cTMD and cTMD-Mix. How-

ever, due to the study's nature, these results are neither representative nor conclusive enough to establish a cause-effect relationship.

During diet elaboration, food needs to reach a temperature above 65 °C, and should not be exposed to temperatures between 15 °C and 45 °C to guarantee microbiological safety (12). The three types of diet studied showed a good safety profile. In all cases, microbiological safety temperatures (> 65 °C) were reached during preparation, and the presence of microorganisms was not detected in the analysis. It should be noted that only cTMD-Mix remained above 65 °C until the time of serving.

As far as we know, this is the first study conducted in geriatric nursing homes to evaluate the use of resources and costs of TMD production for patients with dysphagia or mastication difficulties using natural and commercial foods, and three different processing methods (hTMD, cTMD, cTMD-Mix).

Raw material costs were lower for hTMD when compared to cTMD and cTMD-Mix. However, the time required to prepare each portion was much lower in the latter. Approximately, 1.5 minutes for commercial diets and 11 minutes for hTMD were required. Considering the resources needed to prepare the diets, the cost per portion and the cost per 100 g served were lower for the commercial diets than for the home-made ones.

Previous studies suggest that patients who switch from hTMD to fortified TMD (13,31) or cTMD (32) experience significant weight gain improvements. Additionally, two observational studies showed that cTMD-Mix was a simple and effective way to improve nutritional status in elderly institutionalized residents with swallowing disorders (36,37). The scope of our study was not to assess the nutritional status of patients. Considering that most of our participants were already receiving TMD before the study started, our results appear to be in line with the above studies. In this respect, the percentage of patients with malnutrition was higher in the nursing home serving hTMD and lower in the nursing home serving hTMD-Mix.

Our study presents some limitations. The first one relates to the study population, as patient recruitment was made according to convenience, and the number of participants was relatively low, preventing a robust statistical analysis. Despite the small sample size, since study participants include elderly residents with a high level of dependence, we can assume that the study population is representative of nursing home residents. Similarly, the study involved a reduced number of kitchens, each one with specific characteristics and peculiarities, so their practices may not be representative. However, it is important to keep in mind that the study objective was descriptive rather than comparative. Some participants might have required help to eat; however, since such data were not collected, it cannot be established whether having support influenced the amount of food consumed. Finally, since the nursing home staff collected the data, there may be inter-staff variability in measurements. In order to reduce this possible bias, the staff responsible for data collection received appropriate training, both in person and online; a graduated scale widely used in observational studies was used to assess the amount ingested; only objective symptoms related to dysphagia disorders

such as choking, fractional swallowing, throat clearing, coughing during and after ingestion were reported. Despite these limitations, the study provides useful information for future analyses of the potential benefits and costs of TMD as prepared using different processes.

In conclusion, in patients with dysphagia and/or chewing difficulties, concentrated, nutrient-dense food products (Meritene®, Nestlé Health Science), particularly those produced using the MixxPro® automatic food mixer, ensure a high caloric intake, even in subjects with lower compliance rates. Moreover, they allow quick and safe food preparation, reducing both kitchen workload and production costs. Further studies with a larger sample of patients and centers are needed to obtain representative results and establish whether there is any cause-effect of such diets on patient nutritional status.

REFERENCES

- Mioche L, Bourdiol P, Peyron M-A. Influence of age on mastication: effects on eating behaviour. *Nutr Res Rev* 2004;17:43-54. DOI: 10.1079/NRR200375
- Ortega O, Martín A, Clavé P. Diagnosis and Management of Oropharyngeal Dysphagia Among Older Persons, State of the Art. *J Am Med Dir Assoc* 2017;18(7):576-82. DOI: 10.1016/j.jamda.2017.02.015
- Vandewoude MFJ, van Wijngaarden JP, De Maesschalck L, Luijing YC, Van Gossum A. The prevalence and health burden of malnutrition in Belgian older people in the community or residing in nursing homes: results of the NutriAction II study. *Aging Clinical and Experimental Research* 2019;31(2):175-83. DOI: 10.1007/s40520-018-0957-2
- Bell CL, Lee ASW, Tamura BK. Malnutrition in the nursing home. *Curr Opin Clin Nutr Metab Care* 2015;18(1):17-23. DOI: 10.1097/MCO.000000000000130
- Teramoto S, Fukuchi Y, Sasaki H, Sato K, Sekizawa K, Matsuse T. High incidence of aspiration pneumonia in community- and hospital-acquired pneumonia in hospitalized patients: a multicenter, prospective study in Japan. *J Am Geriatr Soc* 2008;56(3):577-9. DOI: 10.1111/j.1532-5415.2008.01597.x
- Painter V, Le Couteur D, Waite L. Texture-modified food and fluids in dementia and residential aged care facilities. *Clin Interv Aging* 2017;12:1193-203. DOI: 10.2147/CIA.S140581
- Streicher M, Wirth R, Schindler K, Sieber CC, Hiesmayr M, Volkert D. Dysphagia in Nursing Homes—Results From the NutritionDay Project. *J Am Med Dir Assoc* 2018;19(2):141-7.e2. DOI: 10.1016/j.jamda.2017.08.015
- Wright L, Cotter D, Hickson M, Frost G. Comparison of energy and protein intakes of older people consuming a texture modified diet with a normal hospital diet. *J Hum Nutr Diet* 2005;18(3):213-9. DOI: 10.1111/j.1365-277X.2005.00605.x
- Massoulard A, Bonnabau H, Gindre-Poulvelarie L, Baptiste V, Preux PM, Villemonteix C, et al. analysis of the food consumption of 87 elderly nursing home residents, depending on food texture. *J Nutr Heal Aging* 2011;15(3):192-5. DOI: 10.1007/s12603-010-0271-y
- Nowson CA, Sherwin AJ, McPhee JG, Wark JD, Flicker L. Energy, protein, calcium, vitamin D and fibre intakes from meals in residential care establishments in Australia. *Asia Pac J Clin Nutr* 2003;12(2):172-7.
- Beck AM, Hansen KS. Meals served in Danish nursing homes and to meals-on-wheels clients may not offer nutritionally adequate choices. *J Nutr Elder* 2010;29(1):100-9. DOI: 10.1080/01639360903574742
- Velasco C, García-Peris P. Tecnología de alimentos y evolución en los alimentos de textura modificada; del triturado o el deshidratado a los productos actuales. *Nutr Hosp* 2014;29(3):465-9.
- Germain I, Dufresne T, Gray-Donald K. A Novel Dysphagia Diet Improves the Nutrient Intake of Institutionalized Elders. *J Am Diet Assoc* 2006;106(10):1614-23. DOI: 10.1016/j.jada.2006.07.008
- Volkert D, Beck AM, Cederholm T, Cruz-Jentoft A, Goisser S, Hooper L, et al. ESPEN guideline on clinical nutrition and hydration in geriatrics. *Clin Nutr* 2019;38(1):10-47. DOI: 10.1016/j.clnu.2018.05.024
- Rocamora JAI, García-Luna PP. El menú de textura modificada; valor nutricional, digestibilidad y aportación dentro del menú de hospitales y residencias de mayores. *Nutr Hosp* 2014;29(4):873-9.
- Caracuel García A. Alimentación adaptada en el adulto: propiedades nutricionales y funcionales de los alimentos comerciales de textura modificada. In: Andrés Carretero M, editor. *Actualización en bromatología hospitalaria*; 2009. p. 181-90.
- Rosas R. Contaminaciones alimentarias. *Offarm* 2007;26(6):95-100.
- Camps N, Comapry M, Follia N. Toxiinfección alimentaria producida por virus norwalk en una residencia geriátrica asistida. *Gac Sanit* 2001;15(2):31-130.
- Lund B, O'Brien S. Microbiological safety of food in hospitals and other healthcare settings. *J Hosp Infect* 2009;73(2):109-20. DOI: 10.1016/j.jhin.2009.05.017
- Precios de alimentos frescos y de temporada en mercados mayoristas de Mercasa. Available from: <https://www.mercasa.es/red-de-mercasa/precios-y-mercados-mayoristas>
- Estadísticas—mercadmadrid. Available at: <https://www.mercamadrid.es/estadisticas/>
- Agricultura, Medio Ambiente y Mundo Rural. *Boletín Agrario*; 2019. Available at: <https://boletinagrario.com/>
- Precios Medios. Available from: https://www.mapa.gob.es/app/precios-medios-nacionales/pmn_tabla.asp
- Ministerio de empleo y seguridad social. Resolución de 16 de febrero de 2018, de la Dirección General de Empleo, por la que se registran y publican las tablas salariales para los años 2017 y 2018 del Convenio colectivo estatal del sector laboral de restauración colectiva. *BOE* 2018;53:25459-80.
- Ministerio de empleo y seguridad social. *Boletín Oficial Del Estado* no 209; jueves 31 de agosto de 2017. Anexo I tablas salariales 2017. Sec. III. Pág. 86300.
- Actualización de rentas con el IPC general. Available at: <https://www.ine.es/calcula/>
- Tarifas de gas. 1er trimestre 2019. Preciogas; 2019. Available at: <https://preciogas.com/faq/factura-gas-natural>
- Tarifas de luz. 1er trimestre. Preciogas; 2019. Available at: <https://preciogas.com/faq/factura-luz>
- Colodny N. Dysphagic Independent Feeders' Justifications for Noncompliance With Recommendations by a Speech-Language Pathologist. *Am J Speech-Language Pathol* 2005;14(1):61-70. DOI: 10.1044/1058-0360(2005/008)
- Rubio M, Cabrerizo L, Galaz R. Estudio comparativo y aleatorio entre la dieta triturada habitual hospitalaria frente a una fórmula hipercalórica e hiperproteica (Vegenat-MED). *Nutr Hosp* 2002;17(S3):105.
- Ott A, Senger M, Lötzeyer T, Gefeller O, Sieber CC, Volkert D. Effects of a Texture-Modified, Enriched, and Reshaped Diet on Dietary Intake and Body Weight of Nursing Home Residents with Chewing and/or Swallowing Problems: An Enable Study. *J Nutr Gerontol Geriatr* 2019;38(4):361-76. DOI: 10.1080/21551197.2019.1628158
- De Luis DA, Izaola O, Prieto R, Mateos M, Aller R, Cabezas G, et al. Effect of texture modified diets in elderly ambulatory patients. *Nutr Hosp* 2009;24(1):87-92.
- Miles A, Dennison K, Oad MA, Shasha L, Royal M. Consumer Satisfaction of Texture Modified Meals Served in Residential Aged-Care Facilities. *Int J Food Sci Nutr Res* 2019;1(1). DOI: 10.31546/IJFSNR.1005
- Sukkar SG, Maggi N, Travalca Cupillo B, Ruggiero C. Optimizing Texture Modified Foods for Oro-pharyngeal Dysphagia: A Difficult but Possible Target? *Front Nutr* 2018;5:68. DOI: 10.3389/fnut.2018.00068
- O'Keefe ST. Use of modified diets to prevent aspiration in oropharyngeal dysphagia: is current practice justified? *BMC Geriatr* 2018;18(1):167. DOI: 10.1186/s12877-018-0839-7
- Khoo M, Lee S, Fong R, Tang L. Effect of a concentrated texture-modified diet on the nutritional status of elderly in a long-term care facility in Hong Kong. *Eur Geriatr Med [Internet]* 2019;10(September):S292. Available from: <http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L631722629%0Ahttp://dx.doi.org/10.1007/s41999-019-00221-0>. DOI: 10.1007/s41999-019-00221-0
- Ianes A, Giordani A, Ardoino G, D'Ostuni F. Impact of texture-modified diets based on concentrated commercial food products for dysphagia prepared with the food mixer MixxPro® on the body weight of institutionalized elderly patients. *Eur Geriatr Med [Internet]* 2019;10(September):S291. Available from: <http://www.embase.com/search/results?subaction=viewrecord&from=export&id=L631722449%0Ahttp://dx.doi.org/10.1007/s41999-019-00221-0>. DOI: 10.1007/s41999-019-00221-0