



Trabajo Original

Association between birthweight, cardiovascular risk factors, and depression in young Mexican adults

Asociación entre peso al nacer, riesgo cardiovascular y depresión en adultos jóvenes mexicanos

Luis Adolfo Santos-Calderón¹, Juan Manuel Vargas-Morales², Omar Sánchez-Armáss Capello³, Margarita Terán-García⁴, Patricia Elizabeth Cossío-Torres⁵, Marisol Vidal-Batres⁶, Joan D. Fernández-Ballart⁷, Celia Aradillas-García⁸, and UP AMIGOS Team

¹Facultad de Medicina. Universidad Autónoma de San Luis Potosí. San Luis Potosí, S.L.P. Mexico. ²Facultad de Ciencias Químicas. Universidad Autónoma de San Luis Potosí. San Luis Potosí, S.L.P. Mexico. ³Facultad de Psicología. Universidad Autónoma de San Luis Potosí. San Luis Potosí, S.L.P. Mexico. ⁴University of Illinois Urbana-Champaign. Urbana-Champaign, Illinois. USA. ⁵Departamento de Salud Pública. Facultad de Medicina. Universidad Autónoma de San Luis Potosí. San Luis Potosí, S.L.P. Mexico. ⁶Coordinación para la Innovación y Aplicación de la Ciencia y la Tecnología (CIACYT). Universidad Autónoma de San Luis Potosí. San Luis Potosí, S.L.P. Mexico. ⁷Medicina Preventiva y Salud Pública. Facultad de Medicina y Ciencias de la Salud. Universidad Rovira i Virgili. Tarragona, Spain. ⁸Centro de Investigación Aplicada en Ambiente y Salud (CIAAS). Facultad de Medicina/CIACYT. Universidad Autónoma de San Luis Potosí. San Luis Potosí, S.L.P. Mexico

Abstracts

Background: the aim of this study was to investigate the association between birthweight, cardiovascular disease (CVD) risk factors, and depression in young Mexican adults.

Methods: birthweight reports, family history of CVD and diabetes-related diseases, anthropometrics, serum lipid profile (total cholesterol [TC], triglycerides [TG], high-density lipoprotein-cholesterol [HDL-C], low-density lipoprotein-cholesterol [LDL-C], and very-low density lipoprotein-cholesterol [VLDL-C]), and depressive symptoms were measured in 778 subjects of the UP-AMIGOS cohort study. To investigate the association between birthweight categories and CVD risk factors and depression, a one-way analysis of variance with post-hoc test was performed of quantitative variables, and χ^2 test for qualitative variables.

Results: mean age was 17.8 years and 469 (60.3 %) of patients were female (n = 469, 60.3 %). The percentage of patients with low birthweight (LBW) was 8.1 % (n = 63), and 3.3 % (n = 26) reported high birthweight (HBW). Young adults with HBW were associated with elevated diastolic blood pressure (DBP), and high weight and body mass index (BMI) when compared to LBW subjects, the difference being statically significant (p < 0.05). Birthweight had no significant association with depression (p > 0.67).

Conclusion: the findings from this population-based study revealed a positive relation between birthweight categories and some CVD risk factors. Depression was not related to birthweight.

Keywords:

Birthweight.
Cardiovascular risk.
Depression. Mexican population.

Received: 28/01/2021 • Accepted: 20/05/2021

Author contributions: conceptualization: S.L.A. and A.C.; formal analysis: S.L.A. and C.P.E.; writing-original draft preparation: S.L.A.; writing-review editing: S.L.A., S.O., C.P.E., V.M, F.J.D. and V.J.M.; supervision: A.C. and C.P.E.; investigation: V.J.M., S.O. and T.M.

Funding: this research was funded by grants from the UASLP Hormones Laboratory at the School of Medicine, Clinical Biochemistry Laboratory, Chemical Sciences School, and the University Health Center under agreement support C09-PIFI-030606 (to C. Aradillas-García); the University of Illinois at Urbana-Champaign Research Board (#09070) and Center on Health, Aging, and Disability (to F. Andrade); and the USDA National Institute of Food and Agriculture, Hatch Projects #600108-793000-793323 (to M. Raffaelli) and #600109-698000-698354 (to M. Terán-García).

Acknowledgments: the data presented here represent a subproject of the UP AMIGOS study group (universities of San Luis Potosí and Illinois: A multidisciplinary Investigation on Genetics, Obesity and Social environment). We are grateful to all the research staff and study participants. The authors also thank MCF. Gicela de Jesús Galván Almazán for her contribution and assistance in the management of databases.

Conflicts of interest: the authors declare no conflicts of interest and that the funders had no role in the design of the study; in the collection, analyses, or interpretation of the data; in the writing of the manuscript; or in the decision to publish the results.

Santos-Calderón LA, Vargas-Morales JM, Sánchez-Armáss Capello O, Terán-García M, Cossío-Torres PE, Vidal-Batres M, Fernández-Ballart JD, Aradillas-García C, and UP AMIGOS Team. Association between birthweight, cardiovascular risk factors, and depression in young Mexican adults. *Nutr Hosp* 2021;38(4):833-838

DOI: <http://dx.doi.org/10.20960/nh.03547>

Correspondence:

Celia Aradillas-García. Centro de Investigación Aplicada en Ambiente y Salud (CIAAS). Facultad de Medicina/CIACYT. Universidad Autónoma de San Luis Potosí. Av. Sierra Leona nº 550-2ª. 78210 San Luis Potosí, S.L.P. Mexico
e-mail: celia@uaslp.com

Resumen

Introducción: el objetivo de este estudio fue investigar la asociación entre el peso al nacer, los factores de riesgo de las enfermedades cardiovasculares (ECV) y la depresión en adultos mexicanos jóvenes.

Métodos: se obtuvieron como variables el peso al nacer, los antecedentes heredofamiliares de ECV y diabetes, la antropometría, el perfil lipídico (colesterol total [CT], triglicéridos [TG], lipoproteína de alta densidad [C-HDL], lipoproteína de baja densidad [C-LDL] y lipoproteína de muy baja densidad [C-VLDL]) y los síntomas de depresión de 778 participantes del estudio de cohortes UP-AMIGOS. Se realizó un análisis de la varianza de 1 vía con pruebas *post hoc* para investigar la asociación entre las categorías de peso al nacer, riesgo ECV y depresión entre las variables cuantitativas; para las variables cualitativas se realizaron pruebas del χ^2 .

Resultados: la edad media fue de 17,8 años y 469 (60,3 %) de los participantes eran mujeres. El porcentaje de pacientes con bajo peso al nacer (BPN) fue del 8,1 % (n = 63) y el 3,3 % (n = 26) tenían elevado peso al nacer (EPN). Se encontró una asociación en el grupo de EPN con la elevación de la presión arterial diastólica (PAS), el peso y el índice de masa corporal (IMC) al compararlo con el grupo de BPN, con una diferencia significativa de $p < 0,05$. No se encontró ninguna asociación entre el peso al nacer y la depresión ($p > 0,67$).

Conclusiones: se encontró una relación positiva entre las categorías de peso al nacer con algunos factores de riesgo de ECV. La depresión no se asoció con el peso al nacer según los resultados de este estudio poblacional.

Palabras clave:

Peso al nacer. Riesgo cardiovascular.
Depresión. Población mexicana.

INTRODUCTION

The “developmental origins of health and disease” hypothesis explains that an adverse environment during a critical period of growth and development may have lifelong consequences on organ development and mental health status, and may increase the risk for certain non-communicable diseases (NCDs) in the mid and long term (1-3).

In 2015, 20.5 million newborns were registered worldwide, and 14.6 % of them had low birthweight (LBW < 2500 g). In that year in Mexico, it was estimated that 5-10 % of newborns had LBW (4). Furthermore, the incidence of high birthweight (HBW > 4000 g) varies across different regions of the world, depending on the cut-off value of minimum weight. Worldwide estimates range from 3 % to 10 % of all newborns, while in Mexico 4.8 % of newborns have HBW (7). Both LBW and HBW are risk factors for certain NCDs.

Currently, the association between birthweight, components of the metabolic syndrome (MetS) and mental illness remains controversial. Results are inconsistent, with some studies showing strong associations while others do not (6-20). Regarding birthweight and MetS, a recent cross-sectional study in Chinese and Spanish adolescents reported that HBW is associated with hyperglycemia, low high-density lipoprotein-cholesterol (HDL-C), hypertriglyceridemia, and abdominal obesity (7). HBW is associated with early development of insulin resistance, type-2 diabetes mellitus (T2DM), obesity, and MetS (6,7). Regarding high blood pressure, several studies have shown an inverse relationship between birthweight and blood pressure (BP) (8,9). LBW in combination with a positive family history for T2DM in the maternal branch is a risk factor for MetS, independently of weight status (6,10). Previous reports indicated a J-shape or U-shape relationship between birthweight and fat mass later in life (6).

A recent study indicates that LBW, used as a proxy for intrauterine growth retardation, is causally related to increased susceptibility to coronary artery disease and T2DM. The authors reported that this relationship was not mediated by adult obesity or hypertension (11). Also, childhood cardiometabolic risk exhibits specific patterns for race and gender. For example, birthweight and high waist circumference (WC) in black children follow a U-shaped relation-

ship, as well as high systolic blood pressure (SBP) in Mexican and Hispanic children. LBW with high SBP and low HDL-C appears to be more prominent in boys, while an inverse association with high HbA1c is more evident in girls (12).

All these findings differ from those presented by Dos Santos et al., who reported that birthweight is not a risk factor for the development of MetS during adolescence (13), and those presented by de Rooij et al., where LBW is not associated with an increased prevalence of MetS in adulthood (14).

Few studies have evaluated the potential association between birthweight and mental illness. There is evidence that LBW increases the risk for psychiatric problems such as stress, major depressive disorder (MDD), and anxiety disorders when compared with infants weighing > 3500 g at birth (15). Other studies show a U-shape relationship between birthweight, mental disorders, and insomnia. HBW and LBW adolescents have a higher risk for mental problems and sleep disorders (6,16). A recent study refuted the hypothesis from observational studies about the association between birthweight and mental disorders with Mendelian randomization — a novel method that tests causality on the basis of genetic data (17).

Some studies show a correlation between mental problems and development of cardiovascular diseases (CVD) later in life (18,19). A recent study with primary care patients found that depression and anxiety disorders were associated with an increased risk for arterial hypertension, obesity, and hyperlipidemia (20).

To our knowledge, there is no data about the relationship between birthweight, depression, and CVD risk factors among Mexican young adults; therefore, we decided to investigate the possible association between these three variables in young adult students.

MATERIALS AND METHODS

STUDY POPULATION

Participants were selected from the 2009 UP AMIGOS cross-sectional study (Universities of San Luis Potosí and Illinois:

A Multidisciplinary Investigation on Genetics, Obesity, and Social Environment). This was a cross-sectional study of young adults aged 17 years 6 months and one day to 25 years, applicants to a public university in Mexico, who were seeking admission for the 2010 school cycle between February and July of 2009.

As part of the university application process, all participants were submitted to a health screen at the university clinic. The health screen consisted of: 1) anthropometric measurements, including height, weight, WC, and BP; 2) a medical interview and physical exam conducted by a physician; 3) a blood draw following an overnight fast for biological markers; and 4) the self-reported depressive symptoms questionnaire Center for Epidemiological Studies Depression Scale (CESD-10) (21). Individuals willing to participate were apprised of study goals and their rights before signing the informed consent form. The Institutional Review Boards of both the Autonomous University of San Luis Potosí and the University of Illinois at Urbana-Champaign approved the study protocol.

Briefly, a total of 6,400 samples and valid questionnaires were collected. Only 778 were found to be eligible on the basis of the selection criteria with complete demographics, lipid profiles, and birthweight data. Birthweight was registered and categorized as < 2500 g, 2500 to 4000 g, and > 4000 g.

ANTHROPOMETRIC MEASURES

All measures were taken by trained health professionals at the university clinic. For all measurements, students were barefoot and wearing light clothes. Height was measured using a stadiometer to the nearest 0.5 cm, and weight was measured on a calibrated scale and recorded to the nearest 0.1 kg (Torino, Mexico). Weight and height were measured twice and the mean of the two values was reported. For the body mass index (BMI) calculation, weight (kg) was divided by the square height (m²) and classified in accordance with the World Health Organization's international classification: underweight (≤ 18.49 kg/m²), normal weight (18.50-24.99 kg/m²), overweight (25.00-29.99 kg/m²) and obese (≥ 30.00 kg/m²). WC was measured using a flexible, non-stretching nylon tape at the level between the last rib and iliac crest following normal expiration, with the subject in the standing position, and was recorded to the nearest 0.1 cm. WC risk level for MetS was determined according to the International Diabetes Federation (IDF)/American Heart Association (AHA) population- and sex-specific joint definition: at risk males (≥ 90 cm) and at risk females (≥ 80 cm) (22).

BLOOD PRESSURE

BP was measured by trained and certified healthcare providers according to a protocol adapted from the procedures recommended by the American Heart Association. BP was taken on the dominant arm after at least a 5-min rest while participants were in a sitting position. Arms were bare and supported at heart level,

and we used appropriately sized Welch Allyn cuffs. According to the harmonized definition, high blood pressure was defined as SBP ≥ 130 mmHg and/or DBP ≥ 85 mmHg (22).

FAMILY HISTORY

The family history of CVD and T2DM was assessed during the medical interview, including family members (sisters, brothers, parents, grandparents) known to have the disease.

ASSESSMENT OF DEPRESSIVE SYMPTOMS

Participants completed the CESD-10 questionnaire. Items evaluate depressed affect, positive affect, somatic complaints, and interpersonal problems. In a previous Mexican study, the CESD-10 tool showed good psychometric properties with Cronbach's $\alpha = 0.72$ (23). Participants rated how often they experienced each symptom during the past week using 4 response options from "Rarely or none of the time" (less than 1 day) to "All of the time" (5-7 days). Positive items were reversed, and responses assigned numeric values (0 to 3) before being summed. Possible scores range from 0 to 30, with a score of 10 or higher considered a positive screen for depression (21,24).

BIOCHEMICAL MEASUREMENTS

Blood samples were collected after an overnight fast (≥ 8 hours). All samples were measured in the Alcyon 300 autoanalyzer (Abbott, Illinois, USA, with reagents from BioSystems, Mexico). Fasting blood glucose (FBG) was determined according to the glucose oxidase peroxidase method. Impaired fasting glucose (IFG) was defined as a FG ≥ 100 mg/dL (22). Serum total cholesterol (TC), HDL-C, and triglycerides (TG) were determined by standard methods at the Clinical Research Laboratory, Chemistry School, UASLP in Mexico. Low-density lipoprotein-cholesterol (LDL-C) and very-low-density lipoprotein-cholesterol (VLDL-C) were calculated using Friedewald's equation. Hypertriglyceridemia was defined as a TG concentration ≥ 150 mg/dL, hypercholesterolemia as TC ≥ 200 mg/dL, high LDL-C as a concentration > 130 mg/dL, and low HDL-C was defined as ≤ 40 mg/dL in males and ≤ 50 mg/dL in females (25).

STATISTICAL ANALYSIS

Participants' general characteristics were categorized separately by birthweight category. A one-way analysis of variance (ANOVA) with post-hoc test was performed to evaluate significant differences in quantitative variables, and the χ^2 test was used for qualitative characteristics. Mean values for CV risk factors, such as serum lipid profile and glucose levels, as well as BP, were reported by using the ANOVA (*post-hoc* test) approach across

the different categories of birthweight. Statistical analyses were performed using the SPSS for Windows software (version 26.0. SPSS, Chicago, IL, USA), and p-values < 0.05 were considered statistically significant.

RESULTS

The number of women and men included in the present study was 469 (60.2 %) and 309 (39.7 %), respectively. The means and standard deviations (SDs) of general characteristics, CESD-10 scores, and family history across birthweight categories are shown in table I. Both groups with birthweight > 4000 g had the highest amount of weight and BMI ($p < 0.05$). No significant relationships were found between CESD-10. However, a significant relationship was found only for WC in men and for height in women.

Means and SDs for BP, lipid profile, and serum glucose levels according to birthweight categories are shown in table II. There was a significant association between SBP and DBP, but only DBP had statistical significance across birthweight categories. Participants with birthweight > 4000 g, had higher means for DBP ($p < 0.05$).

DISCUSSION

The gestational stage and early childhood represent a window of phenotypic plasticity, and represent a sensitive period as related to cardiometabolic risk programming. Maternal weight status, lipid profile, smoking habit, gestational diabetes *mellitus*,

or hypertensive disease during pregnancy condition birthweight and gestational age (26-29).

All these prenatal factors, along with postnatal factors like breastfeeding and rapid weight gain during childhood, are associated with cardiometabolic risk later in life (29-33). Genetic factors and epigenetic mechanisms play an important role in programming the genes correlated to cardiovascular and mental disease (34,35).

There are very few studies in developing countries that review the association between birthweight, components of MetS, and mental illness. We decided to investigate the possible association between these three variables in young Mexican adults. Our results are quite similar to those of previous studies (3). In the present study only HBW had an association with overweight and obesity. The reviewed studies demonstrated a J- or U-shaped relationship between these two variables (6,7). We also observed, in men only, a significant association between birthweight and central obesity ($p < 0.05$), which was only marginally identified among women ($p = 0.05$).

We found that SBP had statistical significance but was inconclusive at the post-hoc test. Only HBW was associated with high DPB. The results from studies examining the associations between HBW and hypertension are inconsistent. Several studies showed an inverse relationship between birthweight and BP (8,9). Nevertheless, a recent study in adolescents showed a positive association between HBW and hypertension (36).

There are studies that support the statement that cardiovascular risk factors are related to the transmission of mitochondrial DNA (6,37). Our study did not find a significant p-value for a family history of diabetes or cardiovascular disease in the maternal branch, but it could have been otherwise with more cases.

Table I. General characteristics, CESD-10 score, and family history of participants according to birthweight (g) categories

	Men birthweight (g)			p-value*	Women birthweight (g)			p-value*
	< 2500	2500-4000	> 4000		< 2500	2500-4000	> 4000	
Age (y)	17.8 (0.5)	17.9 (1.4)	18.8 (2)	< 0.05 [†]	17.6 (0.9)	17.8 (1.2)	18.1 (0.3)	0.42
BMI (kg/m ²)	21.7 (1.9)	23.8 (4.2)	25.7 (5.6)	< 0.05 [‡]	21.7 (3.5)	23.1 (4.4)	26.2 (6.9)	< 0.05 [‡]
Weight (kg)	62.1 (6.2)	70.1 (14.2)	77.3 (18.2)	< 0.05 [‡]	54.1 (9.8)	58.9 (12.7)	70.7 (20.8)	< 0.05 [§]
Height (m)	1.69 (0.05)	1.71 (0.06)	1.73 (0.07)	0.21	1.58 (0.05)	1.59 (0.06)	1.63 (0.07)	< 0.05 [†]
Waist (cm)	78.1 (5.9)	81.3 (11.3)	88.9 (12.5)	< 0.05	73.5 (8)	77.4 (11.9)	83.4 (16.2)	0.05
CESD-10	5 (2)	6 (5)	6 (4)	0.51	7 (5)	7 (4)	7 (6)	0.68
Family history of T2DM, % (n)								
Father	23.5 (4)	33.8 (93)	47 (8)	0.73	28.2 (13)	32.6 (135)	66.6 (6)	0.66
Mother	11.8 (2)	33.4 (92)	47 (8)		32.6 (15)	38.8 (161)	77.7 (7)	
Family history of CVD, % (n)								
Father	5.9 (1)	16.3 (45)	11.7 (2)	0.14	23.9 (11)	16.9 (70)	11.1 (1)	0.14
Mother	47.1 (8)	21.8 (60)	29.4 (5)		17.3 (8)	25.6 (106)	44.4 (4)	

Values are mean (SD) or percentage (number) if % is indicated. BMI: body mass index; CESD-10: center for epidemiological studies depression scale; T2DM: diabetes mellitus type 2; CVD: cardiovascular disease. *p-values result from an analysis of variance. [†]According to Tukey's post-hoc test, the p-value between birthweight > 4000 g and birthweight = 2500-4000 g is $p < 0.05$. [‡]According to Tukey's post-hoc test, the p-value between birthweight > 4000 g and birthweight < 2500 g is $p < 0.05$. [§]According to Tukey's post-hoc test, the p-value between all groups is $p < 0.05$. ^{||}According to Tukey's post-hoc test, the p-value between birthweight > 4000 g and the two other groups is $p < 0.05$.

Table II. Blood pressure, lipid profile, and serum glucose level (mean [SD]) according to birthweight categories

	Birthweight, g			p-value*
	< 2500	2500-4000	> 4000	
SBP (mmHg)	108 (9)	111 (10)	112 (8)	< 0.05 [†]
DBP (mmHg)	70 (8)	72 (8)	76 (8)	< 0.05 [‡]
TC (mg/dL)	170 (40)	163 (36)	161 (37)	0.39
TG (mg/dL)	110 (61)	107 (44)	105 (36)	0.83
HDL-C (mg/dL)	47 (11)	48 (11)	49 (12)	0.83
LDL-C (mg/dL)	100 (34)	95 (33)	92 (36)	0.35
VLDL-C (mg/dL)	22 (12)	21 (9)	21 (7)	0.83
FBG (mg/dL)	86 (8)	88 (8)	89 (5)	0.10

SBP: systolic blood pressure; DBP: diastolic blood pressure; TC: total cholesterol; TG: triglycerides; HDL-C: high density lipoprotein-cholesterol; LDL-C: low-density lipoprotein-cholesterol; VLDL-C: very low density lipoprotein-cholesterol; FBG: fasting blood glucose. *p-values result from the analysis of variance. [†]Statistical significance according to p-value but no difference between groups in Tukey's post-hoc test. [‡]According to Tukey's post-hoc test, the p-value between birthweight > 4000 g and the two other groups is $p < 0.05$.

Other CVD risk factors and components of MetS, including low HDL-C, and high levels of LDL-C, TC, TG, SBP and FBG, had no significant association with birthweight categories. Also, in this study we observed no association between birthweight categories and CESD-10 score (17).

We mentioned above that Dos Santos et al. found that birthweight was not a risk factor for the development of MetS during adolescence (13). This result was probably conditioned by the small size of their Brazilian cohort ($n = 172$), a possible bias in the selection of participants, or methodological differences in the evaluation of adiposity. Nevertheless, a study with a Mendelian randomization analysis studied 7 single nucleotide polymorphisms (SNPs) independently contributing to birthweight at genome-wide significance to estimate the effect of birthweight on risk of coronary ischemic disease, and they concluded that there was no strong evidence between birthweight, coronary disease, and lipid profile (38).

Despite this inconsistency among studies, there is a need to develop public health policies that adopt preventive measures to promote adequate maternal-fetal and child development, and enable an early diagnosis of metabolic abnormalities (28), as well as improvements in the policies related to healthy eating, demanding proper food labeling, promotion of exclusive breastfeeding for at least 6 months, and community-based primary prevention programs regarding lifestyle, among other measures (29,39). A study reveals iron supplementation in children with LBW as a measure to reduce systolic blood pressure on childhood (40). That is a novel observation suggesting that CVD risk factors like hypertension might be reduced with early interventions.

A low number of cases is the main limitation of this study. We only had 778 patients who fulfilled the selection criteria, 689 cases of NBW, 63 cases (8 %) of LBW, and 26 cases (3.3 %) of HBW, which corresponds to the percentages of LBW and HBW in the Mexican literature. The observed results have to be carefully ana-

lyzed. Other prenatal confounders like maternal obesity, smoking habit, and gestational diabetes *mellitus*, and postnatal confounders like breastfeeding duration, socioeconomic status, complementary feeding, catch-up growth, diet and physical activity, were not available and were not considered in the analysis. Due to the cross-sectional design we were unable to determine causality.

In conclusion, the findings of this population-based study revealed a positive relationship between birthweight categories and some CVD risk. Young adults that were born with HBW had association with high DPB, weight, and BMI. Depression was not related to birthweight categories.

CONCLUSIONS

The findings of this population-based study (young Mexican adults) revealed that depression was not related to birthweight. However, a positive relation between birthweight categories and some CVD risk factors such as DBP, weight, and BMI was shown in both genders. Furthermore, WC and height had a relationship with birthweight in men and women, respectively.

REFERENCES

1. Barker DJ. The origins of the developmental origins theory. *J Intern Med* 2007;261(5):412-7. DOI: 10.1111/j.1365-2796.2007.01809.x
2. Mandy M, Nyirenda M. Developmental Origins of Health and Disease: the relevance to developing nations. *International Health* 2018;10:66-70.
3. Azadbakht L, Kelishadi R, Saraf-Bank S, Qorbani M, Ardalan G, Heshmat R, et al. The association of birth weight with cardiovascular risk factors and mental problems among Iranian school-aged children: the CASPIAN-III study. *Nutrition* 2014;30:150-8. DOI: 10.1016/j.nut.2013.06.005
4. UNICEF-WHO. Low birth weight prevalence, by country and region; 2015
5. Ponce-Saavedra AS, González-Guerrero O, Rodríguez-García R, Echeverría-Landa A, Puig-Nolasco A, Rodríguez-Guzmán LM. Macrosomia prevalence and associated factors. *Rev Mex Pediatr* 2011;78:139-42.

6. Guerrero-Romero F, Aradillas-García C, Simental-Mendía LE, Monreal-Escalante E, Cruz-Mendoza E, Rodríguez-Moran M. Birth weight, family history of diabetes, and metabolic syndrome in children and adolescents. *J Pediatr* 2010;156:719-23. DOI: 10.1016/j.jpeds.2009.11.043
7. Wang J, Perona JS, Schmidt-RioValle J, Chen Y, Jing J, Gonzalez-Jimenez E. Metabolic Syndrome and Its Associated Early-Life Factors among Chinese and Spanish Adolescents: A Pilot Study. *Nutrients* 2019;11:1568. DOI: 10.3390/nu11071568
8. Richardson LJ, Hussey JM, Strutz KL. Origins of disparities in cardiovascular disease: birth weight, body mass index, and young adult systolic blood pressure in the national longitudinal study of adolescent health. *Ann Epidemiol* 2011;21:598-607. DOI: 10.1016/j.annepidem.2011.02.012
9. Chen W, Srinivasan SR, Yao L, Li S, Dasmahapatra P, Fernandez C, et al. Low birth weight is associated with higher blood pressure variability from childhood to young adulthood: the Bogalusa Heart Study. *Am J Epidemiol* 2012;176:S99-105. DOI: 10.1093/aje/kws298
10. Reinehr T, Kleber M, Toschke AM. Small for gestational age status is associated with metabolic syndrome in overweight children. *Eur J Endocrinol* 2009;160:579-84. DOI: 10.1530/EJE-08-0914
11. Zanetti D, Tikkanen E, Gustafsson S, Priest JR, Burgess S, Ingelsson E. Birth-weight, type 2 diabetes mellitus, and cardiovascular disease: Addressing the Barker Hypothesis with Mendelian Randomization. *Circ Genom Precis Med* 2018;11:e002054. DOI: 10.1161/CIRCGEN.117.002054
12. Sun D, Wang T, Heianza Y, Huang T, Shang X, Lv J, et al. Birthweight and cardiometabolic risk patterns in multiethnic children. *Int J Obes (Lond)* 2018;42:20-7. DOI: 10.1038/ijo.2017.196
13. Dos Santos-Alves PJ, Henriques AC, Pinto LR, Mota RM, Alencar CH, Alves RS, et al. Endothelial and metabolic disorders in adolescence: Low birth weight is not an isolated risk factor. *J. Pediatr Endocrinol. Metab* 2015;28:407-413. DOI: 10.1515/jpem-2014-0146
14. Rooij SR, Painter RC, Holleman F, Bossuyt PM, Roseboom TJ. The metabolic syndrome in adults prenatally exposed to the Dutch famine. *Am J Clin Nutr* 2007;86:1219-24. DOI: 10.1093/ajcn/86.4.1219
15. Nomura Y, Wickramaratne PJ, Pilowsky DJ, Newcorn JH, Bruder-Costello B, Davey C, et al. Low birth weight and risk of affective disorders and selected medical illness in offspring at high and low risk for depression. *Compr Psychiatry* 2004;48:470-8. DOI: 10.1016/j.comppsy.2007.04.005
16. Alati R, Najman JM, O'Callaghan M, Bor W, Williams GM, Clavarino A. Fetal growth and behaviour problems in early adolescence: findings from the Mater University Study of Pregnancy. *Int J Epidemiol* 2009;38:1390-400. DOI: 10.1093/ije/dyp252
17. Arafat S, Minică CC. Fetal Origins of Mental Disorders? An Answer Based on Mendelian Randomization. *Twin Res Hum Genet* 2018;21:485-94. DOI: 10.1017/thg.2018.65
18. Huffman JC, Celano CM, Januzzi JL. The relationship between depression, anxiety, and cardiovascular outcomes in patients with acute coronary syndromes. *Neuropsychiatr Dis Treat* 2010;6:123-36. DOI: 10.2147/ndt.s6880
19. O'Neil A, Williams ED, Stevenson CE, Oldenburg B, Berk M, Sanderson K. Comorbid cardiovascular disease and depression: sequence of disease onset is linked to mental but not physical self-rated health. Results from a cross-sectional, population-based study. *Soc Psychiatry Psychiatr Epidemiol* 2012;47:1145-51. DOI: 10.1007/s00127-011-0421-5
20. Santos-Veloso MAO, Souza Lacerda de Melo MI, Neves Cavalcanti RA, Soares Bezerra L, Chavez-Markman AV, Goncalves de Lima S. Prevalence of depression and anxiety and their association with cardiovascular risk factors in Northeast Brazil primary care patients. *Rev Assoc Med Bras* 2019;65:801-9. DOI: 10.1590/1806-9282.65.6.801
21. Raffaelli M, Andrade CD, Wiley AR, Sanchez-Armass O, Edwards LL, Aradillas-García C. Stress, social support, and depression: A test of the stress-buffering hypothesis in a Mexican sample. *Journal of Research on Adolescence* 2012;23:283-9. DOI: 10.1111/jora.12006
22. Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, et al. Harmonizing the metabolic syndrome: A joint interim statement of the International Diabetes Federation task force on epidemiology and prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 2009;20:1640-45. DOI: 10.1161/CIRCULATIONAHA.109.192644
23. Bojorquez CI, Salgado SN. Psychometric characteristics of the Center for Epidemiological Studies-Depression Scale (CES-D), 20-and10 item versions, in women from a Mexican rural area. *Salud Mental* 2009;33:299-307.
24. Raffaelli M, Drumond FC, Poppe EM, Sanchez-Armass O, Vazquez-Vidal I, Aradillas-García C. Gender and age-related differences in depressive symptoms and health behaviors among Mexican youth. *Int J Psycho* 2013;47:83-90. DOI: 10.30849/rip/ijp.v47i1.204
25. Jellinger PS, Handelsman Y, Rosenblit PD, Bloomgarden ZT, Fonseca VA, Garber AJ, et al. American Association of Clinical Endocrinologists and American College of Endocrinology Guidelines for management of dyslipidaemia and prevention of cardiovascular disease. *Endocr Pract* 2017;23:1-87. DOI: 10.4158/EP171764.APPGL
26. Yu Z, Han S, Zhu J, Sun X, Ji C, Guo X. Pre-pregnancy body mass index in relation to infant birth weight and offspring overweight/obesity: a systematic review and meta-analysis. *PLoS One* 2013;8:e61627. DOI: 10.1371/journal.pone.0061627
27. Herrera E, Ortega-Senovilla H. Lipid metabolism during pregnancy and its implications for fetal growth. *Curr Pharm Biotechnol* 2014;15:24-31. DOI: 10.2174/1389201015666140330192345
28. Martínez García RM, Jiménez Ortega AI, González Torres H, Ortega RM. Prevención de la obesidad desde la etapa perinatal. *Nutr Hosp* 2017;34:53-7. DOI: 10.20960/nh.1572
29. González-Jiménez E, Montero-Alonso MA, Schmidt-RioValle J, García-García CP. Metabolic syndrome in Spanish adolescents and its association with birth weight, breastfeeding duration, maternal smoking, and maternal obesity: a cross-sectional study. *Eur J Nutr* 2015;54:589-97. DOI: 10.1007/s00394-014-0740-x
30. Fujita Y, Kouda K, Nakamura H, Iki M. Association of rapid weight gain during early childhood with cardiovascular risk factors in Japanese adolescents. *J Epidemiol* 2013;23:103-8. DOI: 10.2188/jea.JE20120107
31. Lurbe E, Garcia-Vicent C, Torro MI, Aguilar F, Redon J. Associations of birth weight and postnatal weight gain with cardiometabolic risk parameters at 5 years of age. *Hypertension* 2014;63:1326-32. DOI: 10.1161/HYPERTENSIONAHA.114.03137
32. Okada T, Takahashi S, Nagano N, Usukura Y, Hosno S. Early postnatal alteration of body composition in preterm and small-for-gestational-age infants: implications of catch-up fat. *Pediatric Research* 2015;77:136-42. DOI: 10.1038/pr.2014.164
33. Kelishadi R, Haghdoost AA, Jamshidi F, Aliramezany N, Moosazadeh M. Low birthweight or rapid catch-up growth: which is more associated with cardiovascular disease and its risk factors in later life? A systematic review and meta-analysis. *Pediatrics and International Child Health* 2014;35:2:110-23. DOI: 10.1179/2046905514Y.0000000136
34. Lin X, Wu D, Gao L, Zhang J, Pan H, Wang H, et al. Altered DNA methylation in neonates born large-for-gestational age is associated with cardiometabolic risk in children. *Oncotarget* 2016;7:86511-21. DOI: 10.18632/oncotarget.13442
35. Carson C, Lawson HA. Epigenetics of metabolic syndrome. *Physiol Genomics* 2018;50(11):947-55. DOI: 10.1152/physiolgenomics.00072.2018
36. Kuciene R, Dulskiene V. Associations between high birth weight, being large for gestational age and high blood pressure among adolescents: a cross-sectional study. *Eur J Nutr* 2018;57:373-81. DOI: 10.1038/s41598-019-45956-9
37. Wilson FH, Hariri A, Farhi A, Zhao H, Petersen KF, Toka HR, et al. A cluster of metabolic defects caused by mutation in a mitochondrial tRNA. *Science* 2004;306:1190-4. DOI: 10.1126/science.1102521
38. Au Yeung, Lin S, Li A, Schooling M. Birth weight and risk of ischemic heart disease: A Mendelian randomization study. *Sci Rep* 2016;6:38420. DOI: 10.1038/srep38420
39. Glistrap LG, Malhotra R, Peltier-Saxe D, Slicas D, Pineda E, Culhane-Hermann C, et al. Community-based primary prevention programs decrease the rate of metabolic syndrome among socioeconomically disadvantaged women. *Journal of Women's Health* 2013;22:322-9. DOI: 10.1089/jwh.2012.3854
40. Lindberg J, Norman M, Westrup B, Domellöf M, Berglund SK. Lower systolic blood pressure at age 7 in low birth weight children who received iron supplements. *Am J Clin Nutr* 2017;106(2):475-80. DOI: 10.3945/ajcn.116.150482