



Trabajo Original

Paciente anciano

Impact of memory bias in the dietary estimation of older adults measured through a food frequency questionnaire and weighed food records – A cross-sectional study in older adults and primary caregivers

Impacto del sesgo de memoria en la estimación dietética de adultos mayores medido a través de un cuestionario de frecuencia de alimentos y registros de pesos de alimentos: estudio transversal en adultos mayores y cuidadores principales

Marcos Galván¹, Laura Ivonne Arellano-Martínez², Ivonne Ramírez-Silva³, Celina Ramírez-Ramírez¹, Guadalupe López-Rodríguez¹, Nely Isabel Cruz-Serrano⁴, Vidalma del Rosario Bezares-Sarmiento⁴, María Elena Rivera-Madrigal⁵

¹Academic Group of Nutritional Epidemiology. Facultad de Ciencias de la Salud. Universidad Autónoma del Estado de Hidalgo. Pachuca, Hidalgo. Mexico. ²Master's Program in Clinical Nutrition. Facultad de Ciencias de la Salud. Universidad Autónoma del Estado de Hidalgo. Pachuca, Hidalgo. Mexico. ³Center for Nutrition and Health Research. Instituto Nacional de Salud Pública. Cuernavaca, Morelos. Mexico. ⁴Lifestyle Research Group. Facultad de Ciencias de la Nutrición y Alimentos. Universidad de Ciencias y Artes de Chiapas. Tuxtla, Chiapas. México. ⁵Faculty of Ciencias de la Salud. Universidad Autónoma de Tlaxcala. Tlaxcala, Tlaxcala. Mexico

Abstract

Introduction: changes in cognitive performance and memory of older adults (OA) can interfere in their reporting their diet.

Objective: to evaluate the impact of memory bias in dietary estimation between OA and their primary caregivers (PC) through the food frequency questionnaire (FFQ) for Mexican OA and weighed food records (WFR).

Methods: the present analysis uses the estimated dietary information based on the response provided by 51 older adults (OA) and their primary caregivers (PC) from the validation study of the FFQ for Mexicans OA was conducted during lockdowns for COVID-19. The personnel who applied FFQ and WFR were trained with standardized instruments and procedures. The Wilcoxon test was used to compare the intake per day of the foods and food groups, the Spearman correlation coefficient was used to evaluate the grams of intake per day of the food groups, and kappa coefficient was used to compare the level of food items and food groups between OA and PC.

Results: in 11 of 14 food groups, no significant differences were observed between the amounts of intake reported by OA and PC. In the groups of dairy products, fruits, vegetables, and legumes, moderate agreement was observed ($\kappa = 0.63$ to 0.79), and in the rest of the groups was strong to perfect ($\kappa \geq 0.80$). Correlation of the amount of intake between OA and PC was high in all food groups ($r \geq 0.87$).

Conclusion: the high correlation and high agreement between the amounts and frequencies of the food groups consumed as reported by the older adults and primary caregivers indicate that the information from both respondents is reliable.

Keywords:

Memory. Food frequency questionnaire. Diet records. Caregivers. Older adults.

Received: 15/07/2023 • Accepted: 18/02/2024

Funding: this research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Acknowledgments: in this research, Fellows of the Youth Building the Future program (Mexico, 2020-2021), nutritionists from the Universidad de Ciencias y Artes de Chiapas, Universidad Autónoma de Tlaxcala, and Universidad Autónoma del Estado de Hidalgo participated. The forth author had a national postgraduate scholarship (No. 1034686) from the National Council of Science and Technology (CONACYT) of Mexico.

Conflicts of interest: the authors declare no conflicts of interest.

Artificial intelligence: the authors declare not to have used artificial intelligence (AI) or any AI-assisted technologies in the elaboration of the article.

Galván M, Arellano-Martínez LI, Ramírez-Silva I, Ramírez-Ramírez C, López-Rodríguez G, Cruz-Serrano NI, del Rosario Bezares-Sarmiento V, Rivera-Madrigal ME. Impact of memory bias in the dietary estimation of older adults measured through a food frequency questionnaire and weighed food records — A cross-sectional study in older adults and primary caregivers. *Nutr Hosp* 2024;41(3):574-584
DOI: <http://dx.doi.org/10.20960/nh.04858>

Correspondence:

Marcos Galván. Academic Group of Nutritional Epidemiology. Facultad de Ciencias de la Salud. Universidad Autónoma del Estado de Hidalgo. Ex-Hacienda la Concepción, s/n. 42160 Pachuca, Hidalgo. Mexico
e-mail: marcos_galvan3112@uaeh.edu.mx

Resumen

Introducción: los cambios cognitivos y la memoria que presentan los adultos mayores (AM) pueden interferir al momento de reportar los alimentos de su dieta.

Objetivo: evaluar el impacto del sesgo de memoria en la estimación de la dieta entre AM y sus cuidadores principales (CP) a través del cuestionario de frecuencia de alimentos (CFA) para AM mexicanos y el registro de peso de alimentos (RPA).

Métodos: el presente análisis utiliza la información dietética estimada con base en la respuesta proporcionada por 51 adultos mayores (AM) y sus cuidadores principales (CP) del estudio de validación del CFA para AM mexicanos que se realizó durante el confinamiento por COVID-19. El personal que aplicó CFA y RPA fue capacitado con instrumentos y procedimientos estandarizados. Se utilizó prueba de Wilcoxon para comparar la ingesta por día de los alimentos y grupos de alimentos, el coeficiente de correlación de Spearman para evaluar los gramos de ingesta por día de los grupos de alimentos y el coeficiente kappa para comparar el nivel de alimentos y grupos de alimentos entre AM y CP.

Resultados: en 11 de 14 grupos de alimentos no se observaron diferencias significativas entre las cantidades de ingesta reportadas por AM y CP. En los grupos de lácteos, frutas, verduras y legumbres se observó concordancia moderada ($\kappa = 0,63$ a $0,79$), y en el resto de los grupos fue de fuerte a perfecta ($\kappa \geq 0,80$). La correlación de la cantidad de ingesta entre AM y CP fue alta en todos los grupos de alimentos ($r \geq 0,87$).

Conclusión: La alta correlación y la alta concordancia entre las cantidades y frecuencias de los grupos de alimentos consumidos según lo informado por los adultos mayores y los cuidadores principales indican que la información de ambos encuestados es confiable.

Palabras clave:

Memoria. Cuestionario de frecuencia de alimentos. Registros de dieta. Cuidadores. Adultos mayores.

INTRODUCTION

The correct measurement of dietary intake in older adults (OA) is important to evaluate and identify unhealthy or risky eating patterns, monitor compliance with dietary recommendations, and promote health (1). In Mexico, all people older than 60 years are considered OA; this stage of life is characterized by changes in cognitive performance and long-term memory (2,3). Therefore, when evaluating diet in this age group, measures should be taken to reduce measurement errors. Accuracy in the evaluation of diet can be hampered by sociodemographic factors such as age, sex, culture, education, income, family environment, and deterioration in cognitive function and memory. The latter can affect the awareness of consumption and interfere with the understanding of the breakdown of ingredients, describing and reporting meals, recalling portion sizes, and knowledge of the foods and beverages consumed (4-6). There is limited research on the impact of cognitive function on the integrity of dietary data reported by OA (7).

One kind of instrument that has been most used to evaluate diet in different populations is FFQ. Lately, short-term questionnaires have begun to be used more, which are designed to evaluate the usual diet by asking about the frequency with which certain foods are consumed during a reference period (8). Compared to other methods of dietary evaluation, they are easy to administer, have a relatively low cost, can establish an association between diet and disease, and can be used to identify eating patterns related to inadequate consumption of specific nutrients (9-11). However, the lack of response can affect certain foods differently. For example, foods can be omitted because they were not consumed or because of difficulty remembering the frequency or amount consumed (12). Thus, the accuracy of the information collected can be substantially lowered.

The above problem has been little investigated in the of OA population of Mexico and Latin America (8,13,14). Therefore, the objective of the present study was to evaluate the impact of memory bias in dietary estimation between OA and their primary caregivers (PC) through the food frequency questionnaire (FFQ) for Mexican OA and weighed food records (WFRs).

METHODS

STUDY DESIGN

The information on the diet of older adults derived from the validation study of the Food Frequency Questionnaire for Mexican Older Adults was analyzed. The present analysis uses the estimated dietary information based on the response provided by the older adults (OA) and the primary caregivers (PC).

PARTICIPANTS

The original study on the validation of FFQ for Mexican OA was conducted during lockdowns for COVID-19, for which the OA who lived in the same house with students of university Nutrition programs were invited to participate through a video in the states of Hidalgo, Chiapas, and Tlaxcala, Mexico. To be eligible for the study, the OA had to be 60 years or older, have an oral diet, have no diagnosis of dementia, and have independent ambulation, and the PC had to be of legal age, live in the same house as the OA, and be responsible for feeding the OA. The sample size needed to detect a correlation equal to $r = 0.46$ of the total energy consumption between the OA and PC, with an alpha level of 0.05 and power of 0.90, was 50 participants. No restriction on the sexes was set.

PROCEDURE AND MEASURES

Food frequency questionnaire (FFQ) for Mexican older adults

FFQ is a semiquantitative questionnaire that consists of 194 food and beverage items to assess the current diet of OA. It was designed to resemble the FFQ developed for Mexico's National Health and Nutrition Surveys 2012 and 2016 (15,16). FFQ consists of 14 food groups: dairy products, fruits, vegetables, fast food, meats, sausages and eggs, fish, legumes, cereals and

tubers, corn products, beverages, snacks, sweets and desserts, and soups, creams and pastes, miscellaneous, and *tortillas*. It evaluates the number of times the respondent has consumed each food category in the last 7 days, which ranges from never to six times a day. The portion sizes of the foods come in predefined categories. The list of foods included in the questionnaire and the sizes of the standard portions were estimated from analyses performed on the foods most consumed in older adults from 24-hour recall national surveys conducted in 2012 and 2016.

The personnel who applied FFQ were trained virtually for 40 hours in theoretical-practical workshops, with standardized instruments and procedures. Two FFQ were delivered on paper to perform the recording, as well as a digital version with macros in Microsoft Excel, identical to the printed version of FFQ, to record the data. It was applied during the period from November 2020 to July 2021 in the homes of the OA and PC. The evaluators asked the study subjects to remember all the foods and their portion sizes consumed in the 7 days before the interview. Subjects who reported not having consumed the standard serving size were asked to select an alternative serving size, ranging from “very small” to “extra-large”. They were then asked to specify the number of days and the number of times per day that they consumed the food during the 7 days before the interview.

Weighed food records (WFR)

The weight or volume of food and beverages ingested over 7 continuous days of the week was recorded directly. To do this, it was necessary to weigh and measure all the foods that the older adult was served, and then the leftovers were weighed to obtain the total grams of the food consumed by difference. The personnel were trained to collect the information in a format designed for this study to make sure they used the scale correctly to weigh the foods and beverages.

Sociodemographic characteristics

The sociodemographic characteristics (age, sex, marital status, level of schooling, health perception, number of diseases, and number of medications taken) were gathered through an online electronic questionnaire, which was administered by the staff before applying FFQ.

STUDY VARIABLES

Food consumption

A result of no food consumption was recorded when the respondent chose the frequency category called “Never (code 01) for the different items.” The corresponding result was recorded if food consumption was reported in the frequency category from 1 time (code 02) to > 6 times per day (code 09).

Amount of food consumed

From the FFQ responses, the amount of each food and drink consumed was calculated by multiplying the number of days, the number of times per day, the serving size in grams (g) or milliliters (ml), and the number of servings consumed in each meal. The total grams and milliliters were divided by 7 days to obtain the daily intake, and these totals were added to the appropriate food groups.

For the questionnaire on weights and measures, the amount of food consumed per day was estimated, yielding the total amount served minus the amount leftover, giving the total amount of food or drink consumed. The grams consumed per day were added and divided by 7 to obtain the average daily intake. As for FFQ, these data were summed for each corresponding food group.

Percentage contribution of each food group

The percentages represented by the 14 food groups were categorized and calculated according to FFQ answered by the OA and PC. First the percentage contributions, including that of the beverage group, were calculated, and then the beverage group was discarded to compare the distribution of the food groups.

STATISTICAL ANALYSIS

Measures of central tendency and dispersion were estimated for continuous variables, as were frequencies and percentages for categorical variables, to describe the sociodemographic and health characteristics of all participants. To compare the dietary intake of each food and of the main food groups of FFQ between that reported by the OA and that reported by the PC, the medians and interquartile ranges (25th to 75th percentile) were estimated. The Wilcoxon test of paired samples was used to compare the intake per day of the foods and food groups reported by the OA and PC. To analyze the differences between consumers and non-consumers of each food group and the differences between sociodemographic groups, the chi-square test was applied.

To evaluate the grams of intake per day of the food groups between that reported by the OA and PC, Spearman correlation coefficients were calculated. Concordance tests were performed between the report on the consumption or not of the food given by the OA and PC, using the kappa coefficient at the level of food items and food groups. To evaluate the agreement on dietary intake in grams at the food item level and by food group, the subjects were classified into tertiles of consumption per day. The degree of agreement with respect to dietary intake was calculated by examining the proportion of subjects who fell into the same tertile or an adjacent tertile as that defined by the reference method (weights and measures) when they were classified by the OA or PC. The FFQ consumption estimated by the OA was compared with the consumption reported in weights and measures by the OA using the kappa coefficient. All statistical analyses were performed with the statistical software STATA version 14.0.

RESEARCH ETHICS AND PATIENT CONSENT

The original validation study of FFQ for Mexican OA was conducted according to the guidelines of the Declaration of Helsinki, and written informed consent was obtained from the participants. The project was approved by the Ethics Committee of the Autonomous University of the State of Hidalgo of the Institute of Health Sciences, which reviewed and approved the informed consent forms and the study protocol with number 2020-17.

RESULTS

The validation study was conducted in 51 older adults; 57 % were women, and 43 % were men. The average age of the OA was 70.4 ± 10.2 years; the average age of the PC was 29.8 ± 10.2. Fifty-five percent of the OA reported being married, 39.2 % reported having a primary education level, and the rest reported a higher level. Fifty-one percent perceived their state of health as normal, 54.9 % reported taking at least one medication, and 45.1 % had more than three diseases diagnosed by a doctor (Table I).

CONCORDANCE OF FOOD CONSUMPTION BETWEEN THE OA AND PC BY FFQ

The concordance value of the consumption of food groups reported by the OA and PC was almost perfect or strong (> 0.90) in fast food, meats, sausages and eggs, fish, corn products, beverages, snacks, sweets and desserts, soups, creams, pastas, and miscellaneous. Moderate agreement (0.64 to 0.79) was found for dairy products, fruits, vegetables, legumes, cereals and tubers, and *tortillas*. In all food groups, the percentage of agreement ranged from 92 % to 100 % (Table II). The previous concordances were corroborated with the analysis performed by item of each food (*n* = 178), 90 % of which presented an almost perfect or strong agreement; 7.3 % (*n* = 13 foods) showed a moderate agreement of 0.60-0.79 (items such as pastry, amaranth, rice, natural fruit water without sugar, fresh chillies, green tomato and tomato, strawberries with sugar or bananas with cream, baked toast without anything, sweets without sugar or light, cajeta, walnuts or almonds, and popcorn); 0.6 % (*n* = 1 food, animal butter) had a weak agreement of 0.40-0.59; and 1.7 % (*n* = 3 foods) no agreement of 0-0.20 (paste, empanada, or sweet vol-au-vent; vegetable juice; and fried foods) (Supplementary material: <https://www.nutricionhospitalaria.org/anexos/04858-01.pdf>).

CONCORDANCE OF FOOD CONSUMPTION BETWEEN OA AND PC BY FFQ AND WFR

The level of agreement of the consumption of food groups reported by the OA in FFQ compared to what they reported in the WFR was almost perfect or strong (> 0.94) for meats, sau-

Table I. Sociodemographic characteristics of the study population

Variable	Value
<i>Sex of older adult (OA), n (%)</i>	
Male	22 (43.0)
Female	29 (57.0)
<i>Sex of primary caregiver (PC), n (%)</i>	
Male	9 (18.0)
Female	42 (82.0)
<i>Age</i>	
PC (x ± DE)	29.8 ± 10.2
OA (x ± DE)	70.4 ± 10.2
<i>Marital status of OA, n (%)</i>	
Single	5 (9.8)
Married	28 (54.9)
Widowed	10 (19.6)
Divorced	8 (15.7)
<i>Level of schooling of OA, n (%)</i>	
None	8 (15.7)
Primary	20 (39.2)
Secondary	8 (15.7)
Baccalaureate	3 (5.9)
Bachelor	12 (23.5)
<i>Health perception of OA, n (%)</i>	
Good	23 (45.1)
Bad	2 (3.9)
Normal	26 (51.0)
<i>Frequency of OA diseases, n (%)</i>	
1-2	16 (31.4)
> 3	23 (45.1)
No	12 (23.5)
<i>Frequency of use of OA medications, n (%)</i>	
1-2	16 (31.4)
> 3	12 (23.5)
No	23 (45.1)

sages and eggs, cereals and tuber vegetables, beverages, and *tortillas*. A moderate agreement (0.63 to 0.67) was observed for dairy and fish. Weak agreement (0.40 to 0.46) was found for fruits, vegetables, fast food, corn products, beverages, snacks, sweets and desserts, soups, creams, and pastas; and there was a minimum level of agreement (22 to 26) for maize and legume products. The level of agreement between what the PC reported on FFQ and on the WFR was almost perfect or strong (> 0.94) for dairy products, meats, sausages and eggs, cereals and tubers, beverages, and *tortillas*; it was moderate (0.63 to 0.67) for vegetables and fish; it was weak (0.40 to 0.46) for fast food, snacks, sweets and desserts, soups, creams, and pastas; and there was minimum agreement (22 to 26) for fruits, corn products, and legumes (Table III).

Table II. Concordance of the consumption of food groups reported by older adults (OA) and primary caregivers (PC) with food frequency questionnaire (FFQ) for Mexican older adults

Food groups	Older adult <i>n</i> = 51 If consumed (%)	Primary caregiver <i>n</i> = 51 If consumed (%)	% agreement	κ	Level of agreement
Dairy	88.2	94.1	94.1	0.638	Moderate
Fruits	94.1	96.1	98.0	0.790	Moderate
Vegetables	94.1	96.1	98.0	0.790	Moderate
Fast food	45.0	45.0	96.1	0.920	Almost perfect
Meats, sausages and eggs	100.0	100.0	100.0	1.000	Almost perfect
Fish	32.3	37.2	98.0	0.957	Almost perfect
Legumes	82.3	84.3	94.1	0.788	Moderate
Cereals and tubers	98.0	100.0	98.0	0.790	Moderate
Corn products	82.3	84.3	98.0	0.929	Almost perfect
Beverages	98.0	98.0	100.0	1.000	Almost perfect
Snacks, sweets and desserts	66.7	70.6	96.0	0.909	Strong
Soups, creams and pastas	66.7	66.7	100.0	1.000	Almost perfect
Miscellaneous	43.1	50.9	92.2	0.843	Strong
<i>Tortillas</i>	100.0	98.0	98.0	0.790	Moderate

% agreement: number of participants who give the same response to FFQ-OA and FFQ-PC / total number of participants. κ : kappa coefficient.

CORRELATION OF THE AMOUNT OF FOOD INTAKE BETWEEN THE OA AND PC IN FFQ

Table IV shows the median and interquartile range of the intake in grams per day of food groups by the OA and PC in FFQ. The OA reported higher dairy intake (229.6 g) than the PC (142.2 g) ($p < 0.003$), as well as in the group of snacks, sweets, and desserts, with 53.8 g reported by OA compared to 50.6 g reported by PC ($p < 0.04$). The correlation coefficients of consumption in grams by food group between the OA and PC ranged from 0.87 for meats, sausages, and eggs to 0.98 for fruits, snacks, sweets and desserts, soups, creams, and pastas. In general, the correlations for all food groups were considered high (Table IV). In the vegetable group, the OA reported a lower consumption (119.5 g) than the PC (124.6 g) ($p < 0.13$). For the other 11 groups, no significant differences were observed between the consumption reported by the OA and PC. Only in 1.1 % ($n = 2$ foods) of the 178 foods was there a significant difference in the median consumption reported by the OA compared to the PC, which were the items added onion and fresh cheese (Supplementary material: <https://www.nutricionhospitalaria.org/anexos/04858-01.pdf>).

CONCORDANCE OF FOOD INTAKE BETWEEN THE OA AND PC IN FFQ

To evaluate the concordance of food intake between the OA and PC with FFQ, a cross-classification analysis was performed. We defined that a subject was correctly classified if their intake per day was in the same tertile or in an adjacent tertile. The concordance of consumption between what was reported by the OA and PC, estimated at the level of food groups in tertile 1, varied from 78.9 % (soups, creams, and pastas) to 100 % (fast food, meats, sausages and eggs, fish, and miscellaneous), with a discrepancy of less than 18 %. In tertile 2, the agreement ranged from 75 % (fish) to 100 % (fast food, corn products, miscellaneous, and *tortillas*), and the disagreement was less than 25 %. In tertile 3, the agreement between the subjects was the best classified, ranging from 97.9 % (vegetables) to 100 % (dairy, fruits, fast food, meats, sausages and eggs, fish, legumes, snacks, sweets, desserts, miscellaneous, and *tortillas*), with a discrepancy of less than 7 % (Table V).

Table III. Concordance of food consumption between older adults and primary caregivers in a Mexican older adult food frequency questionnaire and weighed food records

Food groups	Older adult		% agreement	κ	Level of agreement	Primary caregiver		% agreement	κ	Level of agreement
	Weights and measures if you consumed n (%)	Frequency of Consumption Questionnaire if you consumed n (%)				Weights and measures if they consumed n (%)	Frequency of consumption questionnaire if they consumed n (%)			
Dairy	48 (94)	45 (88)	94.1	0.63	Moderate	48 (94)	48 (94)	100.0	1.00	Almost perfect
Fruits	46 (90)	48 (94)	92.2	0.46	Weak	46 (90)	49 (96)	90.2	0.24	Minimum
Vegetables	50 (98)	48 (94)	92.2	0.46	Weak	50 (98)	49 (96)	94.1	0.62	Moderate
Fast food	19 (37)	23 (45)	80.4	0.59	Weak	19 (37)	23 (45)	80.4	0.59	Weak
Meats, sausages and eggs	51 (100)	51 (100)	100.0	1.00	Almost perfect	51 (100)	51 (100)	100.0	1.00	Almost perfect
Fish	22 (43)	18 (35)	84.3	0.67	Moderate	22 (43)	19 (37)	86.3	0.71	Moderate
Legumes	43 (84)	42 (82)	78.4	0.22	Minimum	43 (84)	43 (84)	80.4	0.25	Minimum
Cereals and tubers	51 (100)	50 (98)	98.0	0.94	Almost perfect	51 (100)	51 (100)	100.0	1.00	Almost perfect
Corn products	48 (94)	42 (82)	84.3	0.26	Minimum	48 (94)	42 (82)	84.3	0.26	Minimum
Beverages	51 (100)	50 (98)	98.0	0.86	Strong	51 (100)	50 (98)	98.0	0.94	Almost perfect
Snacks, sweets and desserts	42 (82)	34 (67)	76.5	0.40	Weak	42 (82)	36 (70)	80.4	0.46	Weak
Soups, creams and pastas	43 (84)	34 (67)	78.4	0.44	Weak	43 (84)	34 (67)	78.4	0.44	Weak
Miscellaneous	51 (100)	22 (43)	43.1	0.00	None	51 (100)	26 (50)	51.0	0.00	None
Tortillas	51 (100)	51 (100)	100.0	1.00	Almost perfect	51 (100)	50 (98)	98.0	0.94	Almost perfect

% agreement: number of participants who give the same response to FFQ-OA and FFQ-PC / total number of participants. κ: kappa coefficient.

Table IV. Correlation of the amount of food group intake per day as reported by the older adult and the primary caregiver on FFQ

Food groups	Older adult n = 51 M (p25, p75)	Primary caregiver n = 51 M (p25, p75)	p-value	Correlation ^a
Dairy	229.6 (35.3, 584.6)	142.2 (38.9, 333.9)	0.003*	0.94
Fruits	217.9 (62.9, 401.5)	211.5 (63.6, 390.4)	0.090	0.98
Vegetables	119.5 (54.3, 266.2)	124.6 (73.6, 243.1)	0.013*	0.91
Fast food	27.4 (14.1, 39.4)	31.4 (14.1, 39.4)	0.317	0.94
Meats, sausages and eggs	171.3 (96.4, 254.7)	196.7 (106.6, 264.9)	0.062	0.87
Fish	16.4 (10.0, 38.6)	12.8 (10, 38.6)	0.317	0.97
Legumes	67.0 (30.7, 154.3)	57.8 (15.1, 147.3)	0.317	0.91
Cereals and tubers	129.3 (82.1, 182.4)	114.4 (83.8, 181.2)	0.312	0.90
Corn products	121.5 (48.3, 217.9)	128.6 (48.3, 224.1)	0.114	0.96
Beverages	1005.3 (672.2, 2047.1)	1148.2 (758.6, 2061.1)	0.147	0.94
Snacks, sweets and desserts	53.8 (19.7, 87.5)	50.6 (17.8, 78.8)	0.041*	0.98
Soups, creams and pastas	131.1 (68.9, 235.2)	128.9 (68.9, 211.9)	0.340	0.98
Miscellaneous	9.5 (3.9, 14.3)	8.2 (2.8, 14.3)	0.083	0.89
Tortillas	170.0 (75.0, 267.1)	170.0 (75.0, 252.8)	0.754	0.88

FFQ: consumption frequency questionnaire for Mexican older adults; p25: 25th percentile; p75: 75th percentile; *p < 0.001, p < 0.01, or p < 0.05, Wilcoxon test; ^aSpearman correlation.

Table V. Agreement by tertile of consumption of food groups per day between older adults (OA) and primary caregivers (PC) on FFQ

Food groups	Tertile 1		Tertile 2		Tertile 3		% agreement	κ
	Concordant n (%)	Discordant n (%)	Concordant n (%)	Discordant n (%)	Concordant n (%)	Discordant n (%)		
Dairy	39 (84.8)	7 (15.2)	34 (87.2)	5 (12.8)	42 (100.0)	0	90.5	0.858
Fruits	68 (97.1)	2 (2.9)	47 (92.2)	4 (7.8)	60 (100.0)	0	96.7	0.949
Vegetables	89 (91.7)	8 (8.3)	86 (93.5)	6 (6.5)	93 (97.9)	2 (2.1)	94.4	0.915
Fast food	12 (100.0)	0	10 (100.0)	0	11 (100.0)	0	100.0	1.000
Meats, sausages and eggs	82 (100.0)	0	89 (97.8)	2 (2.2)	66 (100.0)	0	98.3	0.974
Fish	10 (100.0)	0	3 (75.0)	1 (25.0)	6 (100.0)	0	95.0	0.918
Legumes	18 (81.8)	4 (18.2)	17 (89.5)	2 (10.5)	19 (100.0)	0	90.0	0.850
Cereals and tubers	60 (95.2)	3 (4.8)	57 (98.3)	1 (1.7)	59 (96.7)	2 (3.3)	96.7	0.950
Corn products	28 (96.6)	1 (3.4)	28 (100.0)	0	28 (96.5)	1 (3.5)	97.7	0.965
Beverages	68 (98.6)	1 (1.4)	61 (91.0)	6 (9.0)	66 (95.7)	3 (4.3)	94.7	0.919

(Continues on next page)

Table V (cont.). Agreement by tertile of consumption of food groups per day between older adults (OA) and primary caregivers (PC) on FFQ

Food groups	Tertile 1		Tertile 2		Tertile 3		% agreement	κ
	Concordant n (%)	Discordant n (%)	Concordant n (%)	Discordant n (%)	Concordant n (%)	Discordant n (%)		
Snacks, sweets and desserts	25 (89.3)	3 (10.7)	25 (86.2)	4 (13.8)	26 (100.0)	0	91.6	0.873
Soups, creams and pastas	15 (78.9)	4 (21.1)	11 (84.6)	2 (15.4)	14 (93.3)	1 (6.7)	85.1	0.777
Miscellaneous	15 (100.0)	0	18 (100.0)	0	12 (100.0)	0	100.0	1.000
Tortillas	21 (91.3)	2 (8.7)	24 (100.0)	0	21 (100.0)	0	97.1	0.955

FFQ: consumption frequency questionnaire for Mexican older adults; % agreement: number of participants who give the same response to FFQ-OA and FFQ-PC / total number of participants. κ: kappa coefficient.

CONTRIBUTION OF FOOD GROUPS TO THE OA DIET

To evaluate the percentage contribution to the diet made by the different food groups according to what the OA or PC reported, two analyses were performed. In Analysis A, the beverage group was included. The highest percentage contribution was made by beverages (OA: 41 %, PC: 45 %), followed by fruits (OA: 9 %, PC: 8 %), dairy (OA: 9 %, PC: 5.6 %), and

meats, sausages, and eggs (OA: 6 %, PC: 8 %). In analysis B, which excluded beverages, the greatest contribution was made by fruits (OA: 15 %, PC: 15 %), followed by dairy products (OA: 16 %, PC: 10 %), meats, sausages, and eggs (OA: 12 %, PC: 14 %), *tortillas* (OA: 12 %, PC: 12 %), and soups, creams and pastas (OA: 9 %, PC: 9 %). There were significant differences between OA and PC in the reported median consumption of dairy products (0.003), vegetables (0.013), and snacks (0.041) (Table VI).

Table VI. Percentage of contribution to the diet by food groups reported by older adults and primary caregivers with FFQ

Food groups	A. With beverages				A. No beverages				p-value
	Older adult		Primary caregiver		Older adult		Primary caregiver		
	g/day (M)	%	g/day (M)	%	g/day (M)	%	g/day (M)	%	
Dairy	229.6	9.3	142.2	5.6	229.6	15.6	142.2	10.3	0.003*
Fruits	217.9	8.8	211.5	8.4	217.9	14.8	211.5	15.4	0.090
Vegetables	119.5	4.8	124.6	4.9	119.5	8.2	124.6	9.0	0.013*
Fast food	27.4	1.1	31.4	1.2	27.4	1.9	31.4	2.3	0.317
Meats, sausages and eggs	171.3	6.9	196.7	7.8	171.3	11.7	196.7	14.3	0.062
Fish	16.4	0.7	12.8	0.5	16.4	1.1	12.8	0.9	0.317
Legumes	67.0	2.7	57.8	2.3	67.0	4.6	57.8	4.2	0.317
Cereals and tubers	129.3	5.2	114.4	4.5	129.3	8.8	114.4	8.3	0.312
Corn products	121.5	4.9	128.6	5.1	121.5	8.3	128.6	9.3	0.114
Beverages	1005.3	40.7	1148.2	45.5					0.147
Snacks, sweets and desserts	53.8	2.3	50.6	2.0	53.8	3.7	50.6	3.7	0.041*
Soups, creams and pastas	131.1	5.3	128.9	5.1	131.1	9.0	128.9	9.4	0.340

(Continues on next page)

Tabla VI (cont.). Percentage of contribution to the diet by food groups reported by older adults and primary caregivers with FFQ

Food groups	A. With beverages				A. No beverages				<i>p</i> -value
	Older adult		Primary caregiver		Older adult		Primary caregiver		
	g/day (M)	%	g/day (M)	%	g/day (M)	%	g/day (M)	%	
Miscellaneous	9.5	0.4	8.2	0.3	9.5	0.7	8.2	0.6	0.083
<i>Tortillas</i>	170.0	6.9	170.0	6.8	170.0	11.6	170.0	12.3	0.754
Total	2538.5	100.0	2558.7	100.0	1445.7	100.0	1451.2	100.0	

FFQ: consumption frequency questionnaire for Mexican older adults; M: median; g/day: grams per day; * $p < 0.001$, $p < 0.01$, or $p < 0.05$, Wilcoxon test.

As additional data, the consumption of individual beverages was analyzed, which showed that the most consumed beverage was plain water (OA: 27.7 %; PC: 32.3 %), followed by natural fruit water without sugar (OA: 24.9 %; PC: 8.7 %), and coffee (OA: 8.5 %; PC: 10.3 %). There were no significant differences between the consumption reported by the OA and PC (Supplementary material: <https://www.nutricionhospitalaria.org/anexos/04858-01.pdf>).

DISCUSSION

A relevant correlation was seen between the intake reported by the OA and the PC in FFQ, with an almost perfect or a high concordance for most food groups. Were no significant differences in the amounts reported in grams or milliliters. However, based on the estimated agreement between the consumption of foods obtained from FFQ and that reported on the weights and measures questionnaire, foods from the dairy group, vegetables, and beverages were better captured when the questionnaire was answered by the PC. These results indicate that the repeatability in general and therefore the reliability of FFQ is good. According to what was reported by the OA and PC in terms of weights and measures, the findings coincide with other studies, in which approximately 80 % of the items were significantly correlated between two administrations of the questionnaire, using weighed food records as a reference method 17. In this study, 85 % of the participants were classified as concordant in the intake, with the highest observed agreement of 100 % for the intake of dairy products; meats, sausages, and eggs; cereals and tubers; beverages; and *tortillas*. Therefore, only 15 % of the participants were misclassified, with the lowest value being 3 % for miscellaneous. The results of agreement in this study are comparable to those reported in a validation study that applied a printed FFQ to 62 women aged 31-60, approximately 70 % of whom reported intakes in the same quintile or in an adjacent quintile (18). On the other hand, in a study with 115 men and 115 women in Brazil aged 20-50, comparing a FFQ and a 24-hour recall questionnaire, a percentage of agreement was found that varied from 45.1 % for energy to 26.2 % for unsaturated fats. Therefore, it

can be said that the agreements are higher when weights and measures are used as the reference method.

Regarding the amount of intake of food groups reported by OA and PC, significant differences were only found in three of the 14 food groups, which could be because OA tend to overestimate their real intake when asked to remember the frequency of many foods consumed. This has been observed, for example, by Liu et al. and Dehghan et al. (11,20), who found that FFQ overestimate energy and nutrient values. Another study has reported that the estimated intake of energy and nutrients derived from an FFQ was higher than the estimates derived from the 24-hour questionnaires (21). Bijani et al, in their study of OA, reported that the medians in the FFQ were significantly higher for whole grains, dairy products, vegetables, fruits, sugar, eggs, tea, and coffee ($p < 0.05$), and the consumption of sandwiches and desserts was significantly different among women (22). A study in Mexico with subjects between 18 and 71 years of age reported that a habitual intake of food and nutrient groups through FFQ tended to be higher than those reported through a logbook (23).

In the total grams of intake reported for food groups, a relatively high proportion of subjects were classified correctly (in the same tertile), and few were classified erroneously (less than 7 %). In a study in Brazil, with people older than 20 years filling out a FFQ in duplicate, concordances of 0.36 to 0.65 were found, which were lower than those in our study (24). Thompson et al, in a study of adults between 25 and 70 years old filling out a FFQ, found a percentage of agreement of 84.5 %, a value lower than that found in our study (25). Kowalkowska et al., in their study of subjects between 15 and 65 years old filling out a self-administered FFQ compared to one administered by an evaluator, found a concordance of 70.3 % to 99.4 %, values lower than those reported in the present study (26). Thus, in our study, a greater agreement was obtained for FFQ when applied to the OA and PC, which gives validity to the diet data from both respondents.

We also found a high correlation of consumption between the 14 food groups reported by the OA and the PC ($r = 0.87$ $r = 0.98$). Several studies in adults and OA have found low to moderate correlations, ranging from 0.50 for fish and cereals, 0.40-0.50 for fruits, 0.30-0.45 for vegetables, and 0.18 for

white/red meats (12,27-30). Selem et al reported a correlation between the application of two FFQ of 0.36 to 0.69, which was lower than that in the present study (24). The information obtained from the OA and PC in this study is reliable for evaluating dietary habits and can be used to support the design of interventions in nutrition at the individual and collective levels. Even so, we must remember that the correlations vary not only by food group but also according to the nutrient, so in a second phase of analysis, it would be convenient to evaluate these (10,31-33). The high correlations found in our study indicate that a rigorous process of training of the evaluators was followed, since high correlations are associated with a better result when FFQ are administered by a professional than when self-administered (34).

In the evaluation of the percentage contributions of the food groups to the diet of the OA, this research showed that more than 50 % of the diet was obtained from fruits, dairy products, meats, sausages, eggs, *tortillas*, and cereals and tubers. Rodríguez et al reported that *tortillas* and bread were the two main sources of energy for OA (21). The contribution of *tortillas* to total energy consumption was estimated at 42 % by 24-hour dietary recall and 37.6 % by an FFQ, which indicated a more varied diet that had more fruits and vegetables when reported using the FFQ than when using the 24-hour recall, similar to what we found. Geri et al reported different dietary patterns among OA in Argentina (35). One was the traditional varied diet, which was characterized by consuming similar proportions of milk, cheese, egg, beef, vegetables, fresh fruits, bread, oil, free sugars, and sausages. Anderson et al, in participants between 70 and 79 years old from the USA, evaluated the group of healthy foods, characterized by a relatively higher intake of low-fat dairy products, fruits, whole grains, poultry, fish, and vegetables and a lower consumption of meat, fried foods, sweets, beverages with high energy content, and added fats (36). Their findings differed from what was reported in this study, where the diet of OA was characterized by foods typical of a traditional diet; however, we did find a low proportion of vegetable consumption.

Our study has several limitations that should be considered. First, the sample was small (51 OA), which made it hard to analyze the data of men and women separately. Not all the results could be compared with similar studies since little evidence was found in the literature. Its main strength lies in the fact that this is the first study in Mexican OA that compares the diet reported by the OA and the PC. It shows that in healthy OA, it is possible to apply FFQ, since their memory is still accurate. FFQ designed specifically for Mexican OA are effective at obtaining dietary information, as demonstrated by Patterson et al., keeping in mind the need to modify FFQ for populations with unique dietary patterns (37). It should be noted that the present study was conducted in OA with good cognitive functioning, and it is possible that our conclusions are not applicable to OA with cognitive impairment related to age or disease.

This study found no evidence that there is an impact of memory bias in the dietary estimation of Mexican OA, as no

significant differences were found in the consumption of different food groups estimated by OA and PC. There was a high correlation and agreement between the amount and frequency consumed of the food groups reported by the OA and PC. In this study, the characterization of the OA diet was similar to that reported by the OA and PC. Therefore, the information from both informants is reliable enough to support the design of food interventions at the individual level and in groups of OA. Additionally, the design of a FFQ specifically for the elderly population can substantially improve the ability to capture their dietary patterns as completely as possible.

REFERENCES

- Olendzki B, Procter-Gray E, Magee MF, Youssef G, Kane K, Churchill L, et al. Racial Differences in Misclassification of Healthy Eating Based on Food Frequency Questionnaire and 24-Hour Dietary Recalls. *J Nutr Health Aging* 2017;21(7):787-98. DOI: 10.1007/s12603-016-0839-2
- Villegas-Vázquez KG, Montoya-Arce BJ. Condiciones de vida de los adultos mayores de 60 años o más con seguridad social en el Estado de México. *Papeles Poblac* 2014;20(79):133-67.
- Reuter-Lorenz PA, Park DC. Human Neuroscience and the Aging Mind: A New Look at Old Problems. *Journals Gerontol Ser B Psychol Sci Soc Sci* 2010;65B(4):405. DOI: 10.1093/geronb/gbq035
- Reagh ZM, Roberts JM, Ly M, DiProspero N, Murray E, Yassa MA. Spatial discrimination deficits as a function of mnemonic interference in aged adults with and without memory impairment. *Hippocampus* 2014;24(3):303. DOI: 10.1002/hipo.22224
- Higgs S. Memory for recent eating and its influence on subsequent food intake. *Appetite* 2002;39(2):159-66. DOI: 10.1006/appe.2002.0500
- Hebert JR, Ebbeling CB, Matthews CE, Hurley TG, MA Y, Druker S, et al. Systematic errors in middle-aged women's estimates of energy intake: Comparing three self-report measures to total energy expenditure from doubly labeled water. *Ann Epidemiol* 2002;12(8):577-86. DOI: 10.1016/S1047-2797(01)00297-6
- Zuniga K, McAuley E. Considerations in selection of diet assessment methods for examining the effect of nutrition on cognition. *J Nutr Health Aging* 2015;19(3):333. DOI: 10.1007/s12603-014-0566-5
- Willett WC. Food frequency methods. In: Willett WC, edit. *Nutritional Epidemiology*, 3rd edition. Oxford: Oxford University Press 2013:96-141. DOI: 10.1093/acprof:oso/9780199754038.001.0001
- Vian I, Zielinsky P, Zilio AM, Mello A, Lazzeri B, Oliveira A, et al. Development and validation of a food frequency questionnaire for consumption of polyphenol-rich foods in pregnant women. *Matern Child Nutr* 2015;11(4):511-24. DOI: 10.1111/mcn.12025
- Subar AF, Thompson FE, Kipnis V, Midthune D, Hurwitz P, McNutt S, et al. Comparative Validation of the Block, Willett, and National Cancer Institute Food Frequency Questionnaires The Eating at America's Table Study. *Am J Epidemiol* 2001;154(12):1089-99. DOI: 10.1093/aje/154.12.1089
- Liu L, Wang PP, Roebouthan B, Ryan A, Tucker CS, Colbourne J, et al. Assessing the validity of a self-administered food-frequency questionnaire (FFQ) in the adult population of Newfoundland and Labrador, Canada. *Nutr J* 2013;12(1):49. DOI: 10.1186/1475-2891-12-49
- Procter-Gray E, Olendzki B, Kane K, Churchill L, Hayes RB, Aguirre A, et al. Comparison of Dietary Quality Assessment Using Food Frequency Questionnaire and 24-hour-recalls in Older Men and Women. *AIMS public Heal* 2017;4(4):326. DOI: 10.3934/publichealth.2017.4.326
- Block G, Hartman AM. Issues in reproducibility and validity of dietary studies. *Am J Clin Nutr* 1989;50(5):1133-8. DOI: 10.1093/ajcn/50.5.1133
- Cruz-Góngora V, Martínez-Tapia B, Cuevas-Nasu L, Flores-Aldana M, Shamah-Levy T. Dietary intake and adequacy of energy and nutrients in Mexican Older Adults: Results from a Food Frequency Questionnaire of two National Health and Nutrition Surveys, 2012 and 2006. *Salud Publica Mex* 2017;59(3). DOI: 10.21149/7851
- Ramírez-Silva I, Jiménez-Aguilar A, Valenzuela-Bravo D, Martínez-Tapia B, Rodríguez-Ramírez S, Gaona-Pineda EB, et al. Methodology for estimating dietary data from the semi-quantitative food frequency questionnaire of the Mexican National Health and Nutrition Survey 2012. *Salud Publica Mex* 2016;58(6):629-38. DOI: 10.21149/spm.v58i6.7974

16. Denova-Gutiérrez E, Ramírez-Silva I, Rodríguez-Ramírez S, Jiménez-Aguilar A, Shamah-Levy T, Rivera-Dommarco JA. Validity of a food frequency questionnaire to assess food intake in Mexican adolescent and adult population. *Salud Publica Mex* 2016;58(6):617-28. DOI: 10.21149/spm.v58i6.7862
17. Gardener SL, Lyons-Wall P, Martins RN, Rainey-Smith SR. Validation and reliability of the Alzheimer's disease-commonwealth scientific and industrial research organisation food frequency questionnaire. *Nutrients* 2020;12(12):1-17. DOI: 10.3390/nu12123605
18. Lassale C, Guilbert C, Keogh J, Syrette J, Lange K, Cox DN. Estimating food intakes in Australia: validation of the Commonwealth Scientific and Industrial Research Organisation (CSIRO) food frequency questionnaire against weighed dietary intakes. *J Hum Nutr Diet* 2009;22(6):559-66. DOI: 10.1111/j.1365-277X.2009.00990.x
19. da Silva NF, Sichieri R, Pereira RA, da Silva RMVG, Ferreira MG. Reproducibility, relative validity and calibration of a food frequency questionnaire for adults. *Cad Saude Publica* 2013;29(9):1783-94. DOI: 10.1590/0102-311X00120312
20. Dehghan M, Martinez S, Zhang X, Seron P, Lanas F, Islam S, et al. Relative validity of an FFQ to estimate daily food and nutrient intakes for Chilean adults. *Public Health Nutr* 2013;16(10):1782-8. DOI: 10.1017/S1368980012004107
21. Rodríguez MM, Méndez H, Torún B, Schroeder D, Stein AD. Validation of a semi-quantitative food-frequency questionnaire for use among adults in Guatemala. *Public Health Nutr* 2002;5(5):691-8. DOI: 10.1079%20%20PHN2002333
22. Bijani A, Esmaili H, Ghadimi R, Babazadeh A, Rezaei R, Cumming RG, et al. Development and validation of a Semi-quantitative food frequency questionnaire among older people in north of Iran. *Casp J Intern Med* 2018;9(1):78. DOI: 10.22088/cjim.9.1.78
23. Macedo-Ojeda G, Vizmanos-Lamotte B, Fabiola Márquez-Sandoval Y, Patricia Rodríguez-Rocha N, López-Uriarte PJ, Fernández-Ballart JD. Validation of a semi-quantitative food frequency questionnaire to assess food groups and nutrient intake. *Nutr Hosp* 2013;28(6):2212-20. DOI: 10.3305/nh.2013.28.6.6887
24. Selem SS de C, de Carvalho AM, Verly-Junior E, Carlos JV, Teixeira JA, Marchioni DML, et al. Validity and reproducibility of a food frequency questionnaire for adults of São Paulo, Brazil. *Rev Bras Epidemiol* 2014;17(4):852-9. DOI: 10.1590/1809-4503201400040005
25. Thompson FE, Subar AF, Brown CC, Smith AF, Sharbaugh CO, Jobe JB, et al. Cognitive research enhances accuracy of food frequency questionnaire reports: results of an experimental validation study. *J Am Diet Assoc* 2002;102(2):212-25. DOI: 110.1016/s0002-8223(02)90050-7
26. Kowalkowska J, Wadolowska L, Czarnocinska J, Czlapka-Matysik M, Galinski G, Jezewska-Zychowicz M, et al. Reproducibility of a Questionnaire for Dietary Habits, Lifestyle and Nutrition Knowledge Assessment (KomPAN) in Polish Adolescents and Adults. *Nutrients* 2018;10(12). DOI: 10.3390/nu10121845
27. Kristjansdottir AG, Andersen LF, Haraldsdottir J, de Almeida MDV, Thorsdottir I. Validity of a questionnaire to assess fruit and vegetable intake in adults. *Eur J Clin Nutr* 2005;60(3):408-15. DOI: 10.1038/sj.ejcn.1602332
28. Rodríguez IT, Ballart JF, Pastor GC, Jordà EB, Val VA. Validation of a short questionnaire on frequency of dietary intake: reproducibility and validity. *Nutr Hosp* 2008;23(3):242-52.
29. Shatenstein B, Payette H. Evaluation of the Relative Validity of the Short Diet Questionnaire for Assessing Usual Consumption Frequencies of Selected Nutrients and Foods. *Nutrients* 2015;7(8):6362. DOI: 10.3390/nu7085282
30. Beck KL, Houston ZL, McNaughton SA, Kruger R. Development and evaluation of a food frequency questionnaire to assess nutrient intakes of adult women in New Zealand. *Nutr Diet* 2020;77(2):253. DOI: 10.1111/1747-0080.12472
31. Gosadi IM, Alatar AA, Otayf MM, Aljahani DM, Ghabbani HM, Alrajban WA, et al. Development of a saudi food frequency questionnaire and testing its reliability and validity. *Saudi Med J* 2017;38(6):636-41. DOI: 10.15537/smj.2017.6.20055
32. Gardener SL, Lyons-Wall P, Martins RN, Rainey-Smith SR. Validation and Reliability of the Alzheimer's Disease-Commonwealth Scientific and Industrial Research Organisation Food Frequency Questionnaire. *Nutrients* 2020;12(12):1-17. DOI: 10.3390/nu12123605
33. Serna Gutiérrez A, Esparza-Romero J. Diseño y validación de un cuestionario de frecuencia de consumo de alimentos para evaluar la dieta en indígenas yaquis de Sonora, México. *Acta Univ* 2019;29:1-16. DOI: 10.15174/au.2019.2248
34. Cade J, Thompson R, Burley V, Warm D. Development, validation and utilisation of food-frequency questionnaires a review. *Public Health Nutr* 2002;5(4):567-87. DOI: 10.1079/PHN2001318
35. Geri M, Gutiérrez E, González G. Dietary patterns of older adults in an Argentine region and their relationship with socioeconomic factors. *Rev Salud Pública* 2019;21(6):1-5. DOI: 10.15446/rsap.v21n6.82930
36. Anderson AL, Harris TB, Tylavsky FA, Perry SE, Houston DK, Hue TF, et al. Dietary Patterns and Survival of Older Adults. *J Am Diet Assoc* 2011;111(1):84. DOI: 10.1016/j.jada.2010.10.012
37. Patterson RE, Kristal AR, COates RJ, Tylavsky FA, Ritenbaugh C, Van Horn L, et al. Low-fat diet practices of older women: prevalence and implications for dietary assessment. *J Am Diet Assoc* 1996;96(7):670-6. DOI: 10.1016/s0002-8223(96)00186-1