



Original article

## Cognitive profile associated with functional and anthropometric aspects in elderly



W. Araújo de Brito<sup>a</sup>, L. Mendes<sup>a</sup>, M. Magalhães Sales<sup>b</sup>, J.B. Neto<sup>b</sup>, C.J. Brito<sup>e</sup>, M.E. da Silva Grigoletto<sup>c</sup>, A. Pimentel Ferreira<sup>d,\*</sup>

<sup>a</sup> Curso de Enfermagem das Faculdades Integradas Promove de Brasília e Faculdades ICESP, Brasília, Brazil

<sup>b</sup> Programa de Pós-Graduação em Educação Física da Universidade Católica de Brasília – UCB, Brasília, Brazil

<sup>c</sup> Programa de Pós Graduação em Educação Física da Universidade Federal de Sergipe – UFS, Sergipe, Brazil

<sup>d</sup> Núcleo Interdisciplinar de Pesquisa – NIP das Faculdades Integradas Promove de Brasília e Faculdades ICESP, Brasília, Brazil

<sup>e</sup> Universidade Federal de Juiz de Fora, UFJF, Minas Gerais, Brazil

### ARTICLE INFO

*Article history:*

Received 23 February 2014

Accepted 16 February 2015

Available online 6 September 2016

*Keywords:*

Aging

Dementia

Anthropometry

### ABSTRACT

**Objective:** To identify the cognitive profile of the elderly and its association with anthropometric and functional aspects.

**Methodology:** It is a cross-sectional study conducted in 84 elderly. Measured up the following anthropometric variables: body weight, height, waist circumference, hip circumference for subsequent calculation of the conicity index, waist/hip ratio and waist/height ratio. Furthermore, the functional aspects were measured based on the battery of tests adapted from Functional Fitness Test and cognitive impairment using the Mini Mental State Examination.

**Results:** Dementia was found in 7% of the elderly, with no difference between genders. The waist/height ratio above 0.67 cm increased by 5.6 times the chance to develop dementia, while the balance showed a protective factor (15% less likely).

**Conclusion:** There was a low prevalence of dementia, the Waist Circumference and Hip Circumference indexes demonstrated a predictive capacity for dementia while the balance pattern was considered a protective factor.

© 2016 Consejería de Turismo y Deporte de la Junta de Andalucía. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Perfil cognitivo asociado a aspectos funcionales y antropométricos en personas mayores

### RESUMEN

**Objetivo:** Identificar el perfil cognitivo de personas ancianas y su asociación con los aspectos funcionales y antropométricos.

**Metodología:** Estudio transversal descriptivo realizado en 84 personas ancianas. Fueron recogidos datos referentes a las medidas antropométricas: masa corporal, estatura, circunferencia de la cintura, circunferencia de la cadera, para posterior cálculo del índice de conicidad, relación cintura/cadera y la relación cintura/estatura. Además, fueron medidos datos funcionales basados en una serie de pruebas adaptadas del *Functional Fitness Test* y el déficit cognitivo fue evaluado a través del Mini Examen del Estado Mental.

**Resultados:** La demencia encontrada fue del 7% en las personas mayores de edad. La relación cintura/estatura mayor de 0.67 cm aumentó en 5.6 veces las posibilidades de desarrollar demencia, mientras que el equilibrio presentó un factor de protección (15% menos de probabilidad de presencia de demencia).

**Conclusión:** Hubo baja frecuencia de demencia, la circunferencia de la cintura y la relación cintura/cadera presentaron una capacidad predictiva, mientras que el equilibrio fue considerado un factor de protección.

© 2016 Consejería de Turismo y Deporte de la Junta de Andalucía. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

*Palabras clave:*

Personas mayores

Demencia

Antropometría

\* Corresponding author.

E-mail addresses: [cidopimentel@yahoo.com.br](mailto:cidopimentel@yahoo.com.br), [cidopimentel@gmail.com](mailto:cidopimentel@gmail.com) (A. Pimentel Ferreira).

## Perfil cognitivo associado a aspectos funcionais e antropométricos em idosos

### R E S U M O

#### Palavras-chave:

Idoso  
Demência  
Antropometria

**Objetivo:** Identificar o perfil cognitivo de idosos e a sua associação com aspectos funcionais e antropométricos.

**Métodos:** Estudo transversal descritivo realizado em 84 idosos. Mensuraram-se as seguintes variáveis antropométricas: massa corporal, estatura, circunferência da cintura (CC), circunferência do quadril (CQ), para posterior cálculo do índice de conicidade (IC), relação cintura/quadril (RCQ) e relação cintura/estatura (RCE). Ademais, foram medidos os aspectos funcionais baseados na bateria de testes adaptada do *Functional Fitness Test* e o déficit cognitivo, por meio do Mini Exame do Estado Mental.

**Resultados:** A demência foi encontrada em 7% dos idosos, não havendo diferença entre gêneros. A RCE acima de 0.67 cm aumentou em 5.6 vezes a chance para desenvolver demência, enquanto o equilíbrio apresentou fator de proteção (15% menos chances).

**Conclusão:** Houve baixa prevalência de demência, a CC e RCE apresentaram capacidade preditiva, enquanto o equilíbrio foi considerado fator de proteção.

© 2016 Consejería de Turismo y Deporte de la Junta de Andalucía. Publicado por Elsevier España, S.L.U. Este é um artigo Open Access sob uma licença CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### Introduction

During the last few decades, there is a new demographic behavior in the elderly, in which there is a progressive increase in the prevalence of the elderly population due to improvements in life expectancy and the increase in the proportion of elderly in the world's population. However, the search for strategies that seek to prolong the life expectancy free of functional and cognitive impairment is relevant. The elderly population is a group characterized by heterogeneity, so that we can find in the same age group, elderly with preserved functional capacity and the ones with memory impairment levels in an advanced state.<sup>1</sup>

The aging is a continuous process in which there is a progressive decline in physiological processes. Among the most apparent physiological changes during aging, there is a decrease in height by about 1 cm per decade and structural and functional changes in the nervous system, due to reduction in the number of neurons. It is also usual to have increased arterial stiffness, increased blood pressure, a decreased cardiac output at rest and during exercise, and reduced bone mineral density.<sup>2–4</sup>

Specifically, regarding the changes that affect the central nervous system with the course of the aging process, there is the ventriculomegaly and reduced cortical mantle, usually accompanied by progressive cognitive impairment. However, regardless of degenerative diseases, cognitive dysfunction and dementia were observed. Dementia is the progressive reduction or loss of cognitive abilities, partial or complete, permanent or momentary form, enough to cause a loss of autonomy and functional decline.<sup>2</sup>

For Pedrosa and Holland,<sup>5</sup> loss of independence and low levels of cardiorespiratory fitness are associated with risk of morbidity and mortality from chronic diseases, on the other hand the performance of daily activities of life and the maintenance of independence are linked to improved general health. In this sense, low levels of muscular strength and obesity are associated with functional decline and cognitive impairment.<sup>3,6</sup> In addition, important measures related to functional capacity and maintenance of daily activities as the amount of motor units, the contraction speed, neurons driving speed, the reaction time, and volume of brain matter show worsening over the decrease in walk velocity.<sup>1,4,7</sup>

Accordingly, it appears that functional capacity, maintaining the autonomy are associated with a lower incidence of chronic degenerative diseases. Thus, there is a constant need for assessments and maintaining physical fitness, since it features associated with general health in the elderly, particularly in relation to cardiovascular

disease, cognitive response and hospitalization.<sup>2</sup> In this light, the aim of this study was to evaluate the cognitive profile of the elderly and its association with anthropometric and functional aspects.

### Methods

#### Subjects

The sample consisted of 84 elderly of both sexes (15 male and 69 female) aged between 61 and 90 years living in the Federal District, selected by convenience. The inclusion criteria were: have over 60 years of age, a resident in own home, sign the informed consent term, participate in all data collections, show no neuropathy or musculoskeletal problem that precludes participation in physical tests.

Before data collection, the Ethics Committee on Research of the Hospital of the Forces Army (HFA) under protocol 016/2011/CEP/HFA approved this study. Volunteers or their guardians signed a free and informed consent form (ICF).

#### Experimental design

After contact with the elderly in their homes, we collected data related to anthropometric measurements, the Mini Mental State Examination (MMSE), and functional testing. The adapted Functional Fitness Test was used in this study.<sup>8–12</sup>

**Romberg test:** The researcher asked each volunteer to stay with his/her feet together, barefoot, with open eyes for about 30 s and then with eyes closed for 30 s. The standing position without loss of balance clocked the permanence time and the test was considered positive when the volunteer has failed to spend more than 10 s with their eyes closed without body oscillations. In the time each volunteer had to have the first oscillation, considered equilibration time, was recorded similar to the procedure adopted by Cho et al.<sup>8</sup>

**Handgrip test:** After familiarization, the participants were seated, with the shoulder in neutral position, elbows bent at 90° and wrist in a neutral position. The subjects were instructed to perform a maximal isometric contraction to verbal command of the examiner. Three attempts were made by alternating members, with 60 s interval. No encouragement was given by the researcher during the test. The procedure was carried out respecting the criteria described by Shiratori et al.<sup>9</sup>

**Stand up and walk test (TSW) (Timed up and go – TUG):** This test was measured using a stopwatch and recorded in seconds the time it takes to stand up from a chair, walk a distance of 3 m, turn around an obstacle and sit again. We used a chair 45 cm in height for the seat and 65 cm in height to the armrest with backrest, brand Grosfillex®. The TSW was done according to the procedure described by Hyun et al.<sup>10</sup>

**Chair get up test:** For this test, we used a chair with traditional back, without lateral support, with 45 cm height, and a stopwatch. The chair was supported on the wall to prevent the same from moving during the test period. A pre-test for familiarization was conducted to ensure that the participant sitting occupies most of the seat with his/her feet flat on the floor. He was spoken aloud each time the participant until the fifth time. This test was performed according to procedure described by Yamauchi et al.<sup>11</sup>

**Mini Mental State Examination (MMSE):** The MMSE consists of typical questions grouped into 7 categories, each of them designed in order to assess specific cognitive functions, with them being: time orientation, local, words registration, attention and calculation, memory, language, and visual constructive capacity. Leading questions and punctuation establishment based on pre-established scores on the instrument obtained responses. The cutoff points used in this study were adopted as model, as used previously in another study by Ferreira et al.<sup>12</sup>

**Body mass and height:** For the body mass a digital Britania® balance with a resolution of 0.1 kg was used. For this measurement, individuals were with minimum clothing, positioned to his feet, and remained motionless until the stabilization of the value displayed on device. A tape fixed to the wall was used for the height measurement. The head was positioned according to the Frankfurt plan in which an imaginary line was passing through the lowest point of the inferior orbital rim and the highest point of the upper edge of the right auditory meatus.<sup>13</sup> After these adjustments, the most prominent point of the head to the realization of the measure was noted.

**Waist and hip circumference:** The measurement of waist circumference (WC) was carried out with a tape, 150 cm in length and resolution of 0.1 cm. It was measured in line with the navel, and evaluated without shirt. The measurement of the hip circumference (WHR) was held at the leading region of the hip, assessed with feet together and legs straight.<sup>14</sup>

**Arm circumference:** To measure the circumference of the arm, the same was flexed toward the chest at a 90° angle, and located and marked the midpoint between the acromion and olecranon.

It was requested to the individual to stay with the arm extended along the body and with the palm facing down.<sup>15</sup>

**Thigh circumference:** The tape was placed across, half way between the inguinal line and the upper border of the patella. Feet were slightly apart and body weight distributed equally between them.<sup>14</sup>

**Neck circumference:** The measurement was performed with the participant standing with the back straight and head in the Frankfurt horizontal plane. The tape was positioned in the lower neck circumference just above the laryngeal prominence (Adam's apple).

### Statistical analysis

Initially the data normality was checked, using the Shapiro–Wilk test when the group was stratified to 50 individuals and Kolmogorov–Smirnov when the group was more than 50 individuals. Descriptive statistics and frequency of measures were used to assess the prevalence of hypertension, diabetes, labyrinthitis, osteoporosis, physical activity and nutritional evaluation between different groups. We used the chi-square test to analyze the frequency ratios between the different groups. The test “t” independent was applied to analyze the anthropometric, hemodynamic differences, mental state and performance of functional tests in accordance with the level of physical activity, nutritional classification and sex.

To analyze the cutoff points of anthropometric indicators and performance of the studied functional tests that could identify dementia the ROC curve technique (receiver operating characteristic) was adopted. The bivariate regression models and multivariate logistic regression were used to calculate the Odds Ratio (OR), so that all the functional capacity and anthropometric variables were tested as controls in the regression models. *p* value < 0.05 was adopted to point out the significant differences. Statistical analysis was performed using SPSS 18.0 (SPSS Inc., Chicago, Illinois, USA) for Windows, and the Stata software version 9.1 (College Station, TX: StataCorp LP).

### Results

Tables 1 and 2 presented the characteristics of the sample, with anthropometric data, hemodynamics performance and results of the MMSE test as well as the percentages of dementia and other chronic diseases.

**Table 1**  
Characterization sample with anthropometric data and hemodynamic performance.


	Men n = 15	Women n = 69	Total sample n = 84
Body mass (kg)	76.2 ± 10.0	68.9 ± 12.0*	68.3 ± 13.3
Height (m)	1.6 ± 0.1	1.6 ± 0.1†	1.55 ± 0.1
Body mass index (kg/m <sup>2</sup> )	28.5 ± 2.7	27.4 ± 4.4	27.6 ± 4.2
Conicity index	1.4 ± 0.1	1.4 ± 0.1	1.4 ± 0.1
Waist hip ratio (cm)	1.0 ± 0.1	0.9 ± 0.1†	1.0 ± 0.1
Waist height ratio (cm)	0.64 ± 0.1	0.6 ± 0.1	0.6 ± 0.1
Systolic blood pressure (mmHg)	136 ± 13	131 ± 13	132 ± 13
Diastolic blood pressure (mmHg)	92 ± 13	85 ± 10	86 ± 11
Mean arterial pressure (mmHg)	106 ± 12	100 ± 10	101 ± 11
Neck circumference (cm)	43.0 ± 2.9	38.9 ± 4.0†	39.1 ± 3.9
Arm circumference (cm)	31.4 ± 2.3	32.0 ± 4.2	32.0 ± 4.3
Thigh circumference (cm)	49.1 ± 4	49.0 ± 6.1	48.5 ± 5.6
Hip circumference (cm)	103.0 ± 6.8	103.0 ± 11.1	104.0 ± 9.7
Waist circumference (cm)	104.0 ± 9.3	97.0 ± 9.8†	98.0 ± 10.6
Dominance handgrip (kg)	17.0 ± 8.0	20.4 ± 8.3	19.9 ± 8.3
Non-dominance handgrip (kg)	15.5 ± 8.5	18.7 ± 9.9	18.2 ± 9.7
Test stand and walk (s)	10.2 ± 3.3	11.3 ± 4.1	11.1 ± 4.0
Chair get up (s)	11.2 ± 6.0	12.1 ± 9.4	11.9 ± 3.5
Mini-Mental State Examination (score)	21.4 ± 3.8	21.9 ± 4.0	21.8 ± 4.0

\* Significant difference between men versus women.

**Table 2**  
Characterization of dementia data frequency and chronic diseases.

	Men	Women	Total sample
Obesity	77%	56.5 <sup>*</sup>	57.2%
Hypertension	92.3%	73.2 <sup>*</sup>	76.2%
Diabetes Mellitus	38.5%	31%	32.1%
Labyrinthitis	23%	32.4%	31%
Osteoporosis	15.4%	26.8%	25%
Dementia	7.7%	7%	7.1%

\* Significant difference between men versus women.

**Fig. 1.** ROC curves for the anthropometric indicators and functional tests as predictors of dementia. Abbreviations: BMI, body mass index; DH strength, dominant hand palmar pinch strength; NDH strength, nondominant hand palmar pinch strength; ROC, receiver operating characteristic; WHtR, waist-to-height ratio.

There was a statistically significant difference for the variables: body weight, height, WHR, neck circumference and WC, in which men showed higher values when compared to women.

**Table 3** shows the cutoffs and their sensitivity and specificity of anthropometric indicators and tests that measured the functional capacity in predicting dementia.

According to **Table 3**, the WC and the waist/height ratio (WHeR) were considered predictors of dementia, since the lower limit of the confidence interval was above 0.50 (Li-IC > 0.5). Already body mass index (BMI), taper index, waist/hip ratio (WHR) and performance tests, as carried out in our methodology, showed no reliable values of sensitivity and specificity (Li-CI < 0.5). **Fig. 1** shows the ROC curves of the key variables tested.

**Table 4** presented the odds ratio of anthropometric measures and functional performance for dementia for the total sample.

Initially the sex was tested as modifier effect; however, it showed no change in capacity, and we chose to analyze only the total sample without stratification by sex. Among the anthropometric and functional variables analyzed, only WHeR and the equilibrium showed risk/protection, respectively as significant. Seniors with above 0.67 cm WHeR were 5.63 times more at risk

**Table 3**  
Cutoffs, sensitivity and specificity of anthropometric measures and functional tests as predictors of dementia.

Dementia (n = 84)	Cutoffs	Sensitivity	Specificity	Area under curve
Waist (cm)	100	83.3	65.4	0.75 (0.57–0.93)
Body mass index (kg/m <sup>2</sup> )	28	67	53	0.71 (0.47–0.94)
Conicity index	1.41	50	62	0.57 (0.36–0.78)
Waist hip ratio (cm)	0.92	50	52	0.46 (0.25–0.67)
Waist height ratio (cm)	0.67	67	77	0.77 (0.61–0.93)
Dominance handgrip (kg)	24	67	39	0.38 (0.16–0.59)
Non-dominance handgrip (kg)	28	67	32	0.33 (0.11–0.54)
Test stand and walk (s)	10.6	66	51	0.61 (0.3–0.92)

**Table 4**  
Odds Ratio for dementia for the total sample.

Measures	Total sample
Body mass index (kg/m <sup>2</sup> ) > 27 kg/m <sup>2</sup>	3.75 (0.5–30)
Conicity index > 1.25	1
Waist > 100 cm	8.15 (0.99–66)
Waist height ratio (cm) > 0.67	5.63 (1.11–28) <sup>†</sup>
Sedentary lifestyle	1.36 (0.26–7.0)
Dominance handgrip (kg) > 20	0.53 (0.10–2.7)
Non-dominance handgrip (kg) > 20	0.36 (0.05–2.9)
Test stand and walk (s) > 10.59	0.5 (0.05–2.6)
Balance	0.15 (0.04–0.65) <sup>†</sup>
Chair get up (s)	0.65 (0.12–3.2)

\* Significant odds ratio.

of presenting dementia while older people who have balance were 15% less likely to have dementia.

## Discussion

The main results of this study showed that the prevalence of dementia verified by the results of the MMSE in the elderly was 7%, and that among the anthropometric and functional variables tested, only WC and WHeR showed predictive capacity to dementia. When analyzed the odds ratio, it was found that the maker of the top WHeR to 0.67 cm is 5.6 times more likely to have dementia, while the balance of the presence means protection factor by 15%. It is important to understand that the MMSE results only suggest the presence of dementia, and despite having high correlation with the same, more specific tests are necessary for the diagnosis to be confirmed.

We believe that the highest amount of abdominal fat was verified by WC and WHeR. According to Matsudo et al.,<sup>16</sup> the anthropometric measures (greater deposition of fat in the central region) are associated with lower levels of physical fitness and a less active lifestyle. The gain in body weight as well as body fat accumulation particularly in the central region appears to result from a standard genetically programmed, changes in diet and physical activity level, age-related or an interaction between these factors. Lower levels of muscle strength, as well as obesity (BMI ≥ 30), higher waist circumference and percentages of fat are associated with functional decline.<sup>3</sup>

The grip strength is the only muscle strength measurement recommended for clinical practice, so that it is increasingly used for clinical diagnosis in elderly.<sup>17</sup> Lower strength scores can be considered an independent predictor of overall mortality in the elderly over 85 years noninstitutionalized.<sup>18</sup> The lower level physical fitness, lifestyle and excess weight are closely related to social isolation and physical mobility, and in some cases lead to loss of functional independence and neurological disorders, affecting performance cognitive test, as observed in this study. With increasing age, people become less active, their physical capabilities diminish and psychological changes appear as a feeling of old age, stress and depression. Concurrently, there is the appearance of chronic

diseases that contribute to deteriorate the aging process; however, it seems that the disuse of physiological functions creates more problems than most of the effects of aging.<sup>16</sup>

In this sense, our study found that the balance was considered a protective factor for dementia. Since subjects with higher performance, in the balance of capacity, possibly present greater physical mobility and functional autonomy, which corroborates Borges et al.,<sup>19</sup> appointed that the practice of physical activity and functional autonomy both promote rehabilitation, as maintaining the overall health of the elderly maintaining physical fitness, independence and autonomy in activities of daily living. Currently there are various methods for measuring body stability, which is an important component in maintaining the standing position of equilibrium and during daily activities and movements, especially for elderly people, since they are considered to fall-risk population.<sup>20</sup> The system postural control depends on an intact ability (unharmed) to correctly perceive the environment by peripheral sensory system, as well as the processing and integration of proprioceptive pathways, visual and vestibular in the central nervous system.<sup>20</sup>

The risk of falls increases with aging, and can have a devastating effect on the independence and quality of life. Often it leads to a vicious cycle of inactivity with future decline in functionality, increased risk of falls, and more likely to need an attendant. Balance tests are useful tools to diagnose the loss of this physical ability. Encouraging seniors to improve physical feature can help them maintain independence and reduce the need for live assistance.<sup>21</sup> Corroborating the findings of this study, Hanna et al.,<sup>22</sup> indicated that the moderately active people presented a lower risk by mental disorders than those sedentary, showing that participation in physical exercise programs exerts benefits in the physical domain. Furthermore, psychologically and physically active people probably have a faster cognitive processing. This can be explained in part by a direct and indirect effect of exercise on cognitive function, such that the mechanisms that act directly increase the speed of cognitive processing, leading to an improvement in cerebral circulation and changes in the synthesis and degradation neurotransmitters.

In addition, direct mechanisms, such as low blood pressure, decrease in LDL and triglyceride in the blood plasma and inhibition of platelet aggregation appear to act indirectly by enhancing these functions and the overall functional capacity, reflecting in this way on mental health in the elderly.<sup>23</sup> The prevalence data of our study were similar to those presented by Oliveira et al.,<sup>24</sup> which also show that cognitive impairments in the elderly detected by the MMSE were not associated with the mobility test assessed by TSW. These results coincide with the literature and can be explained by the fact that some cases of dementia, patients have excessive movement and unrest for much of the day. What they are not perceived in Alzheimer's dementia due to lack of extrapyramidal signs and motor slowness.

One limitation of the present data is the sample size and the disproportion between male and female elderly; however, the literature still lacks investigations like this, making relevant the results of this study. Another limitation was the instrument used since the MMSE despite being a widely used test, still lacks studies to better determine its use, as well as the adoption of appropriate points of cuts in the Brazilian population. Applicable, the present study showed low prevalence of dementia in the elderly; however, it is important to keep the concern for variables related to functional capacity, as well as the anthropometric profile, as they present relationship with the cognitive deficits.<sup>25–27</sup> In this sense, health professionals should provide an environment that encourages physical activity safely, maintenance of functional capacity and avoidance of excess weight, particularly in the central part of the body.

In conclusion, the prevalence of dementia in the elderly was considered lower, however similar between male and female. Waist circumference and waist/height ratio showed predictive ability for dementia verified by MMSE, and seniors with greater deposition of fat in the central region of the body showed 5.6 times more likely to have dementia, while those with more balanced presence of a protective factor, with 15%, were less likely to have dementia.

### Conflicts of interest

The authors have no conflicts of interest to declare.

### Acknowledgment

This study was funded by grant from the Brazilian Scholarship Program *Ciência sem Fronteiras* (grant reference #234243/2014-7). The funders had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

### References

- Abellan van Kan GA, Rolland Y, Andrieu S, Bauer J, Beauchet O, Bonnefoy M, et al. Gait speed at usual pace as a predictor of adverse outcomes in community-dwelling older people an International Academy on Nutrition and Aging (IANA) task force. *J Nutr Health Aging*. 2009;13:881–9.
- Cooper R, Kuh D, Cooper C, Gale CR, Lawlor DA, Matthews F, et al. Objective measures of physical capability and subsequent health: a systematic review. *Age Ageing*. 2011;40:14–23.
- Schaap LA, Koster A, Visser M. Adiposity, muscle mass, and muscle strength in relation to functional decline in older persons. *Epidemiol Rev*. 2013;35:51–65.
- Cooper R, Kuh D, Hardy R, Mortality Review Group: FALCon and Halcyon Study Teams. Objectively measured physical capability levels and mortality: systematic review and meta-analysis. *BMJ*. 2010;341:c4467.
- Pedrosa R, Holanda G. Correlação entre os testes da caminhada, marcha estacionária e TUG em hipertensas idosas. *Rev Bras Fisioter São Carlos*. 2009;13:252–6.
- Benedet AL, Moraes CF, Camargos EF, Oliveira LF, Souza VC, Lins TC, et al. Amerindian genetic ancestry protects against Alzheimer's disease. *Dement Geriatr Cogn Disord*. 2012;33:311–7.
- Stenholm S, Maggio M, Lauretani F, Bandinelli S, Ceda GP, Di Iorio A, et al. Anabolic and catabolic biomarkers as predictors of muscle strength decline: the In CHIANTI study. *Rejuvenation Res*. 2010;13:3–11.
- Cho GH, Hwangbo G, Shin HS. The effects of virtual reality-based balance training on balance of the elderly. *J Phys Ther Sci*. 2014;26:615–7.
- Shiratori AP, Iop Rda R, Borges Júnior NG, Domenech SC, Gevaerd Mda S. Evaluation protocols of hand grip strength in individuals with rheumatoid arthritis: a systematic review. *Rev Bras Reumatol*. 2014;54:140–7.
- Hyun J, Hwangbo K, Lee CW. The effects of pilates mat exercise on the balance ability of elderly females. *J Phys Ther Sci*. 2014;26:291–3.
- Yamauchi T, Islam MM, Koizumi D, Rogers ME, Rogers NL, Takeshima N. Effect of home-based wellrounded exercise in community-dwelling older adults. *J Sports Sci Med*. 2005;4:563–71.
- Ferreira LS, Pinho Mdo S, Pereira MW, Ferreira AP. Perfil cognitivo de idosos residentes em Instituições de Longa Permanência de Brasília-DF. *Rev Bras Enferm*. 2014;67:247–51.
- Costa RF. Composição corporal: teoria e prática da avaliação. São Paulo: Manole; 2001.
- Leite MJCIC. Métodos de avaliação da composição corporal. Faculdade de Ciências da Nutrição e Alimentação da Universidade do Porto; 2004. p. 1–55 [Monografia].
- Sperotto FM, Spinelli RB. Avaliação nutricional em idosos independentes de uma instituição de longa permanência no município de Erechim – RS. *Perspectiva Erechim*. 2010;34:105–16.
- Matsudo SM, Matsudo VKR, de Barros Neto TL. Impacto do envelhecimento nas variáveis antropométricas, neuromotoras e metabólicas da aptidão física. *Rev Bras Ciên Mov*. 2000;8:21–32.
- Roberts HC, Denison HJ, Martin HJ, Patel HP, Syddall H, Cooper C, et al. A review of the measurement of grip strength in clinical and epidemiological studies: towards a standardised approach. *Age Ageing*. 2011;40:423–9.
- Takata Y, Ansay T, Soh I, Awano S, Yoshitake Y, Kimura Y, et al. Physical fitness and 6.5-year mortality in an 85-year-old community-dwelling population. *Arch Gerontol Geriatr*. 2012;54:28–33.
- Borges MP, Moreira AK. Influências da Prática de Atividades Físicas na Terceira Idade: Estudo comparativo dos níveis de autonomia para o desempenho nas AVDs e AIVDs entre idosos ativos fisicamente e idosos sedentários. *Motriz Rio Claro*. 2009;15:562–73.
- Ruhe A, Fejer R, Walker B. The test–retest reliability of centre of pressure measures in bipedal static task conditions – a systematic review of the literature. *Gait Posture*. 2010;32:436–45.

21. Smee DJ, Anson JM, Waddington GS, Berry HL. Association between physical functionality and falls risk in community-living older adults. *Curr Gerontol Geriatr Res*. 2012;2012:864516.
22. Antunes HKM, Santos RF, Cassilhas R, Santos RVT, Bueno OFA, de Mello MT. Exercício físico e função cognitiva: Uma Revisão. *Rev Bras Med Esporte*. 2006;12:97e–103e.
23. Carvalho LED [Tese de Pós-Graduação] Sinalização e Homeostase do Cálcio Hipocampal Moduladas Pelo Exercício Físico: Implicações nos processos de memória e neuroproteção. Minas Gerais: Universidade Federal de Minas Gerais; 2011.
24. Oliveira DLC, Goretti LC, Pereira LSM. O desempenho de idosos institucionalizados com alterações cognitivas em atividades de vida diária e mobilidade: Estudo Piloto. *Rev Bras Fisioter*. 2006;10:91–6.
25. Córdova C, Lopes-E-Silva F Jr, Pires AS, Souza VC, Brito CJ, Moraes CF, et al. Long-term resistance training is associated with reduced circulating levels of IL-6, IFN-gamma and TNF-alpha in elderly women. *Neuroimmunomodulation*. 2011;18:165–70.
26. Ferreira AP, Ferreira CB, Brito CJ, Souza VC, Córdova C, Nóbrega OT, et al. The effect of aerobic exercise intensity on attenuation of postprandial lipemia is dependent on apolipoprotein E genotype. *Atherosclerosis*. 2013;229:139–44.
27. Del Pozo-Cruz J, Magaña M, Ballesteros M, Porras M, Rodríguez Bies E, Navas P, et al. Influencia de la capacidad funcional sobre el perfil lipídico, daño muscular y perfil bioquímico en personas mayores no institucionalizadas. *Rev Andal Med Deporte*. 2013;6:57–65.