

Original

mir-199b-5p as a potential biomarker of bone fragility in type 2 diabetes mellitus – Dual role in musculoskeletal pathophysiology

María Dolores López Robles¹, Luis Martínez Heredia^{1,2}, Raquel Sanabria de la Torre^{1,3}, Francisco Luis Andújar Vera^{4,5,6}, Sheila González Salvatierra^{1,2}, Giulio Ferrero⁷, Alessio Naccarati⁷, Manuel Muñoz Torres^{1,2,8,9}, Beatriz García Fontana^{1,2,8}, Cristina García Fontana^{1,2,8}

¹Instituto de Investigación Biosanitaria de Granada. Granada, Spain. ²CIBER on Frailty and Healthy Aging (CIBERFES). Instituto de Salud Carlos III. Madrid, Spain. ³Department of Biochemistry and Molecular Biology III and Immunology. Universidad de Granada. Granada, Spain. ⁴Bioinformatic Unit. Instituto de Investigación Biosanitaria de Granada (Ibs. Granada). Granada, Spain. ⁵Department of Computer Science and Artificial Intelligence. Universidad de Granada. Granada, Spain. ⁶Andalusian Research Institute in Data Science and Computational Intelligence (DaSCI Institute). Granada, Spain. ⁷Italian Institute for Genomic Medicine (IIGM). Torino, Italy. ⁸Endocrinology and Nutrition Unit. Hospital Universitario Clínico San Cecilio. Granada, Spain. ⁹Department of Medicine. Universidad de Granada. Granada, Spain

Abstract

Type 2 diabetes mellitus (T2DM) is associated with increased risk of bone fragility and fractures, yet early diagnostic tools for musculoskeletal complications remain lacking. This study aimed to identify microRNAs (miRNAs) differentially expressed in T2DM patients and assess their potential as biomarkers for bone fragility. Serum samples from 8 T2DM patients and 8 matched healthy controls were analyzed using high-throughput sequencing and RT-qPCR. Among the miRNAs examined, hsa-miR-199b-5p was significantly under-expressed in T2DM patients, particularly in those with degraded trabecular bone score (TBS) and lower bone mineral density (BMD).

Statistical analyses revealed strong positive correlations between miR-199b-5p expression and indicators of bone integrity, including cortical and trabecular volumetric BMD, and serum periostin levels. Conversely, negative correlations were found with fracture risk (FRAX), TBS-adjusted FRAX, and CTX levels, supporting its role in bone metabolism. Literature suggests miR-199b-5p promotes osteogenesis via the GSK-3 β / β -catenin and periostin-mediated Wnt/ β -catenin signaling pathways. Thus, its reduced expression may contribute to impaired bone remodeling in T2DM. Interestingly, miR-199b-5p shows contrasting effects in osteoarthritis (OA), where it is upregulated and contributes to cartilage degradation. Periostin, similarly, promotes bone formation in T2DM but may exacerbate inflammation in OA. These findings underscore the context-dependent roles of miR-199b-5p and highlight its potential as a dual biomarker: protective in T2DM bone loss, yet detrimental in OA. Further research is needed to clarify its therapeutic relevance and ensure disease-specific targeting strategies.

Keywords:

Type 2 diabetes mellitus. microRNAs. Bone fragility. Osteoarthritis. Biomarkers.

Received: 06/04/2025 • Accepted: 07/23/2025

María Dolores López Robles and Luis Martínez Heredia contributed equally to this work.

Manuel Muñoz Torres and Beatriz García Fontana are the corresponding authors of this article.

Funding: This work was supported by the Junta de Andalucía (grant PI-0268-2019) and Instituto de Salud Carlos III (grant PI18-00803), co-funded by the European Regional Development Fund (FEDER) and CIBERFES (CB16/10/00475). C. G. F., and B. G. F. were funded by grants C1-S0033 and CP22/00022 from the Junta de Andalucía and Instituto de Salud Carlos III, respectively.

Conflict of interest: The authors declare no conflict of interest.

Artificial intelligence: The authors declare not to have used artificial intelligence (AI) or any AI-assisted technologies in the elaboration of the article

López Robles MD, Martínez Heredia L, Sanabria de la Torre R, Andújar Vera FL, González Salvatierra S, Ferrero G, Naccarati A, Muñoz Torres M, García Fontana B, García Fontana C. mir-199b-5p as a potential biomarker of bone fragility in type 2 diabetes mellitus – Dual role in musculoskeletal pathophysiology. Rev Osteoporos Metab Miner 2025;17(4):160-169

DOI: 10.20960/RevOsteoporosMetabMiner.00085

Correspondence:

Beatriz García Fontana. Endocrinology and Nutrition Unit. Hospital Regional Universitario de Málaga. Avda. de Carlos Haya, 84. 29010 Málaga, Spain
e-mail: bgfontana@fibao.es

INTRODUCTION

Diabetes *mellitus* is a leading cause of morbidity and mortality worldwide (1), with type 2 diabetes *mellitus* (T2DM) accounting for 90% % of all diabetes cases (2). The recent surge in the diagnosis rates of these and other lifestyle-related metabolic disorders can largely be attributed to the widespread adoption of sedentary lifestyles and excessive energy consumption in industrialized nations (3). Additional factors such as advanced age, hypertension, and family history further increase the predisposition to develop T2DM (4).

Currently, 13.8% % of the Spanish population suffers from T2DM, with an estimated 6% % of individuals unaware of their condition. Furthermore, approximately 12.6% % of the Spanish population is at risk of developing T2DM due to impaired glucose tolerance or abnormal fasting glucose levels (5). According to the di@bet.es study, 5.98 million Spaniards have issues with glucose metabolism (5). The International Diabetes Federation (IDF) projects that the number of diabetic patients globally will reach 643 million by 2030, making diabetes one of the “epidemics of the 21st century” (6,7).

In the early stages of T2DM, metabolic disturbances such as hyperglycemia, insulin resistance, dyslipidemia, and hyperinsulinemia contribute to cellular and organ-level damage, resulting in both microvascular complications (e. g, retinopathy, nephropathy, neuropathy) and macrovascular complications (e. g, coronary artery disease, cerebrovascular disease, peripheral artery disease). These vascular issues account for approximately 70–80% % of diabetes-related mortality (8,9). In addition to these vascular issues, growing evidence has highlighted that bone fragility is also prevalent in T2DM (10). The risk of fractures in patients with T2DM has significantly increased due to changes in bone remodeling and microarchitecture. Several studies have demonstrated a higher risk of fractures, particularly at the hip and vertebrae (11,12). This fragility is driven by chronic hyperglycemia, insulin resistance, and the accumulation of advanced glycation end products (AGEs), which impair bone quality and disrupt osteoblast-osteoclast balance (10). Indeed, multiple meta-analyses have confirmed an increased risk of incident hip, vertebral, and non-vertebral fractures in individuals with T2DM, with major consequences for quality of life (QoL) and long-term disability (13,14). On the other hand, recent studies have also drawn a connection between osteoarthritis (OA) and metabolic syndrome, a cluster including insulin resistance, dyslipidemia, and hypertension, further linking diabetes to musculoskeletal deterioration (15,16). The presence of diabetes is believed to accelerate the progression of OA and complicate its management, leading to the proposed diabetes-induced osteoarthritis (DM-OA) phenotype (17). This phenotype suggests that systemic inflammation and oxidative stress in diabetes predispose patients to OA, with chronic hyperglycemia promoting cartilage degradation, joint

inflammation, AGEs accumulation, and matrix stiffening, all of which impair joint cushioning and function (18). Recognizing musculoskeletal complications of type 2 diabetes *mellitus* as clinically significant outcomes, comparable with traditional vascular complications, underscores the urgent need for early detection strategies and targeted therapies to preserve bone and joint health in this growing patient population.

However, there is currently no tool available for the early diagnosis of these musculoskeletal complications. In this context, evaluating the differential expression of molecules such as microRNAs (miRNAs) may offer valuable insights into the development of diagnostic techniques for this population. The potential of miRNAs as biomarkers for musculoskeletal complications is supported by a growing body of scientific literature, which reports differential miRNA responses to various diseases (19,20), further highlighting the importance of investigating their role in T2DM-associated musculoskeletal deterioration.

In addition to miRNAs, proteins such as periostin have also emerged as important biomarkers in T2DM-related bone fragility. Periostin, a protein originally identified as osteoblast-specific factor-2, is involved in bone formation and remodeling but also plays a role in other complications of T2DM, such as cardiovascular disease (21). Th interaction of periostin with pathways such as Wnt/ β -catenin, which regulate osteoblast differentiation and bone metabolism, makes it a potential therapeutic target for improving bone health in diabetes (22). Its levels are altered in various musculoskeletal disorders, including those seen in T2DM, making it a promising biomarker for osteoporosis and bone fragility in these patients.

MicroRNAs (miRNAs) are small endogenous RNA molecules that post-transcriptionally regulate gene expression. They have been shown to play key roles in a wide range of biological processes. Recent advancements in transcriptomic technologies, especially next-generation sequencing and advanced bioinformatics tools, enable more in-depth exploration of messenger RNAs (mRNAs) and non-coding RNAs (ncRNAs), including miRNAs (23). Over 2,000 miRNAs have currently been identified, and it is estimated that they regulate approximately 30% % of all human genes. miRNAs are present in circulating blood, representing an opportunity to use these disease-related circulating miRNAs as potential biomarkers (24). In this study, the miRNome of a group of healthy individuals and patients with T2DM was analyzed to identify differentially expressed miRNAs between the 2 groups. The differential expression of selected miRNAs was subsequently validated using RT-qPCR. Following validation, the potential role of these miRNAs, particularly miR-199b-5p, as biomarkers in T2DM was further investigated through statistical analysis of their correlation with determinants of bone fragility and fracture risk.

MATERIALS AND METHODS

STUDY POPULATION

This study included a total of 16 participants, 8 T2DM patients (males with a mean age of 60 ± 5 years) and 8 control subjects, sex and aged matched. T2DM was diagnosed according to the American Diabetes Association criteria of 2017. T2DM patients were recruited during 2015 from the Endocrinology and Nutrition Unit of Hospital Universitario Clínico San Cecilio (Granada, Spain). Samples from healthy controls were managed and provided by the Andalusian Biobank.

Inclusion criteria for patients with T2DM were absence of cardiovascular disease, history of cardiovascular events, or renal, hepatic, GI, or thyroid disease. All patients were on diabetes drugs, such as metformin, sulfonylureas, insulin, or a combination of these drugs.

Venous blood samples were obtained at the Clinical Analysis Unit of Hospital Universitario Clínico San Cecilio.

The study was approved by the Ethics Committee of Jaén, Andalucía (ID 1630-M1-18/PI-0514-2018; December 20, 2018), and conducted in accordance with relevant ethical guidelines for human and animal research. Written informed consent was obtained from all participants.

CLINICAL EVALUATION, BIOCHEMICAL AND BONE PARAMETERS OF THE STUDY POPULATION

Anthropometric and biochemical measurements

Height, weight, and waist circumference were recorded following standard protocols. BMI was calculated as weight (kg) divided by height squared (m^2). Fasting venous blood samples were collected in the morning, with serum stored at $80^\circ C$ until analysis. Biochemical parameters measured included HbA1c, total cholesterol, HDL-c, LDL-c, triglycerides, creatinine, calcium, phosphorus, and vitamin D, using routine automated methods. Serum bioactive periostin was quantified in duplicate by ELISA (Biomedica Medizinprodukte GmbH), with detection ranges of 20-4,000 pmol/L and intra-/inter-assay variation $\leq 6\%$ and $\leq 3\%$, respectively. Average periostin levels in healthy individuals are ~ 864 pmol/L.

Hypertension and Dyslipidemia Assessment: Blood pressure was measured using an automated sphygmomanometer after 5 minutes of rest, with 2 readings 1-2 minutes apart; additional readings were taken if differences exceeded 10 mmHg. The mean of the last 2 readings was used. Hypertension was defined as sys-

tolic/diastolic $\geq 140/90$ mmHg or antihypertensive treatment. Dyslipidemia was defined by HDL-c < 50 mg/dL, LDL-c > 100 mg/dL, triglycerides > 150 mg/dL, and/or lipid-lowering drug use.

Areal bone mineral density (aBMD)

aBMD of the left hip was measured using a Hologic QDR 4500 densitometer. The shaft region was located 2 cm distal to the midpoint of the lesser trochanter along the shaft axis. Osteoporosis and osteopenia were categorized based on World Health Organization (WHO) criteria. Scans were performed by an experienced operator following ISCD guidelines; the device was calibrated daily with a spine phantom. Laboratory coefficients of variation were 1.8% (femoral neck) and 1.5% (total hip), with prior studies reporting 2.13% and 3.14% for shaft and trochanter, respectively.

Trabecular Bone Score (TBS)

TBS was measured at the LS using the latest version of TBS iNsight (version 3.0.2.0, Medimaps, Merignac, France). TBS was calculated as the mean value of the individual measurements for vertebrae L1-L4, based on a grey-level analysis of the DXA images. The TBS precision error (percentage of the coefficient of variation) was 1.82%. Diagnosis of preserved and degraded microarchitecture was based on the following TBS ranges: patients with TBS values ≥ 1.31 were categorized as preserved TBS and patients with TBS values < 1.31 were categorized as degraded TBS (25).

3D-DXA modeling

Trabecular macrostructure, cortical thickness, and femoral shape were assessed using 3D-Shaper software (v2.2), which generates participant-specific QCT-like models by registering a statistical shape and density model from QCT scans onto DXA images. Cortical thickness and density were derived from fitted mathematical functions along the femur surface. Measures included volumetric BMD (vBMD) of trabecular, cortical, and integral compartments, and cortical surface BMD (sBMD). Strong correlations with Quantitative Computed Tomography (QCT) ($r = 0.86-0.95$) and low coefficients of variation demonstrated high accuracy and precision.

Fracture Risk Assessment (FRAX)

FRAX estimates the 10-year probability of hip and major osteoporotic fractures by integrating clinical risk factors such as age, sex, BMI, fracture history, parental

hip fracture, smoking, glucocorticoid use, rheumatoid arthritis, secondary osteoporosis, alcohol intake adjusted by BMD and TBS values.

RNA EXTRACTION

miRNeasy Serum/Plasma kit (QIAGEN) was used to extract total RNA from 200 μ L of serum samples following the protocol in full compliance with the manufacturer's instructions for use. UniSp2, UniSp4 and UniSp5 sequences were added using RNA Spike-in mix (vial 1 of the miRNeasy Serum/Plasma kit) to be used as a control for this step. The extracted RNA was used to carry out the sequencing and validation steps.

SEQUENCING

Library development and sequencing

The extracted total RNA samples were sequenced through Illumina technology. miRNA libraries development was performed using the QIAseq miRNA Library Kit (QIAGEN). Adapters were ligated with molecular unique identifiers to subsequently back-transcribe the samples to cDNA, which was amplified by PCR. To guarantee the quality of the amplified samples, these were purified by capillary electrophoresis analysis (Alignent DNA 1000 Chip). Libraries were pooled equimolar and quantified by qPCR. Libraries were sequenced using a NextSeq sequencer (Illumina Inc). FASTAQ files of the sequences were created using bcl2fastq software.

Differential gene expression

Differential expression of miRNAs was analyzed using the Empirical Analysis of Differential Gene Expression (DGE) algorithm implemented in CLC Genomics Workbench v20.0.4. The NormFinder program, developed by Andersen et al. (26), was used to identify endogenous control miRNAs and to generate the volcano plot. Visualization of the volcano plot was performed using the ggplot2 and ggrepel packages in RStudio. For miRNAs selected based on the volcano plot criteria, Mann-Whitney tests were conducted on variance stabilizing transformation (VST)-normalized data, retaining those with p -values < 0.05.

VALIDATION

Polyadenylation and reverse transcription

miRNA molecules were reverse transcribed to cDNA with a poly(A) tail at the 3' end using the miRCURY

LNA RT Kit (Qiagen), according to the manufacturer's protocol. UniSp6 6 RNA spike-in was included as a control.

Quality control

RNA sequences added in the total RNA extraction step (UniSp2, UniSp4 and UniSp5) and in reverse transcription step (UniSp6) were amplified to make sure that the process had been conducted correctly. In addition, hemolysis-related sequences (has-miR-451a and has-miR-23a-3p) and endogenous miRNA sequences (hsa-miR-26a-5p and has-miR-30d-5p) were amplified to determine the quality of starting samples. All reactions were performed in triplicate. miRCURY LNA SYBR Green PCR kit was used to perform RT-qPCR following the protocol indicated by the manufacturer. According to the miRCURY LNA miRNA PCR assay protocol, the PCR conditions consisted of 2 steps: first, a denaturation step at 95 °C for 10 seconds, followed by a combined annealing/extension step at 56 °C for 60 seconds. The reaction occurred for a total of 40 cycles.

RT-qPCR

First, miRCURY LNA miRNA Custom PCR Panels were designed (Fig. 1). These plates were designed to amplify: 1) miRNAs identified in the sequencing; 2) 3 endogenous miRNAs; and 3) 2 control miRNAs (UniSp3 and UniSp6) in triplicate. miRCURY LNA SYBR Green PCR kit as well as in the quality control step was used.

miRNA expression analysis

The $\Delta\Delta C_t$ relative quantification method was used to evaluate the differential expression of miRNAs in the study groups. miRNA expression data are expressed as mean $\pm$ standard deviation (SD).

$$\begin{aligned}\Delta C_t (\text{patients}) &= C_t (\text{miRNA study}) - C_t (\text{miRNA problem}) \\ \Delta C_t (\text{controls}) &= C_t (\text{miRNA study}) - C_t (\text{miRNA problem}) \\ \Delta\Delta C_t &= \Delta C_t (\text{patients}) - \Delta C_t (\text{controls}) \\ FC &= 2^{-\Delta\Delta C_t}\end{aligned}$$

Statistical analysis

Student t tests were performed, with $p \leq .05$ considered statistically significant. Binary logistic regression was conducted using the Wald forward stepwise method. Receiver operating characteristic (ROC) curves were generated from the regression results. Statistical analyses were performed with SPSS, version 25.5 (IBM Corp).




Figura 1. miRCURY LNA miRNA Custom PCR Panels scheme.

RESULTS

STUDY POPULATION

Tables I and II illustrate the clinical, biochemical, and bone-related parameters of T2DM patients stratified by preserved vs degraded TBS. Patients with degraded TBS tended to have higher BMI and waist circumference and lower levels of total cholesterol, LDL, and HDL. Regarding bone parameters, this group showed lower values of BMD (both volumetric and areal), low levels of periostin and CTX, and higher PTHi concentrations, suggesting a trend toward impaired bone quality in individuals with degraded TBS.

DIFFERENTIAL EXPRESSION ANALYSIS AND VALIDATION OF SEQUENCING DATA BY RT-QPCR

Differential expression analysis was conducted to identify miRNAs with significant and biologically relevant changes between T2DM patients and healthy control. A volcano plot was generated to visualize differential expression, with cut-off thresholds set at a false discovery rate (FDR) < 0.05 and an absolute log₂ fold change [Log₂FC] > 1. Based on these criteria, hsa-miR-199b-5p and hsa-miR-491-5p were found to be significantly downregulated in T2DM, while hsa-miR-483-5p, hsa-miR-122-5p, hsa-miR-5010-5p, hsa-miR-193b-5p, and hsa-miR-320c were upregulated (Fig. 2A).

To confirm these findings, raw count data were normalized using the variance stabilizing transformation (VST), and a non-parametric hypothesis test was

Table I. Biochemical characteristics of T2DM patients

	Preserved TBS	Degraded TBS
Age (years)	61.75 ± 2.75	60.50 ± 2.08
Weigh (kg)	85.83 ± 10.74	98.33 ± 14.87
Height(cm)	169.75 ± 9.00	170.25 ± 10.34
BMI (kg/m ²)	29.72 ± 2.28	33.96 ± 4.54
Waist (cm)	102.25 ± 6.59	112.25 ± 10.90
HbA1c (%)	7.95 ± 1.42	7.68 ± 1.89
Uric acid (mg/dL)	5.98 ± 1.54	5.93 ± 0.97
Cholesterol (mg/dL)	215 ± 29.92	148 ± 35.42
LDL (mg/dL)	158 ± 29.07	80.25 ± 30.89
HDL (mg/dL)	53.75 ± 9.64	45.50 ± 11.82
TGs (mg/dL)	207 ± 123.94	113 ± 38.03
Creatinine (mg/dL)	0.72 ± 0.12	0.84 ± 0.1
AHT (%)	50 %	50 %
Dyslipidemia (%)	100 %	75 %

BMI: body mass index; HbA1c: glycated hemoglobin; LDL: low-density lipoprotein; HDL: high-density lipoprotein; TGs: triglycerides; AHT: arterial hypertension; T2DM: type 2 diabetes mellitus; TBS: trabecular bone score.

applied to compare miRNA expression levels across groups. Significant differences were observed for all selected miRNAs except hsa-miR-193b-5p (Fig. 2B).

Table II. Bone and fracture risk parameters in T2DM patients

	Preserved TBS	Degraded TBS
TBS	1.28 ± 0.05	1.02 ± 0.10
Hip FRAX	0.30 ± 0.18	0.20 ± 0.22
Hip FRAX_TBS	0.38 ± 0.28	0.5 ± 0.68
Trabecular_vBMD (g/cm ³)	193.29 ± 40.75	170.15 ± 91.13
Cortical_vBMD (g/cm ³)	869.13 ± 72.35	860.24 ± 84.04
Cortical_sBMD (g/cm ³)	186.42 ± 35.52	174.69 ± 21.79
Integral_vBMD (g/cm ³)	339.17 ± 67.44	315.58 ± 107.29
TH_BMD (g/cm ²)	1.205 ± 0.17	1.11 ± 0.25
Periostin (pmol/L)	1799.99 ± 346.31	1339.66 ± 245.23
iPTH (pg/mL)	25.35 ± 1.34	33.83 ± 9.77
VitD (ng/mL)	20.63 ± 5.98	19.08 ± 6.69
Calcemia (mg/dL)	9.98 ± 0.26	9.85 ± 0.24
Phosphorus (mg/dL)	3.20 ± 0.22	3.53 ± 0.50
CTX (ng/L)	0.46 ± 0.31	0.26 ± 0.12

FRAX: BMD-adjusted (with or without TBS) hip Fracture Risk Assessment Tool; vBMD: volumetric bone mineral density; sBMD: subchondral bone mineral density; TH_BMD: total hip bone mineral density; iPTH: intact parathyroid hormone; CTX: C-terminal telopeptide of type I collagen.

These miRNAs were subsequently selected for validation by RT-qPCR. Among them, only hsa-miR-199b-5p was successfully validated. Consistent with sequencing data, hsa-miR-199b-5p was found to be significantly downregulated in T2DM patients vs healthy controls (0.496 ± 0.043 -fold, $p < 0.05$).

CORRELATION BETWEEN HSA-MIR-199B-5P EXPRESSION AND DETERMINANTS OF BONE FRAGILITY AND INCREASED RISK OF FRACTURE IN T2DM PATIENTS (Fig. 3)

Within the T2DM group, the expression of hsa-miR-199b-5p was correlated with several clinical markers of bone fragility and fracture risk. Spearman correlation analysis revealed significant negative correlations between hsa-miR-199b-5p expression and A) FRAX hip fracture risk ($p = -0.74$, 95% %CI, {-0.95, -0.05}, $p = 0.036$); B) FRAX hip fracture risk (TBS adjust) ($p = -0.862$, 95% %CI, {-0.98, -0.38}, $p = 0.005$); C) CTX (pg/ml) ($p = -0.719$, 95% %CI, {-0.95, 0}, $p = 0.0446$).

In contrast, positive correlations were observed between hsa-miR-199b-5p expression and D) serum levels of periostin (pmol/L) ($p = 0.786$, 95% %CI, {0.16, 0.96}, $p = 0.0208$); E) Cortical vBMD (g/cm³) ($p = 0.893$,




Figure 2. Differential expression of miRNAs. A. Volcano plot showing Log2 fold change (x-axis) vs $-\log_{10} p$ -value (y-axis). Dashed lines indicate cut-offs for significance ($FDR < 0.05$) and expression change ($|\log_2 FC| > 1$). Labeled miRNAs passed both thresholds. B. Violin plots of selected miRNAs showing expression levels after variance stabilizing transformation (VST).




Figure 3. Scatter plots showing the correlation between hsa-miR-199b-5p expression and FRAX hip fracture risk (A); FRAX hip fracture risk (TBS adjust) (B); CTX (pg/mL) (C); serum levels of periostin (pmol/L) (D); Cortical vBMD (g/cm³) (E); Cortical sBMD (g/cm³) (F); Trabecular vBMD (g/cm³) (G); Integral vBMD (g/cm³) (H), in T2DM patients ($n = 8$). The p -values between the different associations were obtained via Spearman's correlation coefficients (showing $p < 0.05$ in each scatter plot).

95% %CI, {0.4, 0.99}, $p = 0.0068$); F) Cortical sBMD (g/cm³) ($p = 0.857$, 95% %CI, {0.27, 0.98}, $p = 0.0137$); G) Trabecular vBMD (g/cm³) ($p = 0.893$, 95% %CI, {0.4, 0.99}, $p = 0.0068$), E) Integral vBMD (g/cm³) ($p = 0.893$, 95% %CI, {0.4, 0.99}, $p = 0.0068$).

DISCUSSION

Our results obtained from sequencing and validation steps suggest differential expression of hsa-miR-199b-5p in T2DM patients vs the healthy group. With the aim of exploring the potential of miR-199b-5p as a novel therapeutic target for the prevention and management of bone fragility associated with T2DM, we conducted multiple correlation analyses in T2DM patients. Our work reveals a positive correlation between miR-199b-5p expression and bone mineral density parameters, including cortical vBMD (mg/cm³), cortical sBMD (mg/cm³), trabecular vBMD (mg/cm³), and integral vBMD (mg/cm³). Consistently, we also observed a negative correlation between hsa-miR-199b-5p expression and FRAX hip fracture risk, FRAX hip fracture risk adjusted for trabecular bone score (TBS), and serum C-terminal telopeptide of type I collagen (CTX) levels (pg/mL), all of which are established indicators of demineralization and bone fragility. These clinical findings are in line with emerging molecular evidence supporting the involvement of miRNA, particularly miR-199b-5p, in osteogenesis and bone metabolism. It has been demonstrated that primary mesenchymal stem cells (MSCs) secrete small RNAs via exosomes, which are increasingly recognized for their role in in-

tercellular communication (27). Bioinformatics analysis using DIANA-mirPath revealed that the expression of exosomal miR-199b-5p, among other miRNAs, is impaired during osteogenic differentiation, suggesting that exosomal miR-199b-5p may play a key role in regulating osteoblast differentiation (28). The effects of knockdown and overexpression of miR-199b-5p on osteoblast differentiation have been studied by measuring alkaline phosphatase (ALP) expression and activity, as well as the expression of the osteogenic marker gene Runx2 (29). The findings indicated that miR-199b-5p enhanced the osteogenic potential of bone marrow-derived MSCs (BMSCs) by modulating the GSK-3 β / β -catenin signaling pathway, suggesting that miR-199b-5p and its analogs could serve as promising therapeutic candidates for bone and musculoskeletal disorders. Furthermore, the screening and validation of miRNAs associated with osteoporosis, using high-throughput sequencing, revealed a reduced expression of miR-199b-5p (20). Based on all this evidence, our results suggest that reduced expression of miR-199b-5p in T2DM patients may disrupt the regulation of osteogenesis, potentially resulting in diminished bone mineralization and increased skeletal fragility.

Additionally, our results also demonstrated a positive correlation between miR-199b-5p expression and serum periostin levels (pmol/L). Periostin is a ubiquitous protein originally known as osteoblast-specific factor-2 and belongs to a group of nonstructural extracellular matrix (ECM) proteins. This protein not only plays a role in adverse cardiac remodeling, being associated with cardiovascular disease in T2DM according to the SCORE2-Diabetes algorithm (21), but also

promotes bone formation, regeneration, and repair (22). An experimental study on bone loss prevention demonstrated that periostin-induced downregulation of sclerostin can activate the Wnt/ β -catenin signaling pathway, thereby inhibiting bone loss (30). This mechanism likely involves the interaction of Wnt proteins with the FZD and LRP6 receptors, leading to the inactivation of the cytoplasmic GSK-3 β /Axin2/APC complex. As a result, β -catenin accumulates in the cytoplasm and translocates into the nucleus, where it promotes the transcription of target genes involved in bone formation, ultimately facilitating osteoblast differentiation (31). In this line, our findings suggest that reduced expression of miR-199b-5p in T2DM patients may contribute to impaired osteogenesis through its regulatory effect on periostin. The observed positive correlation between miR-199b-5p expression and serum periostin levels supports the hypothesis that miR-199b-5p may modulate bone metabolism, at least in part, via the periostin–Wnt/ β -catenin signaling pathway.

Overall, our results suggest that miR-199b (with osteoprotective role in T2DM) could serve as a predictive biomarker, potentially enabling the early identification and monitoring of osteoporosis progression associated with T2DM.

Interestingly, while our findings support a protective role for miR-199b-5p in bone metabolism and osteogenesis, particularly in the context of T2DM-related osteoporosis, emerging evidence suggests that this microRNA may exert deleterious effects in osteoarthritis (OA). Multiple studies have highlighted the role of miR-199b-5p in chondrogenesis. A longitudinal bioinformatics analysis identified miR-199b-5p as a key pro-chondrogenic regulator (32). Experimental data from the same study showed that inhibition of miR-199b-5p during the early stages of chondrogenesis led to downregulation of chondrogenic markers. A deeper understanding of the regulatory mechanisms involved in chondrogenesis may offer valuable insights into OA pathogenesis. The mechanisms of differentiation differ between growth plate cartilage and articular cartilage. While the growth plate drives longitudinal bone growth via cartilage-to-bone substitution, articular cartilage maintenance depends on delayed chondrocyte maturation and hypertrophy (33). Therefore, it is plausible that, in OA, the cartilage maturation process becomes aberrantly activated, leading to a loss of biomechanical integrity. In this context, serum exosomal small RNA sequencing from clinical OA patients, along with gene expression data from serum and cartilage samples retrieved from the GEO database, revealed that miR-199b-5p is consistently upregulated (19). In vitro studies further demonstrated that miR-199b-5p negatively affects chondrocyte viability and promotes extracellular matrix degradation. Conversely, inhibition of miR-199b-5p under inflammatory conditions exerted protective effects against tissue damage. Additionally, in an OA model, blocking miR-199b-5p, al-

leviated disease progression, suggesting that, in contrast to its osteoprotective role in T2DM, miR-199b-5p may contribute to cartilage degeneration in OA. Of note, none of the patients included in our study had a diagnosis of osteoarthritis (OA), supporting the interpretation that the observed downregulation of miR-199b-5p is specifically linked to diabetic bone fragility rather than joint-related pathology.

Similarly, periostin, which was positively correlated with miR-199b-5p expression in our T2DM cohort, appears overexpressed in OA cartilage (34). Immunohistochemical analysis localized periostin to chondrocytes and their surrounding matrix, particularly near erosive regions. In vitro, periostin stimulation of chondrocytes induced the upregulation of MMPs, IL-6, and IL-8, supporting its involvement in OA-associated inflammatory responses.

Overall, these findings underscore the context-dependent nature of miR-199b-5p activity. While it may confer protection vs bone fragility in T2DM by promoting osteogenesis through periostin-mediated Wnt/ β -catenin signaling, its upregulation in joint tissues under inflammatory conditions may exacerbate OA disease. Therefore, the development of miR-199b-5p-based therapies requires a nuanced understanding of its distinct effects in bone versus cartilage to ensure the safety and efficacy profile.

Our study has several strengths and limitations. One of the main limitations is its cross-sectional design, which does not allow us to establish causal relationships between miR-199b-5p expression and bone fragility parameters in T2DM. While our findings are consistent with the biological role of this microRNA in osteogenesis, longitudinal studies are required to determine if its downregulation precedes bone loss or results from it. In addition, although the sample size of our population was small, it is acceptable for a pilot study exploring novel molecular markers in a well-defined clinical context. However, caution is advised when extrapolating these findings to broader populations. Moreover, the study cohort included only male participants, which limits the assessment of potential sex-specific differences, so further studies including both, larger populations and women, especially postmenopausal who are at increased risk of osteoporosis, are needed to confirm these results and to enhance generalizability.

Despite these limitations, our study has notable strengths. As far as we know, this is the first study linking reduced serum miR-199b-5p expression to diabetic bone fragility offering novel insights into its potential role as a biomarker. The associations were robust and supported by correlations with multiple clinical parameters, including BMD, TBS-adjusted FRAX, serum periostin and CTX levels. Additionally, the study excluded patients with osteoarthritis, reducing the risk

of confounding from inflammatory joint disease and reinforcing the specificity of the observed associations with bone metabolism. The integration of bioinformatic analysis and molecular validation strengthens the biological plausibility of our findings.

While our study highlights the potential of miR-199b-5p as a biomarker for bone fragility in T2DM, we acknowledge that the use of circulating microRNAs in clinical practice remains limited. Nevertheless, significant progress has been made in recent years toward the development of robust and standardized assays, and circulating miRNAs are increasingly recognized as promising candidates for non-invasive diagnostics. In this context, our findings contribute to the growing foundation for the future incorporation of miRNA profiling into clinical decision-making as a complementary tool for fracture risk assessment.

CONCLUSIONS

This study identifies hsa-miR-199b-5p as a promising biomarker for bone fragility in T2DM patients. Its reduced expression is significantly associated with lower bone mineral density and increased fracture risk, suggesting a potential role in impaired osteogenesis. Furthermore, the strong correlation between miR-199b-5p and periostin levels highlights its involvement in the Wnt/ β -catenin signaling pathway. However, its contrasting role in OA underscores the need for context-specific interpretation. These findings support the potential of miR-199b-5p as both a diagnostic tool and a therapeutic target for skeletal complications in T2DM.

REFERENCES

- Bailes BK. Diabetes mellitus and its chronic complications. *AORN J* 2002;76(2):266-76, 278-82; quiz 283-6. DOI: 10.1016/S0001-2092(06)61066-1
- Guthrie RA, Guthrie DW. Pathophysiology of diabetes mellitus. *Crit Care Nurs Q* 2004;27(2):113-25. DOI: 10.1097/00002727-200404000-00003
- Booth FW, Roberts CK. Linking performance and chronic disease risk: indices of physical performance are surrogates for health. *Br J Sports Med* 2008;42(12):950-2. DOI: 10.1136/bjsm.2008.052589
- Fletcher B, Gulanick M, Lamendola C. Risk factors for type 2 diabetes mellitus. *J Cardiovasc Nurs* 2002;16(2):17-23. DOI: 10.1097/00005082-200201000-00003
- Soriguer F, Goday A, Bosch-Comas A, Bordiú E, Calle-Pascual A, Carmena R, et al. Prevalence of diabetes mellitus and impaired glucose regulation in Spain: the Di@bet.es Study. *Diabetologia* 2012;55(1):88-93. DOI: 10.1007/s00125-011-2336-9
- Magliano DJ, Boyko EJ, IDF Diabetes Atlas 10th edition scientific committee. IDF DIABETES ATLAS [Internet]. 10th ed. Brussels: International Diabetes Federation; 2021 [citado 21 de mayo de 2025]. (IDF Diabetes Atlas). Available at: <http://www.ncbi.nlm.nih.gov/books/NBK581934/>
- Standl E, Khunti K, Hansen TB, Schnell O. The global epidemics of diabetes in the 21st century: Current situation and perspectives. *Eur J Prev Cardiol* 2019;26(2_suppl):7-14. DOI: 10.1177/2047487319881021
- Sanabria-de la Torre R, García-Fontana C, González-Salvatierra S, Andújar-Vera F, Martínez-Heredia L, García-Fontana B, et al. The Contribution of Wnt Signaling to Vascular Complications in Type 2 Diabetes Mellitus. *Int J Mol Sci* 2022;23(13):6995. DOI: 10.3390/ijms23136995
- García-Fontana B, Morales-Santana S, Díaz Navarro C, Rozas-Moreno P, Genilloud O, Vicente Pérez F, et al. Metabolomic profile related to cardiovascular disease in patients with type 2 diabetes mellitus: A pilot study. *Talanta* 2016;148:135-43. DOI: 10.1016/j.talanta.2015.10.070
- Khosla S, Samakkarthai P, Monroe DG, Farr JN. Update on the pathogenesis and treatment of skeletal fragility in type 2 diabetes mellitus. *Nat Rev Endocrinol* 2021;17(11):685-97.
- Schousboe JT, Morin SN, Kline GA, Lix LM, Leslie WD. Differential risk of fracture attributable to type 2 diabetes mellitus according to skeletal site. *Bone* 2022;154:116220. DOI: 10.1016/j.bone.2021.116220
- Schwartz AV. Epidemiology of fractures in type 2 diabetes. *Bone* 2016;82:2-8. DOI: 10.1016/j.bone.2015.05.032
- Koromani F, Oei L, Shevroja E, Trajanoska K, Schoufour J, Muka T, et al. Vertebral Fractures in Individuals With Type 2 Diabetes: More Than Skeletal Complications Alone. *Diabetes Care* 2020;43(1):137-44. DOI: 10.2337/dc19-0925
- Vilaca T, Schini M, Harnan S, Sutton A, Poku E, Allen IE, et al. The risk of hip and non-vertebral fractures in type 1 and type 2 diabetes: A systematic review and meta-analysis update. *Bone* 2020;137:115457. DOI: 10.1016/j.bone.2020.115457
- Seow SR, Mat S, Ahmad Azam A, Rajab NF, Safinar Ismail I, Singh DKA, et al. Impact of diabetes mellitus on osteoarthritis: a scoping review on biomarkers. *Expert Rev Mol Med* 2024;26:e8. DOI: 10.1016/j.bone.2015.05.032
- Courties A, Berenbaum F, Sellam J. The Phenotypic Approach to Osteoarthritis: A Look at Metabolic Syndrome-Associated Osteoarthritis. *Joint Bone Spine* 2019;86(6):725-30. DOI: 10.1016/j.jbspin.2018.12.005
- Courties A, Sellam J. Osteoarthritis and type 2 diabetes mellitus: What are the links? *Diabetes Res Clin Pract* 2016;122:198-206. DOI: 10.1016/j.diabres.2016.10.021
- Verzijl N, Bank RA, TeKoppele JM, DeGroot J. AGEing and osteoarthritis: a different perspective. *Curr Opin Rheumatol* 2003;15(5):616-22. DOI: 10.1097/00002281-200309000-00016
- Feng T, Zhang Q, Li SH, Ping YL, Tian MQ, Zhou SH, et al. Inhibition of miR-199b-5p reduces pathological alterations in osteoarthritis by potentially targeting Fzd6 and Gcnt2. *Elife* 2024;12:RP92645. DOI: 10.7554/eLife.92645
- Huang J, Yang H, Chai S, Lin Y, Zhang Z, Huang H, et al. Identification of miRNAs related to osteoporosis by high-throughput sequencing. *Front Pharmacol* 2024;15:1451695. DOI: 10.3389/fphar.2024.1451695

21. González-Salvatierra S, García-Martín A, García-Fontana B, Martínez-Heredia L, García-Fontana C, Muñoz-Torres M. Bone proteins are associated with cardiovascular risk according to the SCORE2-Diabetes algorithm. *Cardiovasc Diabetol* 2024;23(1):311. DOI: 10.1186/s12933-024-02406-9
22. Yuan C, Li J. Research progress of periostin and osteoporosis. *Front Endocrinol (Lausanne)* 2024;15:1356297. DOI: 10.3389/fendo.2024.1356297
23. Do DN, Dudemaine PL, Mathur M, Suravajhala P, Zhao X, Ibeagha-Awemu EM. miRNA Regulatory Functions in Farm Animal Diseases, and Biomarker Potentials for Effective Therapies. *Int J Mol Sci* 2021;22(6):3080. DOI: 10.3390/ijms22063080
24. Songia P, Chiesa M, Valerio V, Moschetta D, Myasoedova VA, D'Alessandra Y, et al. Direct screening of plasma circulating microRNAs. *RNA Biol* 2018;15(10):1268-72. DOI: 10.1080/15476286.2018.1526538
25. McCloskey EV, Odén A, Harvey NC, Leslie WD, Hans D, Johansson H, et al. A Meta-Analysis of Trabecular Bone Score in Fracture Risk Prediction and Its Relationship to FRAX. *J Bone Miner Res* 2016;31(5):940-8. DOI: 10.1002/jbmr.2734
26. Andersen CL, Jensen JL, Ørntoft TF. Normalization of real-time quantitative reverse transcription-PCR data: a model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Cancer Res* 2004;64(15):5245-50. DOI: 10.1158/0008-5472.CAN-04-0496
27. Baglio SR, Rooijers K, Koppers-Lalic D, Verweij FJ, Pérez Lanzón M, Zini N, et al. Human bone marrow- and adipose-mesenchymal stem cells secrete exosomes enriched in distinctive miRNA and tRNA species. *Stem Cell Res Ther* 2015;6(1):127. DOI: 10.1186/s13287-015-0116-z
28. Xu JF, Yang GH, Pan XH, Zhang SJ, Zhao C, Qiu BS, et al. Altered microRNA expression profile in exosomes during osteogenic differentiation of human bone marrow-derived mesenchymal stem cells. *PLoS One* 2014;9(12):e114627. DOI: 10.1371/journal.pone.0114627
29. Zhao R, Li Y, Lin Z, Wan J, Xu C, Zeng Y, et al. miR-199b-5p modulates BMSC osteogenesis via suppressing GSK-3 β / β -catenin signaling pathway. *Biochem Biophys Res Commun* 2016;477(4):749-54. DOI: 10.1016/j.bbrc.2016.06.130
30. Lv J, Sun X, Ma J, Ma X, Xing G, Wang Y, et al. Involvement of periostin-sclerostin-Wnt/ β -catenin signaling pathway in the prevention of neurectomy-induced bone loss by naringin. *Biochem Biophys Res Commun* 2015;468(4):587-93. DOI: 10.1016/j.bbrc.2015.10.152
31. Robinson JA, Chatterjee-Kishore M, Yaworsky PJ, Cullen DM, Zhao W, Li C, et al. Wnt/ β -catenin signaling is a normal physiological response to mechanical loading in bone. *J Biol Chem* 2006;281(42):31720-8. DOI: 10.1074/jbc.M602308200
32. Patel K, Barter M, Soul J, Clark P, Proctor C, Clark I, et al. Systems analysis of miR-199a/b-5p and multiple miR-199a/b-5p targets during chondrogenesis. *Elife* 2024;12:RP89701. DOI: 10.7554/eLife.89701.4
33. Velasquillo C, Garciadiego D, Ibarra C. Joint cartilage differentiation and osteoarthritis. *Reumatol Clin* 2007;3 Suppl 3:S2-5. DOI: 10.1016/S1699-258X(07)73647-1
34. Chijimatsu R, Kunugiza Y, Taniyama Y, Nakamura N, Tomita T, Yoshikawa H. Expression and pathological effects of periostin in human osteoarthritis cartilage. *BMC Musculoskelet Disord* 2015;16:215. DOI: 10.1186/s12891-015-0682-3